

Scattering Theory

Ting-Kai Hsu

January 29, 2024

Contents

1 Potential Scattering	1
1.1 In-States	1

This article is designed for leading the reader from old scattering theory of quantum mechanics to nowadays scattering theory of quantum field theory.

1 Potential Scattering

In this chapter we would study the theory of scattering in a simple but important case, the elastic scattering of a non-relativistic particle in a local potential, but using modern techniques that could easily be extended to more general problems.

1.1 In-States

Consider a non-relativistic particle of mass μ in a potential $V(\mathbf{x})$, whose Hamiltonian is,

$$H = H_0 + V(\mathbf{x}) \quad (1.1)$$

Here $H_0 = \mathbf{p}^2/2\mu$ is the kinetic energy operator. The potential is assumed to be function of position operator, and tends to zero as $|\mathbf{x}| = r \rightarrow \infty$. Then it is concerned here with a positive-energy particle¹. The particle comes into facing the potential from great distances regarded to have no interaction with momentum $\hbar\mathbf{k}$,

¹Actually, there could possibly be negative energy if it is in bound state, that is, the energy is less than potential at boundaries. But now, we are considering scattering state, which is opposite with bound state.

and is scattered, going out again to infinity, generally along a different direction.

In Heisenberg picture, for a particle with momentum $\hbar\mathbf{k}$ far from the scattering center if the measurement of the particle is made at very early times, and this situation is represented by a time-independent state vector $|\Psi_{\mathbf{k}}^{\text{in}}\rangle$. As mentioned, at very early times the particle is at a location that is far from the scattering center so that the potential is negligible there, so the state has an energy $\hbar^2\mathbf{k}^2/2\mu$, with the following relation,

$$H |\Psi_{\mathbf{k}}^{\text{in}}\rangle = \frac{\hbar^2\mathbf{k}^2}{2\mu} |\Psi_{\mathbf{k}}^{\text{in}}\rangle \quad (1.2)$$

Now switch to Schrödinger picture, the state would evolve as time goes on. As mentioned, the scattering state would be continuous *superposition*² of states with a spread of energies,

$$|\Psi_g(t)\rangle = \int d^3k g(\mathbf{k}) \exp\left(-i\frac{\hbar t\mathbf{k}^2}{2\mu}\right) |\Psi_{\mathbf{k}}^{\text{in}}\rangle \quad (1.3)$$

where $g(\mathbf{k})$ is a smooth function that is peaked around some wave number \mathbf{k}_0 . The eigenstate $|\Psi_{\mathbf{k}}^{\text{in}}\rangle$ satisfies the further condition that for any sufficiently smooth function $g(\mathbf{k})$, in the limit $t \rightarrow -\infty$,

$$|\Psi_g(t)\rangle \rightarrow \int d^3k g(\mathbf{k}) \exp\left(-i\frac{\hbar t\mathbf{k}^2}{2\mu}\right) |\Phi_{\mathbf{k}}\rangle \quad (1.4)$$

where $|\Phi_{\mathbf{k}}\rangle$ are orthonormal eigenstates of the momentum operator \mathbf{P} with eigenvalue $\hbar\mathbf{k}$,

$$\begin{aligned} \mathbf{P} |\Phi_{\mathbf{k}}\rangle &= \hbar\mathbf{k} |\Phi_{\mathbf{k}}\rangle \\ \langle \Phi_{\mathbf{k}} | \Phi_{\mathbf{k}'} \rangle &= \delta^{(3)}(\hbar\mathbf{k} - \hbar\mathbf{k}') \end{aligned} \quad (1.5)$$

and hence eigenstates of H_0 (not H), with eigenvalue $E(\mathbf{k}) = \hbar^2\mathbf{k}^2/2\mu$. Note that $|\Psi_{\mathbf{k}}^{\text{in}}\rangle$ and $|\Phi_{\mathbf{k}}\rangle$ belong to different Hilbert spaces. The

²It would be discrete sum for a bound state.