


# Our Status in AI


---

- We're done with Part I Searching
- Part II: Probabilistic Reasoning
  - Probability
  - Bayesian networks and reasoning
- Part III: Machine Learning



# Artificial Intelligence

## Probability




Instructors: Dan Klein and Pieter Abbeel --- University of California, Berkeley

[These slides were created by Dan Klein and Pieter Abbeel for CS188 Intro to AI at UC Berkeley. All CS188 materials are available at <http://ai.berkeley.edu>.]

# Today

---

- Probability
  - Random Variables
  - Joint and Marginal Distributions
  - Conditional Distribution
  - Product Rule, Chain Rule, Bayes' Rule
  - Inference
  - Independence
- You'll need all this stuff A LOT for the next few weeks, so make sure you go over it now!





# Uncertainty

- Why uncertainty?
  - Partial observability,
  - Nondeterminism
- Intelligent agents must be able to deal with uncertainty.

Autonomated car:




Maybe:

- Lack of observability
- Or, braked but error of sensors

# Uncertainty

- A ghost is in the grid somewhere
- Sensor readings tell how close a square is to the ghost
  - On the ghost: red
  - 1 or 2 away: orange
  - 3 or 4 away: yellow
  - 5+ away: green
- Sensors are noisy, but we know  $P(\text{Color} \mid \text{Distance})$



$P(\text{red} \mid 3)$	$P(\text{orange} \mid 3)$	$P(\text{yellow} \mid 3)$	$P(\text{green} \mid 3)$
0.05	0.15	0.5	0.3

# Uncertainty

- General situation:
  - **Observed variables (evidence):** Agent knows certain things about the state of the world (e.g., sensor readings or symptoms)
  - **Unobserved variables:** Agent needs to reason about other aspects (e.g. where an object is or what disease is present)
  - **Model:** Agent knows something about how the known variables relate to the unknown variables
- Probabilistic reasoning gives us a framework for managing our beliefs and knowledge


0.11	0.11	0.11
0.11	0.11	0.11
0.11	0.11	0.11

0.17	0.10	0.10
0.09	0.17	0.10
<0.01	0.09	0.17

<0.01	<0.01	0.03
<0.01	0.05	0.05
<0.01	0.05	0.81

# Random Variables


- A random variable is some aspect of the world about which we (may) have uncertainty
  - R = Is it raining?
  - T = Is it hot or cold?
  - D = How long will it take to drive to work?
  - L = Where is the ghost?
- We denote random variables with capital letters
- Like variables in a CSP, random variables have domains
  - R in {true, false} (often write as {+r, -r})
  - T in {hot, cold}
  - D in  $[0, \infty)$
  - L in possible locations, maybe  $\{(0,0), (0,1), \dots\}$



# Probability Distributions

- Associate a probability with each value


- Temperature:



$P(T)$

T	P
hot	0.5
cold	0.5

- Weather:



$P(W)$

W	P
sun	0.6
rain	0.1
fog	0.3
meteor	0.0

# Probability Distributions

- Unobserved random variables have distributions

T	P
hot	0.5
cold	0.5

W	P
sun	0.6
rain	0.1
fog	0.3
meteor	0.0

- A distribution is a TABLE of probabilities of values
- A probability (lower case value) is a single number

$$P(W = \text{rain}) = 0.1$$

- Must have:  $\forall x \ P(X = x) \geq 0$  and  $\sum_x P(X = x) = 1$

Shorthand notation:

$$P(\text{hot}) = P(T = \text{hot}),$$

$$P(\text{cold}) = P(T = \text{cold}),$$

$$P(\text{rain}) = P(W = \text{rain}),$$

...

OK if all domain entries are unique

# Joint Distributions

- A *joint distribution* over a set of random variables:  $X_1, X_2, \dots, X_n$  specifies a real number for each assignment (or *outcome*):

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

$$P(x_1, x_2, \dots, x_n)$$

- Must obey:  $P(x_1, x_2, \dots, x_n) \geq 0$

$$\sum_{(x_1, x_2, \dots, x_n)} P(x_1, x_2, \dots, x_n) = 1$$

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3


- Size of distribution if  $n$  variables with domain sizes  $d$ ?
  - For all but the smallest distributions, impractical to write out!

# Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
  - (Random) variables with domains
  - Assignments are called *outcomes*
  - Joint distributions: say whether assignments (outcomes) are likely
  - *Normalized*: sum to 1.0
  - Ideally: only certain variables directly interact
- Constraint satisfaction problems:
  - Variables with domains
  - Constraints: state whether assignments are possible
  - Ideally: only certain variables directly interact


Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3



Constraint over T,W

T	W	P
hot	sun	T
hot	rain	F
cold	sun	F
cold	rain	T



# Events

- An event is a set E of outcomes

$$P(E) = \sum_{(x_1 \dots x_n) \in E} P(x_1 \dots x_n)$$

- From a joint distribution, we can calculate the probability of any event

- Probability that it's hot AND sunny?

- Probability that it's hot?

- Probability that it's hot OR sunny?

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

For any proposition  $\phi$ ,  $P(\phi) = \sum_{\omega \in \phi} P(\omega)$

$$P(a \vee b) = P(a) + P(b) - P(a \wedge b)$$

- Typically, the events we care about are *partial assignments*, like  $P(T=\text{hot})$

# Quiz: Events

- $P(+x, +y) ?$

$$P(X, Y)$$

- $P(+x) ?$

- $P(-y \text{ OR } +x) ?$

X	Y	P
+x	+y	0.2
+x	-y	0.3
-x	+y	0.4
-x	-y	0.1

# Marginal Distributions

- Marginal distributions are sub-tables which eliminate variables
- Marginalization (summing out): Combine collapsed rows by adding

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3



$$P(t) = \sum_s P(t, s)$$




$$P(s) = \sum_t P(t, s)$$

$$P(X_1 = x_1) = \sum_{x_2} P(X_1 = x_1, X_2 = x_2)$$

T	P
hot	0.5
cold	0.5

$P(W)$

W	P
sun	0.6
rain	0.4



# Quiz: Marginal Distributions

$P(X, Y)$

X	Y	P
+x	+y	0.2
+x	-y	0.3
-x	+y	0.4
-x	-y	0.1

$$P(x) = \sum_y P(x, y)$$


$$P(y) = \sum_x P(x, y)$$

$P(X)$

X	P
+x	
-x	

$P(Y)$

Y	P
+y	
-y	




# Conditional Probabilities

- A simple relation between joint and conditional probabilities
  - In fact, this is taken as the *definition* of a conditional probability

$$P(a|b) = \frac{P(a,b)}{P(b)}$$

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3



$$P(W = s | T = c) = \frac{P(W = s, T = c)}{P(T = c)} = \frac{0.2}{0.5} = 0.4$$

$$\begin{aligned} &= P(W = s, T = c) + P(W = r, T = c) \\ &= 0.2 + 0.3 = 0.5 \end{aligned}$$

# Quiz: Conditional Probabilities

- $P(+x \mid +y) ?$

$P(X, Y)$

X	Y	P
+x	+y	0.2
+x	-y	0.3
-x	+y	0.4
-x	-y	0.1

- $P(-x \mid +y) ?$

- $P(-y \mid +x) ?$

# Conditional Distributions

- Conditional distributions are probability distributions over some variables given fixed values of others

Conditional Distributions

$$P(W|T)$$

$P(W T = hot)$	
W	P
sun	0.8
rain	0.2
$P(W T = cold)$	
W	P
sun	0.4
rain	0.6

Joint Distribution

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

# Normalization Trick

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

$$\begin{aligned} P(W = s|T = c) &= \frac{P(W = s, T = c)}{P(T = c)} \\ &= \frac{P(W = s, T = c)}{P(W = s, T = c) + P(W = r, T = c)} \\ &= \frac{0.2}{0.2 + 0.3} = 0.4 \end{aligned}$$



$$\begin{aligned} P(W = r|T = c) &= \frac{P(W = r, T = c)}{P(T = c)} \\ &= \frac{P(W = r, T = c)}{P(W = s, T = c) + P(W = r, T = c)} \\ &= \frac{0.3}{0.2 + 0.3} = 0.6 \end{aligned}$$

$P(W|T = c)$

W	P
sun	0.4
rain	0.6

# Normalization Trick

$$\begin{aligned} P(W = s|T = c) &= \frac{P(W = s, T = c)}{P(T = c)} \\ &= \frac{P(W = s, T = c)}{P(W = s, T = c) + P(W = r, T = c)} \\ &= \frac{0.2}{0.2 + 0.3} = 0.4 \end{aligned}$$

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

**SELECT** the joint probabilities matching the evidence  
→

T	W	P
cold	sun	0.2
cold	rain	0.3

$P(c, W)$

**NORMALIZE** the selection  
(make it sum to one)  
→

W	P
sun	0.4
rain	0.6

$P(W|T = c)$

$$\begin{aligned} P(W = r|T = c) &= \frac{P(W = r, T = c)}{P(T = c)} \\ &= \frac{P(W = r, T = c)}{P(W = s, T = c) + P(W = r, T = c)} \\ &= \frac{0.3}{0.2 + 0.3} = 0.6 \end{aligned}$$

# Normalization Trick

$P(T, W)$		
T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

**SELECT** the joint probabilities matching the evidence  
→

$P(c, W)$		
T	W	P
cold	sun	0.2
cold	rain	0.3

**NORMALIZE** the selection  
(make it sum to one)  
→

$P(W T = c)$	
W	P
sun	0.4
rain	0.6

- Why does this work? Sum of selection is  $P(\text{evidence})!$  ( $P(T=c)$ , here)

$$P(x_1|x_2) = \frac{P(x_1, x_2)}{P(x_2)} = \frac{P(x_1, x_2)}{\sum_{x_1} P(x_1, x_2)}$$

# Quiz: Normalization Trick

- $P(X | Y=-y) ?$

$P(X, Y)$

X	Y	P
+x	+y	0.2
+x	-y	0.3
-x	+y	0.4
-x	-y	0.1

**SELECT** the joint  
probabilities  
matching the  
evidence



**NORMALIZE** the  
selection  
(make it sum to one)



# To Normalize

- (Dictionary) To bring or restore to a **normal condition**

All entries sum to ONE

- Procedure:

- Step 1: Compute  $Z = \text{sum over all entries}$
- Step 2: Divide every entry by  $Z$

- Example 1

W	P
sun	0.2
rain	0.3

Normalize  $\rightarrow$   $Z = 0.5$

W	P
sun	0.4
rain	0.6

- Example 2

T	W	P
hot	sun	20
hot	rain	5
cold	sun	10
cold	rain	15

Normalize  $\rightarrow$   $Z = 50$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

# Probabilistic Inference

- Probabilistic inference: compute a desired probability from other known probabilities (e.g. conditional from joint)
- We generally compute conditional probabilities
  - $P(\text{on time} \mid \text{no reported accidents}) = 0.90$
  - These represent the agent's *beliefs* given the evidence
- Probabilities change with new evidence:
  - $P(\text{on time} \mid \text{no accidents, 5 a.m.}) = 0.95$
  - $P(\text{on time} \mid \text{no accidents, 5 a.m., raining}) = 0.80$
  - Observing new evidence causes *beliefs to be updated*



# Inference by Enumeration

- General case:

- Evidence variables:  $E_1 \dots E_k = e_1 \dots e_k$
- Query\* variable:  $Q$
- Hidden variables:  $H_1 \dots H_r$

$$\left. \begin{array}{l} E_1 \dots E_k = e_1 \dots e_k \\ Q \\ H_1 \dots H_r \end{array} \right\} X_1, X_2, \dots, X_n$$


*All variables*

- We want:


$$P(Q|e_1 \dots e_k)$$

\* Works fine with multiple query variables, too

- Step 1: Select the entries consistent with the evidence



- Step 2: Sum out H to get joint of Query and evidence



- Step 3: Normalize

$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} \underbrace{P(Q, h_1 \dots h_r, e_1 \dots e_k)}_{X_1, X_2, \dots, X_n}$$

$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$

# Inference by Enumeration

- $P(W)?$
- $P(W | \text{winter})?$
- $P(W | \text{winter, hot})?$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W)$ ?

- Step 1: no
- Step 2: sum over S and T

W	P
sun	$0.30+0.10+0.10+0.15= 0.65$
rain	$0.05+0.05+0.05+0.20= 0.35$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W)?$

- Step 1: no
- Step 2:
- Step 3: normalize

W	P
sun	$0.30+0.10+0.10+0.15= 0.65/1$
rain	$0.05+0.05+0.05+0.20= 0.35/1$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W | \text{winter})?$ 
  - Step 1: select  $S=\text{winter}$

S	T	W	P
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W | \text{winter})?$ 
  - Step 2: Sum over T

S	T	W	P
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

S	W	P
winter	sun	0.10+0.15= 0.25
winter	rain	0.05+0.25= 0.25

# Inference by Enumeration

- $P(W | \text{winter})?$ 
  - Step 3: normalize
    - $Z = 0.25 + 0.25 = 0.5$

W	$P(W   \text{Winter})$
sun	$0.25/0.5 = 0.5$
rain	$0.25/0.5 = 0.5$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W \mid \text{winter, hot})?$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W | \text{winter, hot})?$ 
  - Step 1:

S	T	W	P
winter	hot	sun	0.10
winter	hot	rain	0.05

- Step 2: no
- Step 3: normalize
  - $Z = 0.10 + 0.05 = 0.15$

W	P
sun	0.10/0.15
rain	0.05/0.15

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration


---

- Obvious problems:
  - Worst-case time complexity  $O(d^n)$
  - Space complexity  $O(d^n)$  to store the joint distribution

# The Product Rule

- Sometimes have conditional distributions but want the joint

$$P(y)P(x|y) = P(x, y) \quad \longleftrightarrow \quad P(x|y) = \frac{P(x, y)}{P(y)}$$



# The Product Rule

$$P(y)P(x|y) = P(x, y)$$

- Example:


R	P
sun	0.8
rain	0.2

$$P(D|W)$$

D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

$$P(D, W)$$

D	W	P
wet	sun	
dry	sun	
wet	rain	
dry	rain	



# The Chain Rule

---

- More generally, can always write any joint distribution as an incremental product of conditional distributions


$$P(x_1, x_2, x_3) = P(x_1)P(x_2|x_1)P(x_3|x_1, x_2)$$

$$P(x_1, x_2, \dots, x_n) = \prod_i P(x_i|x_1 \dots x_{i-1})$$

- Why is this always true?

# Bayes Rule

---



# Bayes' Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

That's my rule!

- Dividing, we get:

$$P(x|y) = \frac{P(y|x)}{P(y)}P(x)$$

- Why is this at all helpful?

- Lets us build one conditional from its reverse
- Often one conditional is tricky but the other one is simple
- Foundation of many systems we'll see later (e.g. ASR, MT)

- In the running for most important AI equation!



# Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{cause}|\text{effect}) = \frac{P(\text{effect}|\text{cause})P(\text{cause})}{P(\text{effect})}$$

- Example:

- M: meningitis, S: stiff neck

$$\left. \begin{array}{l} P(+m) = 0.0001 \\ P(+s|m) = 0.8 \\ P(+s|-m) = 0.01 \end{array} \right\} \text{Example givens}$$

$$P(+m|s) = \frac{P(+s|m)P(+m)}{P(+s)} = \frac{P(+s|m)P(+m)}{P(+s|m)P(+m) + P(+s|-m)P(-m)} = \frac{0.8 \times 0.0001}{0.8 \times 0.0001 + 0.01 \times 0.999}$$

- Note: posterior probability of meningitis still very small
  - Note: you should still get stiff necks checked out! Why?

# Quiz: Bayes' Rule

- Given:

$$P(W)$$

R	P
sun	0.8
rain	0.2

$$P(D|W)$$

D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

- What is  $P(W | \text{dry})$  ?

# Quiz: Bayes' Rule

- Given:

$P(W)$	
R	P
sun	0.8
rain	0.2

$P(D W)$		
D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

- What is  $P(W | \text{dry})$  ?

$$P(\text{sun}|\text{dry}) = \frac{P(\text{dry}|\text{sun})P(\text{sun})}{P(\text{dry})} = \frac{P(\text{dry}|\text{sun})P(\text{sun})}{P(\text{dry, sun})P(\text{sun}) + P(\text{dry}|\text{rain})P(\text{rain})} = \frac{0.9 \times 0.8}{0.9 \times 0.8 + 0.3 \times 0.2} = 0.92$$

$$P(\text{rain}|\text{dry}) = 1 - P(\text{sun}|\text{dry}) = 0.08$$



# Independence


- Two variables are *independent* in a joint distribution if:

$$P(X, Y) = P(X)P(Y)$$

$$X \perp\!\!\!\perp Y$$

$$\forall x, y P(x, y) = P(x)P(y)$$

- Says the joint distribution *factors* into a product of two simple ones
- Usually variables aren't independent!
- Can use independence as a *modeling assumption*
  - Independence can be a simplifying assumption
  - *Empirical* joint distributions: at best “close” to independent
  - What could we assume for {Weather, Traffic, Cavity}?
- Independence is like something from CSPs: what?



# Example: Independence?

---

$P_1(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

$P(T)$

T	P
hot	0.5
cold	0.5

$P(W)$

W	P
sun	0.6
rain	0.4

$P_2(T, W) = P(T)P(W)$

T	W	P
hot	sun	0.3
hot	rain	0.2
cold	sun	0.3
cold	rain	0.2

# Example: Independence

- N fair, independent coin flips:

$$P(X_1)$$

H	0.5
T	0.5



$$P(X_2)$$

H	0.5
T	0.5

...


$$P(X_n)$$

H	0.5
T	0.5




# Conditional Independence

---



# Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches it doesn't depend on whether I have a toothache:
  - $P(+\text{catch} | +\text{toothache}, +\text{cavity}) = P(+\text{catch} | +\text{cavity})$
- The same independence holds if I don't have a cavity:
  - $P(+\text{catch} | +\text{toothache}, -\text{cavity}) = P(+\text{catch} | -\text{cavity})$
- Catch is *conditionally independent* of Toothache given Cavity:
  - $P(\text{Catch} | \text{Toothache}, \text{Cavity}) = P(\text{Catch} | \text{Cavity})$
- Equivalent statements:
  - $P(\text{Toothache} | \text{Catch}, \text{Cavity}) = P(\text{Toothache} | \text{Cavity})$
  - $P(\text{Toothache}, \text{Catch} | \text{Cavity}) = P(\text{Toothache} | \text{Cavity}) P(\text{Catch} | \text{Cavity})$
  - One can be derived from the other easily



# Conditional Independence

---

- Unconditional (absolute) independence very rare (why?)
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments.
- X is conditionally independent of Y given Z  $X \perp\!\!\!\perp Y | Z$

if and only if:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

or, equivalently, if and only if

$$\forall x, y, z : P(x|z, y) = P(x|z)$$

# Conditional Independence

- What about this domain:

- Traffic
- Umbrella
- Raining



Raining


Traffic

Umbrella

# Conditional Independence

- What about this domain:




- Traffic
- Umbrella
- Raining



# Conditional Independence

- What about this domain:




- Fire
- Smoke
- Alarm



# Conditional Independence

- What about this domain:

- Fire
- Smoke
- Alarm



# Probability Recap

---

- Conditional probability

$$P(x|y) = \frac{P(x,y)}{P(y)}$$

- Product rule

$$P(x,y) = P(x|y)P(y)$$

- Chain rule

$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2)\dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$

- X, Y independent if and only if:  $\forall x, y : P(x,y) = P(x)P(y)$

- X and Y are conditionally independent given Z if and only if:  $X \perp\!\!\!\perp Y | Z$   
 $\forall x, y, z : P(x,y|z) = P(x|z)P(y|z)$

# Probability Recap

---

- **n variables, domain's size: d**
  - All dependence:
    - Size of table:  $d^n$
  - All independence (rare):
    - Size of table:  $d^n$
  - Conditional independence (practical):
    - Size of table:  $d^n \rightarrow d^n$