

# ETC3430: Financial mathematics under uncertainty

Dan Zhu

Monash Business School  
Monash University, Australia

March 23, 2022

## Holding times of Markov Processes

### Poisson Process

Definition

Properties of Poisson Process

The Transition Rate of Poisson Processes

Examples

# Time homogeneous: Holding times

Dan Zhu

Holding times of  
Markov Processes

Poisson Process

Definition

Properties of Poisson  
Process

The Transition Rate of  
Poisson Processes

Examples

## Theorem

*The holding time,  $\tau^i$ , of a time homogeneous Markov jump process with transition rates  $\mu_{i,j}$  given that its initial state is  $i$ , is an exponentially distributed random variable with  $\lambda_i = -\mu_{i,i}$ ,*

$$\mathbb{P}(\tau^i > t) = \mathbb{P}(T_1 > t | X_0 = i) = \exp^{-\lambda_i t}.$$

*Further more*

$$\mathbb{P}(X_{T_1} = j | X_0 = i) = \frac{\mu_{i,j}}{\lambda_i}.$$

*independent of  $T_1$ .*

Here  $\lambda_i$  is the total force out of state  $j$ .

## Proof.

We have

$$\begin{aligned}
 & \mathbb{P}(T_1 > t | X_0 = i) \\
 &= \prod_{j=0}^{n-1} \mathbb{P}(X_s > i, \frac{tj}{n} < s \leq \frac{tj+t}{n} | X_{\frac{tj}{n}} = i) \text{ for all } n \\
 &= \mathbb{P}(X_s > i, 0 < s \leq \frac{t}{n} | X_0 = i)^n \text{ time-homogeneity} \\
 &= \lim_{n \rightarrow \infty} (1 - \lambda_i \frac{t}{n} + o(\frac{t}{n}))^n = \exp^{-\lambda_i t}
 \end{aligned}$$

Hence  $\tau^i \sim \text{Exp}(\lambda_i)$ .

## Proof.

Probability jump is from  $i$  to  $j$  for  $j \neq i$  is

$$\begin{aligned} \lim_{h \rightarrow 0} \mathbb{P}(X_{t+h} = j | X_t = i, X_{t+h} \neq i) &= \lim_{h \rightarrow 0} \frac{\mathbb{P}(X_{t+h} = j | X_t = i)}{\mathbb{P}(X_{t+h} \neq i | X_t = i)} \\ &= \lim_{h \rightarrow 0} \frac{\mathbb{P}_{i,j}^{(h)}}{\sum_{k \neq i} \mathbb{P}_{i,k}^{(h)}} \\ &= \frac{\mu_{i,j}}{\sum_{k \neq i} \mu_{i,k}} \end{aligned}$$



# Counting Processes

Dan Zhu

## Definition (Counting Process)

*A counting process in continuous time,  $\{N_t, t \geq 0\}$ , has jumps of size +1 only, and whose paths are constant in between two jumps, i.e.*

$$N_t = \sum_{k=1}^{\infty} k \mathbb{I}_{[T_k, T_{k+1})}(t) = \sum_{k=1}^{\infty} \mathbb{I}_{[T_k, \infty)}(t)$$

*where  $(T_k)_{k \geq 1}$  is the increasing family of jump times such that  $\lim_{k \rightarrow \infty} T_k = +\infty$ .*

Notice, we can also recover the jump times from the counting process

$$T_k = \inf\{t \in \mathbb{R}_+ : N_t = k\}, k \geq 1.$$

Holding times of  
Markov Processes

Poisson Process

Definition

Properties of Poisson  
Process

The Transition Rate of  
Poisson Processes


Examples

**Definition**

Properties of Poisson  
Process

The Transition Rate of  
Poisson Processes

Examples



# Poisson Random Variables

Dan Zhu

Holding times of  
Markov Processes

Poisson Process

Definition

Properties of Poisson  
Process

The Transition Rate of  
Poisson Processes

Examples

Here, we briefly review some properties of the Poisson random variable that we have discussed in the previous chapters. Remember that a discrete random variable  $X$  is said to be a Poisson random variable with parameter  $\lambda$ , if its range is  $R_X = \{0, 1, 2, 3, \dots\}$ , with

$$P_X(k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!} & \text{for } k \in R_X \\ 0 & \text{otherwise} \end{cases}$$

Here are some useful facts that we have seen before:

- ▶ its mean and variance are equal to  $\lambda$
- ▶ the sum of independent Poissons is Poisson distributed with its parameter being the sum of the  $\lambda$ 's



## Definition

*A time homogeneous Poisson Process is a counting process satisfies the following conditions*

- 1. Independent increments, i.e.  $N_{t_4} - N_{t_3}$  is independent of  $N_{t_2} - N_{t_1}$  as long as  $[t_1, t_2)$  and  $[t_3, t_4)$  are disjoint time intervals in  $\mathbb{R}_+$ .*
- 2. Stationary increments, i.e.  $N_{t+h} - N_{s+h}$  has the same distribution as  $N_t - N_s$  for all  $h > 0$  and  $0 \leq s \leq t$ .*

## Theorem

*The increment of a time homogeneous Poisson process follows a Poisson distribution that*

$N_t - N_s \sim \text{Poisson}(\lambda(t-s))$  with the intensity  $\lambda = \lim_{h \rightarrow 0} \frac{1}{h} \mathbb{P}(N_t = 1)$ .

# Sum of independent Poisson Process

Dan Zhu

Holding times of  
Markov Processes

Poisson Process

Definition

Properties of Poisson  
Process

The Transition Rate of  
Poisson Processes

Examples

Given Poisson random variables  $N^a \sim \text{Poisson}(\lambda_a)$  and  $N^b \sim \text{Poisson}(\lambda_b)$ , their sum

$$N = N^a + N^b \sim \text{Poisson}(\lambda_a + \lambda_b).$$

<sup>1</sup> Hence, the natural extension of this result is that independent Poisson Processes is also a Poisson process with intensity equals the sum of the original intensities.

---

<sup>1</sup>Verify it via MGF.

## Theorem

*Let  $W_1 = T_1$ , and  $W_i = T_i - T_{i-1}$  for  $i = 2, 3, \dots$  denote the sequence of weighting times of a Poisson process with intensity  $\lambda$ . This sequence  $\{W_i\}$  are i.i.d exponential random variables with parameter  $\lambda$ .*

## Proof.

To show they are exponential, we have

$$\mathbb{P}(W_1 \leq t) = \mathbb{P}(N_t > 0) = 1 - \exp^{-\lambda t}$$

$$\mathbb{P}(W_i \leq t) = 1 - \mathbb{P}(N_{T_{i-1}+t} - N_{T_{i-1}} = 0) = 1 - \mathbb{P}(N_t = 0) = 1 - \exp^{-\lambda t}$$

## Proof.

To show independence, we consider

$$\begin{aligned} & \mathbb{P}(W_i > s_i | W_1 = s_1, W_2 = s_1, \dots, W_{i-1} = s_{i-1}) \\ &= \mathbb{P}(N_{\sum_{j=1}^i s_j} = i - 1 | W_1 = s_1, W_2 = s_1, \dots, W_{i-1} = s_{i-1}) \\ &= \mathbb{P}(N_{\sum_{j=1}^i s_j} - N_{\sum_{j=1}^{i-1} s_j} = 0) \\ &= \exp^{-\lambda s_i} = \mathbb{P}(W_i > s_i). \end{aligned}$$

## Question

*Consider a two state Markov chain with  $\mu_{1,1} = \mu_{2,2} = -\lambda$ , find the transition matrix  $\mathbb{P}(t)$ .*

This model has a very simple structure, assume  $X_0 = 0$ ,  $X_t = 0$  if and only if there is even number of transitions. Since the intensity of transition from 0 to 1 and 1 to 0 are the same, hence the time between each transition is  $\exp(\lambda)$ . This implies a Poisson process of parameter  $\lambda$ .

$$\begin{aligned}
 P_{00}(t) &= P(X(t) = 0 | X(0) = 0) \\
 &= P(\text{an even number of arrivals in } [0, t]) \\
 &= \sum_{n=0}^{\infty} e^{-\lambda t} \frac{(\lambda t)^{2n}}{(2n)!} \\
 &= e^{-\lambda t} \sum_{n=0}^{\infty} \frac{(\lambda t)^{2n}}{(2n)!} \\
 &= e^{-\lambda t} \left[ \frac{e^{\lambda t} + e^{-\lambda t}}{2} \right] = \frac{1}{2} + \frac{1}{2} e^{-2\lambda t}.
 \end{aligned}$$

Recall that

$$\begin{aligned}
 \sinh(x) &= \frac{e^x - e^{-x}}{2} = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}, \\
 \cosh(x) &= \frac{e^x + e^{-x}}{2} = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}.
 \end{aligned}$$

# Transition Rate of Poisson Process

Dan Zhu

The Poisson Process is that

$$\mathbb{P}(X_{t+h} = i + 1 | X_t = i) = \lambda h + o(h)$$

$$\mathbb{P}(X_{t+h} = i | X_t = i) = 1 - \lambda h + o(h).$$

Consider  $\mathbb{S} = \{0, 1, 2, 3, \dots\}$ , we have

$$\mathbb{P}_{i,j}^{(h)} = \begin{cases} 1 - \lambda h + o(h) & \text{if } j = i \\ \lambda h + o(h) & \text{if } j = i + 1 \\ 0 & \text{otherwise} \end{cases}$$

and

$$\mu_{i,j} = \begin{cases} -\lambda & \text{if } j = i \\ \lambda & \text{if } j = i + 1 \\ 0 & \text{otherwise} \end{cases}$$

Holding times of  
Markov Processes

Poisson Process

Definition

Properties of Poisson  
Process

The Transition Rate of  
Poisson Processes

Examples



## Example

*The number of customers arriving at a grocery store can be modeled by a Poisson process with intensity  $\lambda = 10$  customers per hour. Find the probability that there are 2 customers between 10:00 and 10:20. Find the probability that there are 3 customers between 10:00 and 10:20 and 7 customers between 10:20 and 11.*

First  $X \sim \text{Pois}(\frac{10}{3})$ ,

$$P(X = 2) = \frac{e^{-\frac{10}{3}} \left(\frac{10}{3}\right)^2}{2!} \\ \approx 0.2.$$

Then, we have two non-overlapping intervals,

$$\begin{aligned} & P\left(3 \text{ arrivals in } I_1 \text{ and } 7 \text{ arrivals in } I_2\right) \\ &= P\left(3 \text{ arrivals in } I_1\right) \cdot P\left(7 \text{ arrivals in } I_2\right) \\ &= \frac{e^{-\frac{10}{3}} \left(\frac{10}{3}\right)^3}{3!} \cdot \frac{e^{-\frac{20}{3}} \left(\frac{20}{3}\right)^7}{7!} \\ &\approx 0.0325 \end{aligned}$$

## Example

Let  $N_t$  be a Poisson process with intensity  $\lambda = 2$ , and let  $X_1, X_2, \dots$  be the corresponding interarrival times.

1. Find the probability that the first arrival occurs after  $t > 0.5$
2. Given that we have had no arrivals before  $t = 1$ , find  $\mathbb{P}(X_1 > 3)$
3. Given that the third arrival occurred at time  $t = 2$ , find the probability that the fourth arrival occurs after  $t = 4$ .
4. I start watching the process at time  $t = 10$ . Let  $T$  be the time of the first arrival that I see. In other words,  $T$  is the first arrival after  $t = 10$ . Find  $\mathbb{E}[T]$  and  $\text{var}(T)$

1.  $X_1 \sim \text{Exp}(2)$ ,

$$P(X_1 > 0.5) = e^{-(2 \times 0.5)} \approx 0.37$$

2.

$$\begin{aligned} P(X_1 > 3 | X_1 > 1) &= P(X_1 > 2) \text{ (memoryless property)} \\ &= e^{-2 \times 2} \approx 0.0183 \end{aligned}$$

3.  $X_4 \sim \text{Exp}(2)$ ,

$$\begin{aligned} P(X_4 > 2 | X_1 + X_2 + X_3 = 2) &= P(X_4 > 2) \text{ (independence of the } X_i\text{'s)} \\ &= e^{-2 \times 2} \approx 0.0183 \end{aligned}$$

4. When I start watching the process at time  $t = 10$ , I will see a Poisson process. Thus, the time of the first arrival from  $t = 10$  is  $\text{Exp}(2)$ . In other words, we can write  $T = 10 + X$  with  $X \sim \text{Exp}(2)$ . Thus,  $\mathbb{E}[T] = 10.5$  and  $\text{Var}(T) = 0.25$ .