AP Physics 1

Name _____

Period _____ Date _____

Practice Test – Introductory Dynamics (Version 2)

On any problems, you must show all work, include units, round properly, and box your final answer. Do the algebra first, before plugging in values.

Rewrite the following statement in the space below: On my honor, before God, I have neither given, nor received, any aid on this test.

For all questions:

- Use g = 9.8 m/s² (9.8 N/kg). Air resistance is negligible unless the question says it is not.
- Free-body diagrams: draw the forces themselves, not components, each as an arrow starting on the dot.
- Round answers to the significant figures of the given values.
- 50 minutes. Four questions, 25 points each (100 points total).


1. (25 points) Students measure the drag force F_D on a soccer ball in a wind tunnel at several air speeds v. The drag force is modeled by $F_D = Kv^2$, where K is a constant.

Speed v (m/s)	**Drag force F_D (N)**	
10.0	0.619	
15.0	1.41	
20.0	2.58	
25.0	3.71	
30.0	5.79	
35.0	7.72	

(a) Using $F_D = Kv^2$, state the quantity you will plot on the vertical axis and the quantity you will plot on the horizontal axis so the data form a straight line, each with its units. State what the slope equals.


(b) Use the blank column in the table to calculate the values you will plot on the horizontal axis.

(c) Plot the data on the grid. Label each axis with the quantity and units, choose a scale, and draw the best-fit line.




(d) Calculate the slope of your best-fit line using two well-separated points on the line. Then give K with its units.

2. (25 points) A 240. g squid has lifted off the floor of an aquarium. It squirts water downward, so the water pushes it upward with a thrust of 0.40 N. After it leaves the bottom, starting from rest just above the floor, it rises 0.60 m in 2.0 s.



Assumptions for this practice problem: *water drag on the squid is negligible; its acceleration is constant; g = 9.8 m/s².*

(a) Draw the free-body diagram of the squid after it has left the bottom.




(b) Calculate the squid's acceleration, magnitude and direction.

(c) With up as positive, write Newton's second law for the squid and solve it for the buoyant force in letters.


(d) Calculate the magnitude of the buoyant force on the squid.

3. (25 points) A loaded toboggan slides down a snowy hill. In the picture the toboggan moves down the slope to the left and is moving at a constant speed. Friction from the snow and drag from the air are **not** negligible. No angle is given.




Assumptions for this practice problem: *the air is still (no wind), so the air moves past the toboggan opposite its motion.*

Draw and label the free-body diagram of the toboggan. Draw the forces themselves, not components. Each force is one arrow that starts on the dot and points in the direction the force acts. No calculations.



4. (25 points) A skater standing on smooth ice pushes a medicine ball horizontally to the right. Find the skater's acceleration during the push.



medicine ball mass	4.00 kg
skater mass	60.0 kg
time her hands push the ball	250. ms
speed the ball leaves her hands, relative to the ground	18.0 km/h

The ball's speed is measured relative to the ground.

Assumptions for this practice problem: *the skater starts at rest and slides freely on the ice (ignore friction and air resistance); the medicine ball starts from rest; its acceleration during the push is constant; the push between the medicine ball and the skater is the only horizontal interaction.*

(a) Draw the free-body diagram of the skater during the push.

(a) skater

(b) Calculate the medicine ball's acceleration relative to the ground during the push.

(c) Calculate the magnitude of the force the skater exerts on the medicine ball, and the magnitude of the force the medicine ball exerts on the skater.

(d) Calculate the skater's acceleration, magnitude and direction.

Answer Key – Practice Test, Version 2 – Introductory Dynamics

Worked step by step: diagram, symbolic equations and algebra first, then numbers with units. g = 9.8 m/s² does not limit significant figures; answers match the significant figures of the givens.

Numerical problems: diagram first, then a = ΣF/m (equation sheet), written a = F_{net}/m. Replace F_{net} with the actual forces and their signs, swap F_g = mg, solve in letters, then plug in numbers with units.
Kinematics: #1 $v = v_0 + at$ · #2 $\Delta x = v_0t + \frac{1}{2}at^2$ · #3 $v^2 = v_0^2 + 2a\Delta x$. Pick the one that leaves out the variable you don't know and don't need.

1. Drag linearization (25 points)

Step 1 **(a) Match $F_D = Kv^2$ to y = mx + b.**

$F_D = K \cdot (v^2) + 0$

vertical (y)	horizontal (x)	slope	intercept
F_D (N)	v^2 (m^2/s^2)	K	0

Vertical: F_D in N. Horizontal: v^2 in m^2/s^2. Slope = K.


Watch out: v^2 is speed squared, in m^2/s^2. It is not an acceleration (m/s^2).

Step 2 **(b) Square each speed; the units square too.**

Example: $(10.0\ m/s)^2 = 100\ m^2/s^2$

v (m/s)	F_D (N)	v^2 (m^2/s^2)
10.0	0.619	100
15.0	1.41	225
20.0	2.58	400
25.0	3.71	625
30.0	5.79	900
35.0	7.72	1225

Step 3 **(c) Graph: labeled axes with units, a scale that uses more than half the grid, one straight line through the middle of the points.**



The best-fit line is not forced through any data point. The red squares are points read off the line, not data points.

Step 4 **(d) Slope from two well-separated points on the line, units carried.**


Points read off the line, away from data points: (150 m²/s², 0.96 N) and (1175 m²/s², 7.44 N)

$$\text{slope} = \frac{7.44\ \text{N} - 0.96\ \text{N}}{1175\ \text{m}^2/\text{s}^2 - 150\ \text{m}^2/\text{s}^2} = \frac{6.48\ \text{N}}{1025\ \text{m}^2/\text{s}^2} = 0.006322\ \text{N}\cdot\text{s}^2/\text{m}^2$$

$K \approx 0.00632\ \text{N}\cdot\text{s}^2/\text{m}^2\ (= \text{kg/m})$. Any value from 0.00582 to 0.00684 N·s²/m² from a reasonable line earns credit.

Units: N ÷ (m²/s²) = N·s²/m², and N = kg·m/s², so N·s²/m² = kg/m.

2. Thrust and buoyancy (25 points)



Step 1 **(a) Forces on the squid after it leaves the bottom.**

F_g down (Earth). Thrust F_t up (the water it pushes down pushes back up on it). Buoyancy F_b up (the surrounding water).

No normal force: it is no longer touching the bottom. Acceleration is up, so $F_t + F_b$ is longer than F_g.

Step 2 **(b) Acceleration from the motion data.**

From rest, time and displacement known, final speed not needed: use #2.

$$\Delta y = v_0 t + \tfrac{1}{2}at^2 \rightarrow \Delta y = 0 + \tfrac{1}{2}at^2 \rightarrow a = \frac{2\Delta y}{t^2} = \frac{2(0.60\text{ m})}{(2.0\text{ s})^2} = \boxed{0.30\text{ m/s}^2\text{, upward}}$$

Step 3 **(c) Newton's second law, up is +, letters first.**

$$a = \frac{F_{net}}{m} \rightarrow a = \frac{F_b + F_t - F_g}{m} \rightarrow F_b + F_t - mg = ma$$

$$\boxed{F_b = m(g + a) - F_t}$$

Step 4 **(d) Now the numbers, units in every parenthesis.**


$m = 240.\ g = 0.240$ kg. $F_b = (0.240\text{ kg})(9.8\text{ m/s}^2 + (0.30\text{ m/s}^2)) - 0.40\text{ N} = (0.240\text{ kg})(10.1\text{ m/s}^2) - 0.40\text{ N} = 2.424\text{ N} - 0.40\text{ N} = 2.024\text{ N}$

$\boxed{F_b = 2.02\text{ N}}$ (the subtraction sets the decimal place; g does not limit the significant figures).

Watch out: keep the sign of a. Using a = 0 just because the squid moves up, or dropping the thrust, gives the wrong buoyant force.

3. Free-body diagram on a slope (25 points)

toboggan, moving downhill to the left



Step 1 **List every force on the toboggan: exactly four.**

Earth pulls on it: F_g. The snow pushes on it two ways: F_N (perpendicular to the surface) and kinetic friction F_f (along the surface). The air pushes on it: drag F_D.

Step 2 **Directions. No angle is needed.**

F_g: straight down, vertical, not perpendicular to the slope.

F_N: perpendicular to the slope, pointing out of the surface, so it tilts away from vertical toward the left.

F_f: along the slope, uphill (up and to the right), opposite the sliding.

F_D: opposite the motion through the air. The air is still and the toboggan moves downhill, so drag also points uphill, parallel to F_f.


Step 3 **Check the diagram.**

Four arrows, each starting on the dot, each labeled. F_f and F_D are two separate arrows, side by side, both uphill. Speeding up, slowing down or moving at a constant speed changes the arrow lengths, not their directions.

Lengths in this key: constant speed, so the two uphill arrows together balance the part of gravity that pulls along the slope, and FN is a little shorter than Fg. Lengths are not graded.

Watch out: do not add the components of F_g (along and perpendicular to the slope) as extra arrows; they are not new forces. Drag is the air resistance, not a fifth force, and there is no forward “motion” force.

4. Recoil (25 points)



Step 1 **(a) Forces on the skater during the push.**

Vertical: the ice pushes up (F_N); Earth pulls down (F_g); the medicine ball presses down ($F_{ball,\ down}$). Nothing accelerates vertically, so $F_N = F_g + F_{ball,\ down}$.

Horizontal: the only force is the medicine ball pushing back on the skater, $F_{ball\ on\ skater}$, to the left. The vertical forces do not enter the horizontal net-force equation. Arrow lengths are not to scale.

Step 2 **Units and frames.**

Convert first: 250. ms = 0.250 s; $18.0\ \text{km/h} \times \frac{1000\ \text{m}}{1\ \text{km}} \times \frac{1\ \text{h}}{3600\ \text{s}} = 5.00\ \text{m/s}$.

Step 3 **(b) Kinematics for the medicine ball.**

Ball, relative to the ground: $v_0 = 0$, $v = 5.00$ m/s, $t = 0.250$ s. Distance is not given: use #1.

$v = v_0 + at \rightarrow a_{ball} = \frac{v - 0}{t} = \frac{5.00\ \text{m/s}}{0.250\ \text{s}} =$ **$20.0\ \text{m/s}^2$**

Step 4 **(c) Second law on the medicine ball, then the third law.**

$a = \frac{F_{net}}{m} \rightarrow F_{skater\ on\ ball} = m_{ball}a_{ball} = (4.00\ \text{kg})(20.0\ \text{m/s}^2) = 80.0\ \text{N} \rightarrow$ **80.0 N, forward on the medicine ball**

Third law: $F_{ball\ on\ skater} = -F_{skater\ on\ ball} \rightarrow$ **80.0 N, backward on the skater**

Step 5 **(d) Second law on the skater, horizontal only.**

$a_{skater} = \frac{F_{ball\ on\ skater}}{M} = \frac{80.0\ \text{N}}{60.0\ \text{kg}} =$ **$1.33\ \text{m/s}^2$, to the left (backward)**

Equal forces, unequal accelerations: the skater has 15 times the mass, so her acceleration is 15 times smaller.

Watch out: convert first: 250. ms is 0.250 s and 18.0 km/h is 5.00 m/s.

1. Drag linearization

Suggested partial credit	pts
(a) F_D (N) on the vertical axis, v^2 (m^2/s^2) on the horizontal axis	4
(a) Slope = K	2
(b) v^2 column correct, with units	4
(c) Axes labeled with quantity and units	2
(c) Scale uses more than half the grid	2
(c) Points plotted correctly	2
(c) One straight best-fit line	2
(d) Two well-separated points read off the line, not data points	3
(d) K from 0.00582 to 0.00684, with units $N \cdot s^2/m^2$ (or kg/m)	4
Total	**25**

2. Thrust and buoyancy

Suggested partial credit	pts
(a) F_g down, thrust and buoyancy up, labeled	4
(a) No normal force	2
Acceleration: correct kinematics equation, magnitude and direction (sign)	6
$F_b + F_t - mg = ma$ written with up as +	4
Solved in letters: $F_b = m(g + a) - F_t$	3
Substitution with units in every parenthesis (and grams to kg if needed)	3
$F_b = 2.02$ N	3
Total	**25**

3. Free-body diagram on a slope

Suggested partial credit	pts
F_g straight down (vertical), labeled	5
F_N perpendicular to the slope, out of the surface, labeled	5
F_f along the slope, uphill, labeled	5
F_D uphill (opposite the motion through still air), labeled, separate from F_f	5
Every arrow starts on the dot; no components and no extra forces	5
Total	**25**

4. Recoil

Suggested partial credit	pts
(a) F_N up, F_g down (the medicine ball pressing down may also be shown); backward push from the medicine ball; no forward force	5
Unit conversions (or none needed) and the matching ground-frame values	3
(b) Kinematics equation chosen and solved in letters; $a_{ball} = 20.0\ m/s^2$	5
(c) $F = m_{ball}a_{ball} = 80.0$ N, and the third law gives the same size back on the skater	5
(d) acceleration = F/M = 1.33 m/s^2, backward	7
Total	**25**