PCBC H Parametric & Sequences Test (Practice A) **Name**_____ _____/27

SHOW ALL WORK FOR FULL CREDIT. ***Calculator*** **You have the entire period.**

Honor Code (copy below & sign): On my honor, before God, I have neither given, nor received any aid on this test.

_____

_____ _____

Signature

Solve the following application problem. Report answers to the thousandths. Include units in your answers. (4 points)

7. A golf ball is hit from the ground at an angle of 28° with an initial speed of 110 ft/s.

a) Write a set of parametric equations for the motion of the golf ball. (Recall: $x = (v\cos\theta)t,\ \ y = (v\sin\theta)t - 16t^2 + h$)

b) Determine the maximum height reached by the golf ball.

c) Determine how long the golf ball is in the air.

d) Determine how far the golf ball traveled in the air.

The given pattern continues. Write down the nth term of the sequence $\{a_n\}$ suggested by the pattern. (3 points)

8. $-\frac{2}{5}, \frac{4}{7}, -\frac{6}{9}, \frac{8}{11}, \cdots$

Write out the sum. (3 points)

9. $\sum_{k=1}^{n} \left(k^2 - 2k\right)$

Find each sum. Show the properties and formulas you use, then use your calculator. (#10 and #11 are 3 points each, #12 is 4 points)

10. $\sum_{k=1}^{35} (2k - 7)$

11. $\sum_{k=1}^{15} \left(k^2 + 3k\right)$

12. $\sum_{k=8}^{20} \left(k^3 - k\right)$

Expand using the Binomial Theorem. (#13 is 3 points, #14 is 4 points)

13. $(x+3)^6$

14. $(2x-1)^5$

Extra practice: more sequence and summation questions

Not part of the 27 points. Worked in the key.

Express the sum using summation notation.

E1. $\frac{1}{4}+\frac{1}{9}+\frac{1}{16}+\cdots+\frac{1}{121}$

The given pattern continues. Write down the nth term of the sequence $\{a_n\}$ suggested by the pattern.

E2. $\frac{2}{3}, \frac{4}{9}, \frac{8}{27}, \frac{16}{81}, \cdots$

Answer Key

Work on the left, the reason beside it. Boxed = final answer.

Day 2: Calculator

7. Golf ball: $h = 0$ ft, $\theta = 28°$, $v = 110$ ft/s. Calculator in **degree** mode, Graph in **Parametric**.

a) $x = (110\cos 28°)t$ — *plug v and θ into the given formula; degree symbol required*

$y = (110\sin 28°)t - 16t^2 + 0$ — *+h: starts at 0 ft*

$x_1(t) = 110\cos(28°)\cdot t,\ \ y_1(t) = 110\sin(28°)\cdot t - 16t^2 + 0$ — *what you type in*

$0 \le t \le 4$, tstep 0.01; x : 0 to 330, y : 0 to 47 — *graph doesn't come back down? raise tmax*

b) $\boxed{y \approx 41.670 \text{ ft}}$ — *trace to the top ($t \approx 1.614$); not Analyze, Maximum*

c) $\boxed{t \approx 3.228 \text{ s}}$ — *trace to $y \approx 0$; trace step 0.01, then 0.001*

d) $\boxed{x \approx 313.480 \text{ ft}}$ — *same point, read x*

Units: feet (the speed was ft/s).

8. $-\frac{1}{5}, \frac{2}{7}, -\frac{3}{9}, \frac{4}{11}, \ldots$

top: $2n$, bottom: $2n+3$ — *write 1 2 3 4 above; top and bottom separately*

$-,+,-,+ \Rightarrow (-1)^n$ — *first term negative*

$\boxed{a_n = (-1)^n \frac{2n}{2n+3}}$ — *check $n = 1$: $-\frac{2}{5}$*

9. $\sum_{k=1}^{n}(k^2 - 2k)$

$(1-2) + (4-4) + (9-6) + \cdots + (n^2 - 2n)$ — *$k = 1,2,3$, then the last term with n*

$\boxed{-1 + 0 + 3 + \cdots + (n^2 - 2n)}$ — *don't add*

10.

$2\sum_{k=1}^{35} k - 7\sum_{k=1}^{35} 1$ — *(2), (3) split; (1) constants out*

$2\left[\frac{35(36)}{2}\right] - 7(35)$ — *(6), (5) with $n = 35$; then the calculator*

$\boxed{1015}$

11.

$\sum_{k=1}^{15} k^2 + 3\sum_{k=1}^{15} k$ — *(2), (3) split; (1) constants out*

$\left[\frac{15(16)(31)}{6}\right] + 3\left[\frac{15(16)}{2}\right]$ — *(7), (6) with $n = 15$; then the calculator*

$\boxed{1600}$

12.

$\sum_{k=8}^{20} k^3 - \sum_{k=8}^{20} k$ — *(2), (3) split; (1) constants out*

$\left[\sum_{k=1}^{20} k^3 - \sum_{k=1}^{7} k^3\right] - \left[\sum_{k=1}^{20} k - \sum_{k=1}^{7} k\right]$ — *(4): starts at 8, so $j = 7$*

$\left[\left[\frac{20(21)}{2}\right]^2 - \left[\frac{7(8)}{2}\right]^2\right] - \left[\frac{20(21)}{2} - \frac{7(8)}{2}\right]$ — *(8), (6); then the calculator*

$\boxed{43134}$

13. $(x+3)^6$: $n = 6$, row 1 6 15 20 15 6 1. The sign goes with its term.

$1x^6(3)^0 + 6x^5(3)^1 + 15x^4(3)^2 + 20x^3(3)^3$ — *blanks first: first term's powers count down, second term's count up*

$+15x^2(3)^4 + 6x^1(3)^5 + 1x^0(3)^6$

$\boxed{x^6 + 18x^5 + 135x^4 + 540x^3 + 1215x^2 + 1458x + 729}$

Second term positive: every term positive.

14. $(2x-1)^5$: $n = 5$, row 1 5 10 10 5 1. The sign goes with its term.

$1(2x)^5(-1)^0 + 5(2x)^4(-1)^1 + 10(2x)^3(-1)^2 + 10(2x)^2(-1)^3$ — *blanks first: first term's powers count down, second term's count up*

$+5(2x)^1(-1)^4 + 1(2x)^0(-1)^5$

$(2x)^5 = 32x^5$, … — *the whole first term gets the power*

$\boxed{32x^5 - 80x^4 + 80x^3 - 40x^2 + 10x - 1}$

Second term negative: odd powers of -1 are negative, so signs alternate.

Extra practice (not scored)

E1. $\frac{1}{4} + \frac{1}{9} + \cdots + \frac{1}{121}$

$\frac{1}{2^2}, \frac{1}{3^2}, \frac{1}{4^2}, \ldots, \frac{1}{11^2}$ — *bottoms are squares, starting at 2^2*

$\boxed{\sum_{k=2}^{11} \frac{1}{k^2}}$ — *lower index 2, not 1*

E2. $\frac{1}{2}, \frac{2}{4}, \frac{3}{8}, \frac{4}{16}, \ldots$

top: 2^n, bottom: 3^n — *write 1 2 3 4 above; top and bottom separately*

$\boxed{a_n = \left(\frac{2}{3}\right)^n}$ — *same as $\frac{2^n}{3^n}$*