

*Non-Calculator***Honor Code (copy below & sign): On my honor, before God, I have neither given, nor received any aid on this quiz.****You have 30 minutes.**_____
Signature**SHOW ALL WORK FOR FULL CREDIT!**

Find the product. Write the product in rectangular form, using exact values. (3 points)

1. $\left[4\sqrt{2}(\cos 165^\circ + i \sin 165^\circ) \right] \left[3\sqrt{2}(\cos 240^\circ + i \sin 240^\circ) \right]$

Convert to trigonometric form, then divide. Write the quotient in rectangular form, using exact values. (3 points)


2. $\frac{-3\sqrt{6} + 3i\sqrt{2}}{\sqrt{3} - 3i}$

Find the power. Write the answer in rectangular form, using exact values. (3 points)

3. $(\sqrt{6} - i\sqrt{2})^5$

Find the fourth roots of the number. Leave answers in trig form. Graph each root as a vector in the complex plane. (3 points)

4. $-8 + 8i\sqrt{3}$



Find all complex number solutions of the equation. Write answers in rectangular form. (3 points)

5. $x^3 + 27i = 0$

ANSWER KEY – Non-Calculator

SHOW ALL WORK FOR FULL CREDIT!*Every line shown. Point splits are an estimate of partial credit (halves are possible).*

Find the product. Write the product in rectangular form, using exact values. (3 points)

Problem: $\left[4\sqrt{2}(\cos 165^\circ + i \sin 165^\circ)\right] \left[3\sqrt{2}(\cos 240^\circ + i \sin 240^\circ)\right]$

1. Multiply the r 's, add the θ 's.

$$r = (4\sqrt{2})(3\sqrt{2}) = 12 \cdot 2 = 24 \quad \theta = 165^\circ + 240^\circ = 405^\circ$$

$$405^\circ - 360^\circ = 45^\circ \quad (\text{keep the angle between } 0^\circ \text{ and } 360^\circ)$$

$$24(\cos 45^\circ + i \sin 45^\circ) = 24\left(\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2}\right)$$

$$= \boxed{12\sqrt{2} + 12\sqrt{2}i}$$

*Trap: $\sqrt{2} \cdot \sqrt{2} = 2$, so $r = 12 \cdot 2 = 24$. Leaving $12\sqrt{4}$, or writing $12\sqrt{2}$, costs the point.**1 pt r, 1 pt reduced angle, 1 pt exact rectangular.*

Convert to trigonometric form, then divide. Write the quotient in rectangular form, using exact values. (3 points)

Problem: $\frac{-3\sqrt{6} + 3i\sqrt{2}}{\sqrt{3} - 3i}$

2. Convert the top and the bottom separately (sketch each for the quadrant).

Top: $r = \sqrt{(-3\sqrt{6})^2 + (3\sqrt{2})^2} = \sqrt{54 + 18} = \sqrt{72} = 6\sqrt{2} \quad \tan \theta = \frac{3\sqrt{2}}{-3\sqrt{6}} = -\frac{1}{\sqrt{3}} \Rightarrow \text{ref } 30^\circ, \text{ QII} \Rightarrow 150^\circ$

Bottom: $r = \sqrt{(\sqrt{3})^2 + (-3)^2} = \sqrt{3 + 9} = \sqrt{12} = 2\sqrt{3} \quad \tan \theta = \frac{-3}{\sqrt{3}} = -\sqrt{3} \Rightarrow \text{ref } 60^\circ, \text{ QIV} \Rightarrow 300^\circ$

$$\frac{6\sqrt{2} \text{ cis } 150^\circ}{2\sqrt{3} \text{ cis } 300^\circ} = \frac{3\sqrt{2}}{\sqrt{3}} \text{ cis}(150^\circ - 300^\circ) = \frac{3\sqrt{6}}{3} \text{ cis}(-150^\circ) = \sqrt{6} \text{ cis}(-150^\circ + 360^\circ) = \sqrt{6} \text{ cis } 210^\circ$$

$$= \sqrt{6} \left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = -\frac{\sqrt{18}}{2} - \frac{\sqrt{6}}{2}i = \boxed{-\frac{3\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i}$$

Trap: $(-3\sqrt{6})^2 = 9 \cdot 6 = 54$ (square the 3 too). Negative angle: add 360° . Rationalize $\frac{3\sqrt{2}}{\sqrt{3}}$ and simplify $\sqrt{18} = 3\sqrt{2}$.

Find the power. Write the answer in rectangular form, using exact values. (3 points)

Problem: $(\sqrt{6} - i\sqrt{2})^5$

3. Convert, then De Moivre: $[r \operatorname{cis} \theta]^n = r^n \operatorname{cis}(n\theta)$.

$$r = \sqrt{(\sqrt{6})^2 + (-\sqrt{2})^2} = \sqrt{6+2} = \sqrt{8} = 2\sqrt{2}$$

$$\tan \theta = \frac{-\sqrt{2}}{\sqrt{6}} = -\frac{1}{\sqrt{3}} \Rightarrow \text{ref } 30^\circ, \text{ QIV} \Rightarrow 330^\circ$$

$$(2\sqrt{2} \operatorname{cis} 330^\circ)^5 = (2\sqrt{2})^5 \operatorname{cis}(5 \cdot 330^\circ) = 2^5 (\sqrt{2})^5 \operatorname{cis} 1650^\circ$$

$$= 32 \cdot 4\sqrt{2} \operatorname{cis} 1650^\circ = 128\sqrt{2} \operatorname{cis} 1650^\circ$$

$$1650^\circ - 4(360^\circ) = 1650^\circ - 1440^\circ = 210^\circ$$

$$128\sqrt{2}(\cos 210^\circ + i \sin 210^\circ) = 128\sqrt{2} \left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i \right) = \boxed{-64\sqrt{6} - 64\sqrt{2}i}$$

Trap: $(\sqrt{2})^5 = \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 4\sqrt{2}$, not $5\sqrt{2}$ or $2\sqrt{5}$.

Find the fourth roots of the number. Leave answers in trig form. Graph each root as a vector in the complex plane. (3 points)

Problem: $-8 + 8i\sqrt{3}$

4. Convert, then the n th root theorem: $\sqrt[n]{r} \operatorname{cis} \left(\frac{\theta}{n} + \frac{360^\circ k}{n} \right), k = 0, 1, \dots, n-1$.

$$r = \sqrt{(-8)^2 + (8\sqrt{3})^2} = \sqrt{64 + 192} = \sqrt{256} = 16$$

$$\tan \theta = \frac{8\sqrt{3}}{-8} = -\sqrt{3} \Rightarrow \text{ref } 60^\circ, \text{ QII} \Rightarrow 120^\circ$$


$$\sqrt[4]{16} = 2 \quad \text{first angle } \frac{120^\circ}{4} = 30^\circ$$

$$\text{spacing } \frac{360^\circ}{4} = 90^\circ$$

$$\boxed{2 \operatorname{cis} 30^\circ, 2 \operatorname{cis} 120^\circ, 2 \operatorname{cis} 210^\circ, 2 \operatorname{cis} 300^\circ}$$

Stop at four roots: the next one, 390° , is 30° again.

Trap: $\sqrt{256} = 16$, then $\sqrt[4]{16} = 2$ (not 4).



four equal vectors, 90° apart

Find all complex number solutions of the equation. Write answers in rectangular form. (3 points)

Problem: $x^3 + 27i = 0$

5. Isolate x^3 first, then take cube roots of the constant.

$$x^3 + 27i = 0 \Rightarrow x^3 = -27i = 0 - 27i \quad (\text{negative imaginary axis})$$

$$r = 27, \quad \theta = 270^\circ \quad x = (27 \operatorname{cis} 270^\circ)^{1/3}$$
$$\sqrt[3]{27} = 3 \quad \frac{270^\circ}{3} = 90^\circ \quad \text{spacing } \frac{360^\circ}{3} = 120^\circ$$
$$x = 3 \operatorname{cis} 90^\circ, 3 \operatorname{cis} 210^\circ, 3 \operatorname{cis} 330^\circ$$

$$3 \operatorname{cis} 90^\circ = 3(0 + i) = 3i$$

$$3 \operatorname{cis} 210^\circ = 3\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = -\frac{3\sqrt{3}}{2} - \frac{3}{2}i$$

$$x = 3i, \quad -\frac{3\sqrt{3}}{2} - \frac{3}{2}i, \quad \frac{3\sqrt{3}}{2} - \frac{3}{2}i$$

$$3 \operatorname{cis} 330^\circ = 3\left(\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = \frac{3\sqrt{3}}{2} - \frac{3}{2}i$$

Check one: $(3i)^3 = 27i^3 = 27(-i) = -27i \checkmark$

Trap: moving $+27i$ across makes it $-27i$, which is 270° , not 90° . Using 90° gives the wrong three roots.

*Non-Calculator***Honor Code (copy below & sign): On my honor, before God, I have neither given, nor received any aid on this quiz.****You have 30 minutes.**_____
Signature**SHOW ALL WORK FOR FULL CREDIT!**

Find the product. Write the product in rectangular form, using exact values. (3 points)

1. $\left[2 \left(\cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4} \right) \right] \left[5 \left(\cos \frac{11\pi}{12} + i \sin \frac{11\pi}{12} \right) \right]$

Find the quotient. Write it in rectangular form, using exact values. (3 points)


2. $\frac{\sqrt{7} + i\sqrt{21}}{-3\sqrt{21} - 3i\sqrt{7}}$

Find the power. Write the answer in rectangular form. (3 points)

3. $(-2 + 3i)^4$

Find the fifth roots of the number. Leave answers in trig form with θ in radians. Graph each root as a vector in the complex plane. (3 points)

4. $16 - 16i\sqrt{3}$



One of the fourth roots of a complex number w is $-\sqrt{2} + i\sqrt{2}$. (3 points)

5. a) Determine the rectangular form of its other three fourth roots.

b) Find w in rectangular form.

ANSWER KEY – Non-Calculator

SHOW ALL WORK FOR FULL CREDIT!

Every line shown. Point splits are an estimate of partial credit (halves are possible).

Find the product. Write the product in rectangular form, using exact values. (3 points)

Problem: $\left[2 \left(\cos \frac{5\pi}{4} + i \sin \frac{5\pi}{4} \right) \right] \left[5 \left(\cos \frac{11\pi}{12} + i \sin \frac{11\pi}{12} \right) \right]$

1. Multiply the r 's, add the θ 's (common denominator first).

$$r = 2 \cdot 5 = 10 \quad \theta = \frac{5\pi}{4} + \frac{11\pi}{12} = \frac{15\pi}{12} + \frac{11\pi}{12} = \frac{26\pi}{12} = \frac{13\pi}{6}$$

$$\frac{13\pi}{6} - 2\pi = \frac{13\pi}{6} - \frac{12\pi}{6} = \frac{\pi}{6} \quad (\text{keep between } 0 \text{ and } 2\pi)$$

$$10 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) = 10 \left(\frac{\sqrt{3}}{2} + \frac{1}{2}i \right) = \boxed{5\sqrt{3} + 5i}$$

Trap: given in radians, stay in radians. $2\pi = \frac{12\pi}{6}$, so subtract $\frac{12\pi}{6}$, not $\frac{2\pi}{6}$.

Find the quotient. Write it in rectangular form, using exact values. (3 points)

Problem: $\frac{\sqrt{7} + i\sqrt{21}}{-3\sqrt{21} - 3i\sqrt{7}}$

2. Convert top and bottom, divide the r 's, subtract the θ 's.

Top: $r = \sqrt{(\sqrt{7})^2 + (\sqrt{21})^2} = \sqrt{7+21} = \sqrt{28} = 2\sqrt{7}$ $\tan \theta = \frac{\sqrt{21}}{\sqrt{7}} = \sqrt{3} \Rightarrow \text{QI}, 60^\circ$

Bottom: $r = \sqrt{(-3\sqrt{21})^2 + (-3\sqrt{7})^2} = \sqrt{189+63} = \sqrt{252} = 6\sqrt{7}$ $\tan \theta = \frac{-3\sqrt{7}}{-3\sqrt{21}} = \frac{1}{\sqrt{3}} \Rightarrow \text{ref } 30^\circ, \text{ QIII} \Rightarrow 210^\circ$

$$\frac{2\sqrt{7} \operatorname{cis} 60^\circ}{6\sqrt{7} \operatorname{cis} 210^\circ} = \frac{1}{3} \operatorname{cis}(60^\circ - 210^\circ) = \frac{1}{3} \operatorname{cis}(-150^\circ) = \frac{1}{3} \operatorname{cis} 210^\circ$$

$$= \frac{1}{3} \left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i \right) = \boxed{-\frac{\sqrt{3}}{6} - \frac{1}{6}i}$$

Trap: both parts negative means QIII. $\tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$ says 30° ; the sketch says 210° . $\sqrt{252} = \sqrt{36 \cdot 7}$.

Find the power. Write the answer in rectangular form. (3 points)

Problem: $(-2 + 3i)^4$

3. De Moivre: $r^4 \operatorname{cis} 4\theta$. θ is not on the unit circle, so don't find it in degrees. Keep $\cos \theta = \frac{x}{r}$, $\sin \theta = \frac{y}{r}$ exact and use the double-angle formulas twice.

$$r = \sqrt{(-2)^2 + 3^2} = \sqrt{13}, \quad r^4 = 13^2 = 169 \quad \cos \theta = \frac{-2}{\sqrt{13}}, \quad \sin \theta = \frac{3}{\sqrt{13}} \quad (\text{QII})$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = \frac{4-9}{13} = -\frac{5}{13} \quad \sin 2\theta = 2 \sin \theta \cos \theta = \frac{2(3)(-2)}{13} = -\frac{12}{13}$$

$$\cos 4\theta = \cos^2 2\theta - \sin^2 2\theta = \frac{25-144}{169} = -\frac{119}{169} \quad \sin 4\theta = 2 \sin 2\theta \cos 2\theta = \frac{2(-12)(-5)}{169} = \frac{120}{169}$$

$$169 \left(-\frac{119}{169} + \frac{120}{169}i \right) = \boxed{-119 + 120i}$$

Trap: $r^4 = (\sqrt{13})^4 = 169$, and it cancels every denominator. $2(-12)(-5)$ is positive. Check: squaring twice, $(-2 + 3i)^2 = -5 - 12i$, then $(-5 - 12i)^2 = -119 + 120i$.

Find the fifth roots of the number. Leave answers in trig form with θ in radians. Graph each root as a vector in the complex plane. (3 points)

Problem: $16 - 16i\sqrt{3}$

4. Convert, then roots. Radians throughout.

$$r = \sqrt{16^2 + (-16\sqrt{3})^2} = \sqrt{256 + 768} = \sqrt{1024} = 32$$

$$\tan \theta = \frac{-16\sqrt{3}}{16} = -\sqrt{3} \Rightarrow \text{ref } \frac{\pi}{3}, \text{ QIV} \Rightarrow \frac{5\pi}{3}$$


$$\sqrt[5]{32} = 2 \quad \text{first: } \frac{5\pi/3}{5} = \frac{\pi}{3} = \frac{5\pi}{15}$$

$$\text{spacing: } \frac{2\pi}{5} = \frac{6\pi}{15}$$

$$2 \text{cis } \frac{\pi}{3}, 2 \text{cis } \frac{11\pi}{15}, 2 \text{cis } \frac{17\pi}{15}, 2 \text{cis } \frac{23\pi}{15}, 2 \text{cis } \frac{29\pi}{15}$$

Common denominator 15 makes the $+\frac{6\pi}{15}$ steps easy. Five roots, last one $\frac{29\pi}{15} < 2\pi$.

Trap: spacing is $\frac{2\pi}{n}$, not $\frac{2\pi}{3}$ or $\frac{\pi}{5}$.



One of the fourth roots of a complex number w is $-\sqrt{2} + i\sqrt{2}$. (3 points)

Problem: a) Determine the rectangular form of its other three fourth roots.

b) Find w in rectangular form.

5. All n th roots have the same r and sit $\frac{360^\circ}{n}$ apart. Find the given root's trig form, then step around.

$$-\sqrt{2} + i\sqrt{2}: r = \sqrt{2+2} = 2 \quad \tan \theta = -1, \text{ QII} \Rightarrow 135^\circ \quad \text{spacing } \frac{360^\circ}{4} = 90^\circ$$

$$\text{a) } 2 \text{cis } 225^\circ = 2 \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) = -\sqrt{2} - i\sqrt{2}$$

$$2 \text{cis } 315^\circ = 2 \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) = \sqrt{2} - i\sqrt{2}$$

$$2 \text{cis } 405^\circ = 2 \text{cis } 45^\circ = 2 \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) = \sqrt{2} + i\sqrt{2}$$

$$\text{b) } w = (2 \text{cis } 135^\circ)^4 = 2^4 \text{cis } 540^\circ = 16 \text{cis } 180^\circ = 16(-1 + 0i) = \boxed{-16}$$

$$\text{a) } -\sqrt{2} - i\sqrt{2}, \sqrt{2} - i\sqrt{2}, \sqrt{2} + i\sqrt{2} \quad \text{b) } w = -16$$

Trap: fourth roots are 90° apart, not 120° (that is cube roots). To get w , raise the root to the 4th with De Moivre.

*Non-Calculator***Honor Code (copy below & sign): On my honor, before God, I have neither given, nor received any aid on this quiz.****You have 30 minutes.**_____
Signature**SHOW ALL WORK FOR FULL CREDIT!**

Find the product. Write the product in rectangular form, using exact values. (3 points)

1. $\left[2\sqrt{2} \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) \right] \left[\sqrt{2} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) \right]$

Convert to trigonometric form, then divide. Write the quotient in rectangular form, using exact values. (3 points)

2. $\frac{-2 - 2i\sqrt{3}}{\sqrt{2} - i\sqrt{6}}$

Find the power. Write the answer in rectangular form, using exact values. (3 points)


3. $\left[2 \left(\cos \frac{7\pi}{12} + i \sin \frac{7\pi}{12} \right) \right]^8$

Solve. Write answers in rectangular form. (3 points)

4. $x^4 + 81 = 0$

Find the cube roots of the number. Leave answers in trig form. Graph each cube root as a vector in the complex plane. (3 points)

5. $2 - 2i\sqrt{3}$



ANSWER KEY – Non-Calculator

SHOW ALL WORK FOR FULL CREDIT!

Every line shown. Point splits are an estimate of partial credit (halves are possible).

Find the product. Write the product in rectangular form, using exact values. (3 points)

Problem: $\left[2\sqrt{2} \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) \right] \left[\sqrt{2} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) \right]$

1. Multiply the r 's, add the θ 's. The sum is $\frac{11\pi}{12} = 165^\circ$, which is *not* on the unit circle, so use the sum formula.

$$r = (2\sqrt{2})(\sqrt{2}) = 4 \quad \theta = \frac{2\pi}{3} + \frac{\pi}{4} = \frac{8\pi}{12} + \frac{3\pi}{12} = \frac{11\pi}{12}$$

$$\cos \frac{11\pi}{12} = \cos \frac{2\pi}{3} \cos \frac{\pi}{4} - \sin \frac{2\pi}{3} \sin \frac{\pi}{4} = \left(-\frac{1}{2}\right) \frac{\sqrt{2}}{2} - \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} = \frac{-\sqrt{2}-\sqrt{6}}{4}$$

$$\sin \frac{11\pi}{12} = \sin \frac{2\pi}{3} \cos \frac{\pi}{4} + \cos \frac{2\pi}{3} \sin \frac{\pi}{4} = \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} + \left(-\frac{1}{2}\right) \frac{\sqrt{2}}{2} = \frac{\sqrt{6}-\sqrt{2}}{4}$$

$$4 \left(\frac{-\sqrt{2}-\sqrt{6}}{4} + \frac{\sqrt{6}-\sqrt{2}}{4} i \right) = \boxed{-\sqrt{2} - \sqrt{6} + (\sqrt{6} - \sqrt{2})i}$$

Trap: $\cos(A+B) = \cos A \cos B - \sin A \sin B$ (minus), $\sin(A+B) = \sin A \cos B + \cos A \sin B$ (plus). QII check: $\cos$ negative, $\sin$ positive.

Convert to trigonometric form, then divide. Write the quotient in rectangular form, using exact values. (3 points)

Problem: $\frac{-2 - 2i\sqrt{3}}{\sqrt{2} - i\sqrt{6}}$

2. Convert top and bottom, divide the r 's, subtract the θ 's.

$$\text{Top: } r = \sqrt{(-2)^2 + (-2\sqrt{3})^2} = \sqrt{4+12} = 4 \quad \tan \theta = \frac{-2\sqrt{3}}{-2} = \sqrt{3} \Rightarrow \text{ref } 60^\circ, \text{ QIII} \Rightarrow 240^\circ$$

$$\text{Bottom: } r = \sqrt{(\sqrt{2})^2 + (-\sqrt{6})^2} = \sqrt{2+6} = \sqrt{8} = 2\sqrt{2} \quad \tan \theta = \frac{-\sqrt{6}}{\sqrt{2}} = -\sqrt{3} \Rightarrow \text{ref } 60^\circ, \text{ QIV} \Rightarrow 300^\circ$$

$$\frac{4 \text{cis } 240^\circ}{2\sqrt{2} \text{cis } 300^\circ} = \frac{2}{\sqrt{2}} \text{cis}(240^\circ - 300^\circ) = \sqrt{2} \text{cis}(-60^\circ) = \sqrt{2} \text{cis } 300^\circ$$

$$= \sqrt{2} \left(\frac{1}{2} - \frac{\sqrt{3}}{2}i \right) = \boxed{\frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i}$$

Trap: $\frac{2}{\sqrt{2}} = \frac{2\sqrt{2}}{2} = \sqrt{2}$. $\sqrt{2} \cdot \sqrt{3} = \sqrt{6}$. Both parts of the top are negative: QIII, 240° (not 60°).

Find the power. Write the answer in rectangular form, using exact values. (3 points)

Problem: $\left[2 \left(\cos \frac{7\pi}{12} + i \sin \frac{7\pi}{12} \right) \right]^8$

3. De Moivre: raise r to the 8th, multiply the angle by 8, then reduce by multiples of 2π .

$$2^8 = 256 \quad 8 \cdot \frac{7\pi}{12} = \frac{56\pi}{12} = \frac{14\pi}{3}$$

$$\frac{14\pi}{3} - 4\pi = \frac{14\pi}{3} - \frac{12\pi}{3} = \frac{2\pi}{3}$$

$$256 \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) = 256 \left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i \right) = \boxed{-128 + 128\sqrt{3}i}$$

Trap: $\frac{7\pi}{12}$ is not a unit-circle angle, but $8 \cdot \frac{7\pi}{12}$ reduces to one. Don't reach for the sum formula before multiplying.

Solve. Write answers in rectangular form. (3 points)

Problem: $x^4 + 81 = 0$

4. Isolate x^4 , write the constant in trig form, take fourth roots, convert each.

$$x^4 = -81 = -81 + 0i \Rightarrow r = 81, \theta = 180^\circ \quad x = (81 \text{ cis } 180^\circ)^{1/4}$$

$$\sqrt[4]{81} = 3 \quad \frac{180^\circ}{4} = 45^\circ \quad \text{spacing } 90^\circ \quad x = 3 \text{ cis } 45^\circ, 3 \text{ cis } 135^\circ, 3 \text{ cis } 225^\circ, 3 \text{ cis } 315^\circ$$

$$3 \text{ cis } 45^\circ = \frac{3\sqrt{2}}{2} + \frac{3\sqrt{2}}{2}i \quad 3 \text{ cis } 135^\circ = -\frac{3\sqrt{2}}{2} + \frac{3\sqrt{2}}{2}i$$

$$3 \text{ cis } 225^\circ = -\frac{3\sqrt{2}}{2} - \frac{3\sqrt{2}}{2}i \quad 3 \text{ cis } 315^\circ = \frac{3\sqrt{2}}{2} - \frac{3\sqrt{2}}{2}i$$

$$x = \pm \frac{3\sqrt{2}}{2} \pm \frac{3\sqrt{2}}{2}i \quad (\text{all four sign combinations})$$

Trap: $x = \pm 3i$ is wrong: $(3i)^4 = 81i^4 = 81$, not -81 . An even power of a pure imaginary is always positive real, so the roots sit between the axes ($45^\circ, 135^\circ, \dots$).

Find the cube roots of the number. Leave answers in trig form. Graph each cube root as a vector in the complex plane. (3 points)

Problem: $2 - 2i\sqrt{3}$

5. Convert, then cube roots. $\sqrt[3]{4}$ does not simplify, so leave it.

$$r = \sqrt{2^2 + (-2\sqrt{3})^2} = \sqrt{4 + 12} = 4$$


$$\tan \theta = \frac{-2\sqrt{3}}{2} = -\sqrt{3} \Rightarrow \text{ref } 60^\circ, \text{ QIV} \Rightarrow 300^\circ$$

$$\sqrt[3]{4} \quad \text{first: } \frac{300^\circ}{3} = 100^\circ \quad \text{spacing } 120^\circ$$

$$\sqrt[3]{4} \text{ cis } 100^\circ, \sqrt[3]{4} \text{ cis } 220^\circ, \sqrt[3]{4} \text{ cis } 340^\circ$$

The graph: three equal vectors of length $\sqrt[3]{4}$ (between 1 and 2), 120° apart (the “peace sign”).

Trap: divide the angle, not the radius: $\sqrt[3]{4}$, not $\frac{4}{3}$. All angles stay positive and under 360° .



three equal vectors, 120° apart