



how is the famous house problem related to co



AI Mode

All

Images

Videos

Short videos

Forums

Shopping

More ▾

Tools ▾

AI Overview



The famous house problem (finding house number 'x' on a street 'n' where left sum equals right sum) connects to continued fractions because its solutions form a pattern related to convergents of the continued fraction for $\sqrt{8}$, specifically $[2; 1, 4...]$, revealing infinite solutions to a related Pell equation, which Ramanujan instantly recognized and solved by extracting values from these convergents, giving integer pairs for (x, n). [🔗](#)

$$2 + \frac{1}{2 + \frac{1}{3 + \frac{1}{1 + \frac{1}{3 + \frac{1}{4 + \frac{1}{5 + \dots}}}}}}$$

The Problem Setup


- The Scenario:** Houses numbered 1 to n on a street, with a house x where $\sum_{i=1}^{x-1} i = \sum_{i=x+1}^n i$.
- The Equation:** This balances to $x(x-1)/2 = (n(n+1)/2) - (x(x+1)/2)$, simplifying to $2x^2 = n(n+1)$ or $2x^2 - n^2 - n = 0$. [🔗](#)

Ramanujan's Insight: The Pell Equation Connection

- Transforming the Equation:** The equation $2x^2 - n^2 - n = 0$ can be manipulated into a Pell-like equation by multiplying by 8: $16x^2 - 8n^2 - 8n = 0$, then adding 2: $(4x)^2 - (2n+1)^2 = -1$ (or $(4x)^2 - 2(2n+1)^2 = -1$ after some steps), leading to $(2x)^2 - 2(n+1/2)^2 = 1/2$, ultimately relating to $\sqrt{2}$ or $\sqrt{8}$.
- The Role of $\sqrt{8}$:** Ramanujan saw this structure as related to approximating $\sqrt{8}$.
- Continued Fraction Expansion of $\sqrt{8}$:** $\sqrt{8} = 2 + 1/(1 + 1/(4 + 1/(1 + 1/(4 + \dots))))$, written as $[2; 1, 4...]$. [🔗](#)

How Solutions Emerge

- Convergents:** Truncating the infinite continued fraction at each step gives "convergents" (fractions like $2/1$, $5/2$, $28/11$, etc.).



interpreted correctly.

3. **Example:** Convergents of $\sqrt{8}$ give pairs like $(x=1, n=1)$, $(x=6, n=8)$, $(x=35, n=49)$, $(x=204, n=288)$ – each representing valid house number solutions, with the numbers in the denominator (like 1, 8, 49, 288) often being the total house count n or related values. [↗](#)

In essence, Ramanujan recognized the problem as a quest for specific rational approximations of $\sqrt{8}$, a classic number theory topic beautifully solved by continued fractions, providing an elegant, infinite family of answers instead of just one. [↗](#)

Dive deeper in AI Mode



The Scientific Teen

<https://www.thescientificteen.org> › post › ramanujan-th...

RAMANUJAN: The Untrained Genius [↗](#)



Medium · Shenan Piyumantira

2 likes · 3 months ago

Ramanujan's Door Number Problem — From Street ...

Continued fractions give the best rational approximations to irrational numbers. Every truncated continued fraction is called a convergent. These ...

People also ask :

What is an example of a continued fraction?



What's the answer to the viral math problem?



Did Ramanujan work on continued fractions?



What is continued as a fraction?



[Feedback](#)



Mathematics Stack Exchange

<https://math.stackexchange.com> › questions › what-was...

What was Ramanujan's solution to the House Number ...

May 19, 2012 — The problem reduces to a Pell equation whose solutions can be obtained from the continued fraction expansion of $\sqrt{2}$.

[4 answers](#) · Top answer: An expansion of Andre's comment into a detailed exposition can be found at ht...

How can this **problem** be solved using **continued** ... [2 answers](#) Dec 22, 2016

Continued Fraction Counting Problem - Math Stack ... [2 answers](#) Aug 3, 2015

[More results from math.stackexchange.com](#)



WordPress.com

<https://infinitesimallysmallcom.wordpress.com> › the-ho...

The House Number Problem - The Other Side Of Mathematics

Apr 24, 2022 — “Please take down the solution” Ramanujan said and went on to dictate a **continued fraction**. This was the solution to the whole class of such ...



Bhāvanā – The mathematics magazine

<https://bhavana.org.in> › timeless-genius-es-celestial-cloc...

Timeless Geniuses, Celestial Clocks and Continued ...



thefederal.com

<https://thefederal.com> › Features

Ramanujan's enchanting journeys in the maze of 'continued ...

Dec 21, 2021 — The continued fraction provides an easy way of computing the increasingly better approximate value for irrational numbers.



YouTube · Mathologer

913.6K+ views · 5 years ago

How did Ramanujan solve the STRAND puzzle?



He promptly answered 'Please take down the solution' and dictated a continued fraction. The first term was the solution which I had obtained. ...

45:10

8 key moments in this video ▼



Quanta Magazine

<https://www.quantamagazine.org> › Puzzles

Solution: 'Puzzles Inspired by Ramanujan'

Aug 8, 2016 — The second method is based on **continued fractions**, the complete theory of which was perfected by Lagrange in the 18th century. This brings us ...



plus.maths.org

<https://plus.maths.org> › content › chaos-numberland-se...

Chaos in Numberland: The secret life of continued fractions

Jun 1, 2000 — **Continued fractions** are one of the most revealing representations of numbers. Numbers whose decimal expansions look unremarkable and featureless are revealed ...



University of Auckland

<https://www.math.auckland.ac.nz> › ~butcher › mi...

MATHEMATICAL MINIATURE 2 On Ramanujan^{PDF}, continued ...

For example, the convergents, formed by truncating the **continued fractions** at successive terms, are. $1 = 1.1. ., 1 + \frac{1}{2} = 3.2. ., 1 + \frac{1}{2 + \frac{1}{2}} = 7.5.$

People also search for

How is the famous house problem related to continued fractions...



Ramanujan house problem





1 2 3 4 5 6 7 8 9 10 [Next](#)

Results are personalized - [Try without personalization](#)

Newington, Connecticut - [From your IP address](#) - [Update location](#)

[Help](#) [Send feedback](#) [Privacy](#) [Terms](#)