

PDF file name: Elsevier-TheGreenJournal-Prognostic importance of radiologic extranodal

Original article word count: 3130

Extracted content word count: 176 (5% of the original word count)

Title of the article: *Prognostic importance of radiologic extranodal extension in nasopharyngeal carcinoma treated in a Canadian cohort*

The content extracted is as follows:

Methods

The performance of OS and DFS prediction was evaluated in all cases (both cN0 and cN+) in the overall TNM staging using our cN classification proposal, as well as those proposed by Lu et al [6] and Mao et al [5], and compared with the current TNM-8 stage grouping schema based on well-established criteria using OS as the primary end-point and DFS as a secondary end-point [18–21]: (1).

Results

C-index and AIC for Lu [6] and Mao's [5] proposal was improved slightly versus TNM-8 cN classification (c-index: 0.729 vs 0.706 vs 0.704; AIC: 535.60 vs 544.94 vs 545.30).

Discussion

Proposal-I classifies all unequivocal rENE as cN3, which appears to be the simplest approach while achieving the top overall ranking for both OS and DFS compared to other proposals.

Lu et al. [6] separated G2-rENE and G3-rENE in their proposal due to differential HRs associated with G2-rENE and G3-rENE.

Both proposals outperformed TNM-8 and other staging schemas [5,6] that included rENE in cN classification for OS.