

Outcomes after Liver Transplantation in Korea: Incidence and Risk Factors from Korean Transplantation Registry

Jong Man Kim^{¶1}, Deok Gie Kim^{¶2}, Jihyun Kim³, Keunsung Lee³, Kwang-Woong Lee⁴, Je Ho Ryu⁵, Bong-Wan Kim⁶, Dong Lak Choi⁷, Young Kyoung You⁸, Dong-Sik Kim⁹, Yang Won Nah¹⁰, Koo Jeong Kang¹¹, Jai Young Cho¹², Geun Hong¹³, Hee Chul Yu¹⁴, Ju Ik Moon¹⁵, Dongho Choi¹⁶, Shin Hwang¹⁷ and Myoung Soo Kim¹⁸

¹ Department of Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea

² Department of Surgery, Yonsei University Wonju College of Medicine, Wonju, South Korea

³ Novartis Korea Ltd., Seoul, South Korea

⁴ Department of Surgery, Seoul National University College of Medicine, South Korea

⁵ Department of Surgery, Division of Hepato-Biliary-Pancreatic Surgery and liver Transplantation, Pusan National University School of Medicine, Pusan National University Yangsan Hospital, South Korea

⁶ Department of Liver Transplantation and Hepatobiliary Surgery, Ajou University School of Medicine, South Korea

⁷ Department of Surgery, Catholic University of Daegu College of Medicine, South Korea

⁸ Department of Surgery, College of Medicine, The Catholic University of Korea, South Korea

⁹ Division of HBP Surgery & Liver Transplantation, Department of Surgery, Korea University College of Medicine, South Korea

¹⁰ Department of Surgery, Ulsan University Hospital, University of Ulsan College of Medicine, South Korea

¹¹ Department of Surgery, Dongsan Medical Center, Keimyung University School of Medicine, South Korea

¹² Department of Surgery, Seoul National University College of Medicine, South Korea

¹³ Department of Surgery, Mokdong Hospital, Ewha Womans University School of Medicine, South Korea

¹⁴ Department of Surgery, Jeonbuk National University Medical School, South Korea

¹⁵ Department of Surgery, Konyang University Hospital, South Korea

¹⁶ Department of Surgery, Hanyang University, South Korea

¹⁷ Department of Surgery, College of Medicine University of Ulsan, Asan Medical Center, South Korea

¹⁸ Department of Surgery, Yonsei University College of Medicine, Seoul, South Korea

[¶] Jong Man Kim and Deok Gie Kim equally contributed to manuscript as co-1st author

Corresponding author:

Professor Myoung Soo Kim, MD, PhD

Department of Surgery,

Yonsei University College of Medicine

50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, South Korea

E-mail: ysms91@yuhs.ac

Orcid: <https://orcid.org/0000-0002-8975-8381>

and Professor Shin Hwang, MD, PhD

Department of Surgery,

College of Medicine University of Ulsan, Asan Medical Center

88, Olympic-ro 43-gil, Songpa-gu, Seoul 0505, South Korea

E-mail: shwang@amc.seoul.kr

Orcid: <https://orcid.org/0000-0002-9045-2531>

Running title: Outcomes after LT in Korea

Accepted

ABSTRACT

Background/Aims: To analyze the incidence and risk factors of outcomes after liver transplantation (LT) in the Korean population. **Methods:** This study analyzed data from the liver cohort of Korean Organ Transplantation Registry (KOTRY) who had LT between May 2014 and December 2017. Study measures included the incidence of post-LT outcomes in recipients of living donor LT (LDLT) and deceased donor LT (DDLT). Cox multivariate proportional hazards model was used to determine the potential risk factors predicting the outcomes. **Results:** A total of 2,563 adult recipients with LT (LDLT, n=1,956; DDLT, n=607) were included, with mean±SD age of 53.9±8.9 years, and 72.2% were male. The post-LT outcomes observed in each LDLT and DDLT recipients were death (4.0% and 14.7%), graft loss (5.0% and 16.1%), rejection (7.0% and 12.0%), renal failure (2.7% and 13.8%), new onset of diabetes (12.5% and 15.4%), and hepatocellular carcinoma (HCC) recurrence (both 6.7%). In both LDLT and DDLT recipients, the most common post-LT complications were renal dysfunction (33.6% and 51.4%), infection (26.7% and 48.4%), and surgical complication (22.5% and 23.9%). Incidence of these outcomes were generally higher among recipients of DDLT than LDLT. Multivariate analysis indicated recipient age and DDLT as significant risk factors associated with death and graft loss. DDLT and ABO incompatible transplant were prognostic factors for rejection, and HCC beyond Milan criteria at pre-transplant was a strong predictor of HCC recurrence. **Conclusions:** This study is a good indicator of the post-LT prognosis in the Korean population and suggests a significant burden of post-LT complications.

Keywords: Liver transplantation; Incidence; Risk factor; Renal dysfunction

Study highlights:

- In the Korean liver transplant population, incidence of graft loss, rejection and death were higher in recipients of deceased donor than living donor
- The most common outcomes occurring in more than 20% of recipients following liver transplantation were renal dysfunction, infection, and surgical complications
- The study highlights the burden of outcomes post liver transplantation, and identifies risk factors requiring special attention to improve prognosis

Accepted

INTRODUCTION

Liver diseases are a public health problem worldwide and are highly prevalent in Asian countries.¹ In Korea, liver diseases have a high prevalence along with diabetes mellitus (DM), respiratory diseases, and hypertension. According to the Korea Statistics Office, liver diseases accounted for 13.3 deaths per 100,000 population in Korea, and ranked the seventh leading cause of death in 2017.²

Liver transplantation (LT) is the gold standard treatment for patients with end-stage liver disease (ESLD). Since the first LT in Korea in 1988, the number of LT has increased dramatically over time.³ To meet such an increased demand, living donor LT (LDLT) has become an inevitable option as it shortens the waiting time and lowers the dropout rate in comparison to deceased donor LT (DDLT).^{4,5}

According to the Korean Network for Organ Sharing (KONOS), the number of LT performed annually in Korea increased more than seven-fold between 2000 (n=205) and 2019 (n=1,579). Of the total LT procedures in 2019, LDLT accounted for 1,188 cases (75.24%).⁶ The most common indication for LT was alcoholic liver disease (21.85%), followed by hepatitis B virus (HBV; 18.43%), malignant neoplasms (17.10%) and acute liver failure (4.05%).⁶ The survival rate reported in 2019 for LDLT was 94.63% at 3 months, 90.41% at 1 year and 81.01% at 5 years post-transplant. DDLT showed a relatively lower survival rate of 82.24%, 77.31% and 68.46% respectively.⁶

Despite advancement in LT surgery techniques and post-LT management, complications after LT are still frequent. Graft loss, rejection, renal dysfunction, and new onset diabetes mellitus (NODM) are well-known post-LT complications.⁷ Recipient and donor age, gender, and blood

group type matching have been identified as representative risk factors along with the need for appropriate management after LT to improve outcomes.⁸⁻¹²

Study on LT outcomes and associated risk factors in Korean population are still scarce. Identification of this data gap utilizing locally generated database will be worthy of understanding the current status of LT prognosis in Korea. The Korean Organ Transplantation Registry (KOTRY), which was established in 2014, is the first integrated national transplant registry in Korea that includes post-transplantation data from five solid organ transplantation cohorts, comprising of lung, kidney, heart, liver, and pancreas transplants.¹³

The objective of the present study was to investigate the incidence and risk factors of post-LT outcomes among LT recipients in Korea by leveraging the KOTRY registry database.

MATERIALS AND METHODS

Study design and data source

This study utilized data from the liver cohort of the KOTRY registry generated from 15 participating transplantation centers in 2015. These centers performed 81.5% of Korean LTs (1139/1398) in 2015.⁴ The liver cohort of KOTRY, a web-based clinical research management system (iCReaT; <http://icreat.nih.go.kr/> study number: C140015), collected data on both the recipient and living donor demographics, comorbidities, immunologic assessment and viral markers at enrollment. For deceased donors, donor profile on cause of brain death and cold ischemic time were collected at enrollment, and for LT recipients, additional data on laboratory assessment, immunosuppressant used, surgical complications, post-transplant outcomes and complications were collected at enrollment and at 1, 6, 12, 24, and 36 months post-transplant.

The KOTRY registry was approved by the Institutional Review Boards of each participating center. All patients provided informed consent prior to their registration in the KOTRY database.

Patient selection

The study population consisted of patients included in the KOTRY registry who had LT between May 1, 2014, and December 31, 2017. The details of KOTRY design and patient enrollment have been described in previous reports.^{4,13} LT recipients who had been followed up for at least six months post LT were included in the analysis. A total of 2,648 LT recipients were identified, of which 2,563 were adults and 85 were pediatrics. Given the heterogeneous characteristics of pediatric LT recipients, the sample size of pediatrics appeared too small to get statistically valid inferences. Thus, we analyzed the data on adults only.

Study measures

Study measures included recipient demographics (age, gender, body mass index [BMI]), comorbidities (hypertension and DM), underlying liver disease, Child-Turcotte-Pugh (CTP) score and Model for End-stage Liver Disease (MELD) score for chronic liver disease or cirrhosis, urgent status in Korean Network for Organ Sharing (KONOS)¹⁴, donor type (LD or DD), donor demographics (age, gender, body mass index [BMI]), graft related variables (graft type, graft-recipient weight ratio and type of donor operation), ABO blood group compatibility, laboratory parameters, and type of immunosuppressant used. If hepatocellular carcinoma (HCC) was the underlying cause for LT, the pre-transplant HCC status including Milan criteria and liver cancer free (no viable tumor) state was additionally inspected. If the donor type was DD, cold ischemic time and macrovesicular steatosis in donor were also examined. All of these variables were assessed to identify the risk factors on post-LT outcomes.

Post-LT outcomes

The post-LT outcomes examined were overall survival, graft survival, malignancies (HCC, non-HCC, or cholangiocarcinoma), graft rejection (acute or chronic), renal failure, renal dysfunction, infection, surgical complication and new onset of DM (NODM).

Graft loss was defined as patient death or re-transplantation. Rejection was identified as diagnosed clinically or biopsy-proven. Renal failure was defined as initiation of dialysis or receipt of kidney transplantation and renal dysfunction was defined as the eGFR <60 mL/min/1.73 m². Infection was recorded when the causative pathogen was isolated and the patient was treated with antibiotics. Surgical complication was noted as one of the following: bleeding at operation site, intra-abdominal abscess, vascular stenosis or thrombosis (hepatic artery, portal vein stenosis or hepatic vein), biliary stricture, or biliary stenosis. NODM was defined as use of insulin or glucose lowering agent in patients who were non-diabetic before transplantation.

Statistical analysis

Continuous variables are expressed as mean \pm standard deviation (SD) or median (range: minimum, maximum) and categorical variables are expressed as counts and percentages.

Survival curves of LDLT and DDLT according to post-LT outcomes were estimated by the Kaplan-Meier method and tested using log-rank test. The censoring time was defined as the final documented date. The date of death with no evidence of outcomes was also considered censoring except in case of outcome death.

To assess the potential risk factors influencing the occurrence of post-LT outcomes, a separate univariate cox regression analysis was conducted with each variable as an independent predictor for each outcome. Significant variables in the univariate analysis ($p < 0.05$) were included in a cox multivariate proportional hazards model to determine which factors independently predicted post-LT outcomes and graft or patient survival. Results obtained from multivariate analysis were expressed as hazard ratio (HR), 95% CI, and p -value. All tests were two-sided, and statistical significance was defined as $p < 0.05$. All analyses were performed using SAS version 9.2 (SAS Institute, NC, USA).

RESULTS

Baseline characteristics

2,563 adult recipients were included in the analysis, with a mean follow-up duration of 14.7 ± 10.6 months based on the data collected until June, 2018. 1,956 (76.3%) recipients had LDLT and 607 (23.7%) had DDLT. The mean \pm SD age of LDLT and DDLT recipients were 54.3 ± 8.4 and 52.6 ± 10.3 years, respectively. Recipient demographics and clinical characteristics according to donor type are summarized in Table 1.

Hepatitis B and alcoholic liver disease were the two most common underlying liver diseases among LDLT (60.5% and 20.5%) and DDLT (41.0% and 36.4%) recipients. Nearly half (49.2%) of all recipients had liver cancer, and HCC was accountable for majority (97.4%) of the neoplasms. Among recipients with HCC, about two-thirds (64.2% in LDLT and 65.7% in DDLT) had pre-transplant HCC within Milan criteria.

A higher proportion of DDLT than LDLT recipients had MELD score ≥ 35 (31.0% vs 3.0%), whereas more recipients of LDLT than DDLT had KONOS status score of 5 (81.4% vs 30.3%).

Left liver graft was most frequently used for LDLT (95.7%), while the whole liver graft was commonly used in DDLT (95.1%) (Table 1).

Donor characteristics

The mean age of overall LT donors was 35.3 ± 14.0 years and 63.9% were male. A small proportion of donors (27.1%) were overweight (BMI 25–29 kg/m²), and 7.2% had hypertension and 3.6% had DM. Living donors (n=1,956) were relatives of recipients in most cases (85.9%), and conventional operation (open surgery) was performed in 80.2% of them.

In deceased donors (n=607), the principal cause of death was progression of underlying disease (45.5%), followed by trauma (32.0%) and suicide (13.3%). The most common etiological causes of death were subarachnoid hemorrhage (33.3%), hypoxic damage (31.5%), and intracranial hemorrhage (26.7%). The mean cold ischemic time was 5.0 ± 4.4 hours. Macrovesicular steatosis was < 10% in majority (84.9%) of the donors and 10%–30% in 13.0% of the donors (data not shown and can be shared upon request to authors).

Treatments after LT

Induction of immunosuppression using steroids or basiliximab was performed in 89.4% and 77.1% of LT recipients, respectively. To maintain immunosuppression, most recipients (97.9%) were given calcineurin inhibitors (CNIs; eg, cyclosporin and tacrolimus), followed by antimetabolite agents (eg, azathioprine, mycophenolate mofetil, and mizoribine; 73.8%) and mammalian target of rapamycin (mTOR) inhibitors (eg, sirolimus and everolimus; 11.0%).

For the prophylaxis of HBV post LT, almost all (98.7%) recipients were prescribed anti-HBV-specific immunoglobulins (HBIG) either as monotherapy (34.1%) or in combination with antiviral therapies (64.6%).

Outcomes after LT

Incidence of death was 4.0% in LDLT group and 14.7% in DDLT group (Figure 1). Graft loss was observed in 5.0% and 16.1% of LDLT and DDLT recipients, respectively. The leading causes of graft loss were death of patient with functioning graft (69.7%), primary non-function of the graft (10.3%) and recurrence of HCC (5.6%) (Figure 2).

Seven percent of LDLT and 12.0% of DDLT recipients experienced rejection. In both LDLT and DDLT recipients, the most common (>20%) post-LT complications were renal dysfunction (33.6% and 51.4%), infection (26.7% and 48.4%), and surgical complication (22.5% and 23.9%).

Incidence of renal failure (2.7% and 13.8%), NODM (12.5% and 15.4%), and HCC recurrence (both 6.7%) were generally low in both LDLT and DDLT groups. Except for NODM and HCC recurrence, the incidence rate of post-LT outcomes was higher in recipients of DDLT than LDLT (Figure 1).

Kaplan-Meier survival curves for each outcome are presented in Figure 3. Survival rate of LDLT recipients were significantly higher than DDLT recipients during the first 3 years post LT (3-year, 90.4% vs 80.4%; $p<0.001$). Similar differences were observed for graft survival (3-year, 89.1% vs 79.2%; $p<0.001$) and rejection free survival (3-year, 90.2% vs 83.3%; $p<0.001$) between the two groups. Event-free survival rates for renal failure (3-year, 95.6% vs 78.3%; $p<0.001$), renal dysfunction (3-year, 50.5% vs 31.1 %; $p<0.001$) and infection (3-year, 64.1% vs 44.2%; $p<0.001$) were also significantly higher in LDLT recipients than DDLT. The survival distribution of

patients not developing NODM were significantly different among LDLT and DDLT (3-years, 83.3% vs 84.0%, $p=0.039$) and LDLT was higher than DDLT. No significant difference between LDLT and DDLT was observed for HCC recurrence free survival ($p=0.527$) and surgical complication ($p=0.538$).

Risk factors for post-LT outcomes

In both univariate and multivariate analysis, increased recipient age (each $p<0.001$) and DDLT (each $p<0.001$) were significantly associated with increased risk of death. Recipient MELD score ≥ 35 ($p<0.001$), underlying acute hepatitis ($p=0.004$), and primary liver disease ($p=0.001$) were identified as independent risk factors of death only in univariate analysis (Supplementary Table 1).

Multivariate analyses showed that increased recipient age ($p=0.012$), DDLT ($p=0.001$), underlying HCC ($p=0.011$), and acute hepatitis ($p=0.044$) were predisposing factors for graft loss after LT (Supplementary Table 2). HCC beyond the Milan criteria at pre-transplant was the only significant factor identified to increase the risk of HCC recurrence post liver transplant ($p<0.001$; Supplementary Table 3). DDLT ($p=0.042$) and ABO incompatible transplant ($p=0.002$) were independent factors associated with increased risk of rejection (Supplementary Table 4). MELD score ≥ 35 ($p<0.001$) was a predictor of increased risk of renal dysfunction, along with increased recipient age ($p<0.001$), DDLT ($p=0.004$), hypertension ($p=0.007$), and DM ($p=0.011$) in multivariate analysis (Supplementary Table 6). Multiple Cox regression analysis of risk factors for renal failure, infection, surgical complication and NODM are presented in Supplementary Tables 5 and 7-9.

DISCUSSION

This study showed the outcomes and their risk factor in patients who underwent LT. Renal dysfunction is the most common post-transplantation outcome in LDLT and DDLT, followed by infections, and surgical complications, while NODM, graft rejection, graft loss, HCC recurrence, and renal failure were other outcomes. With the exception of NODM, the incidence rate of these outcomes was greater among DDLT recipients compared with LDLT recipients. The age of recipient, MELD score, LDLT, underlying liver disease, Milan criteria were representative risk factors for the outcomes.

In the present study, incidence of death during follow-up duration was 6.5% for all recipients (4.0% in LDLT and 14.7% in DDLT) and the survival rates at 1, 2 and 3 years after transplantation were notably higher in LDLT recipients than DDLT. The survival rates at 3-year observed in our study were consistent with recent studies showing superior survival in LDLT recipients than DDLT.^{15,16} In alignment with literature, our study also supports that recipient age is a risk factor for survival in LT recipients.¹⁵⁻¹⁷ We observed that LDLT was associated with lower risk of death compared to DDLT. Several studies have shown better survival outcomes among recipients with LDLT than DDLT^{15,16,18}, indicating this as a preferable option for LT. Evidence suggests that problem of organ shortage for transplantation in Asia has been successfully addressed by increasing the number of LDLT. However, in Western countries, the number of LDLTs have remained stagnant for over a decade, and the demand for DDLT continues to increase.¹⁹ In Asian countries, high incidence of HCC and low organ donation rate from the deceased has evolved LDLT as a new emerging treatment modality for HCC.²⁰

We observed graft loss in 7.6% of our study population (5.0% in LDLT and 16.1% in DDLT). The death of the LT recipient was the most common cause of graft loss, which is consistent with findings from literature.²¹ A study by Olthoff et al. showed recipient age, hepatitis C virus and

HCC as the risk factors for graft loss²². The study by Azzam demonstrated ABO-incompatible, the age and sex of the donor, hepatitis B and C virus, HCC as the independent risk factors for graft loss.¹⁷ Our study revealed that increased age of the recipient, DDLT, HCC, acute hepatitis, use of steroids, and use of anti-metabolites were risk factors for graft loss, which are in alignment with several studies published previously.²²⁻²⁴ Thus, it also becomes important that recipients' characteristics, including the underlying liver disease should be considered during management.

In this study, the recurrence of HCC was observed in 6.7% each of LDLT and DDLT recipients. A study by Sandhu et al reported the low and similar rates of HCC recurrence at 3-year among recipients of LDLT and DDLT (10.7% and 14.8%).²⁵ Previous research by Park et al reported the low occurrence of de novo malignancies in 2.3% of Korean patients after a mean post-LT period of 41 months.²⁶ Another study by Hong et al. reported the recurrence of HCC in 24.5% of the recipients at 5 years.²⁷ Based on the Milan criteria, Lee et al. reported that the recurrence of HCC was 3.3% in patients within Milan and 29.4% in patients beyond Milan.²⁸ Our study also supports this observation, as the multivariate analysis revealed that the risk of HCC recurrence was higher among patients with HCC beyond the Milan criteria compared to those within the criteria. In addition to the Milan criteria, other factors such as male gender, tumor size of more than 5 cm, positive positron emission tomography scan, high serum alpha-fetoprotein level (>1000 ng/mL), and poor histologic grade (Edmonson-Steiner grade III and IV) have also been suggested to be associated with higher recurrence of HCC in Korean studies.²⁶⁻²⁸

Despite the progress made in preoperative evaluation, surgical procedures, postoperative care, and immunosuppressant therapy, chronic renal disease remains a common complication post LT.^{29,30} Approximately 37.8% of all LT recipients (33.6% in LDLT and 51.4% in DDLT) in the

present study were observed to have some degree of renal dysfunction post-transplantation, which is consistent with finding from earlier studies in Korea.^{31,32} A study by Lee et al. evaluating the early postoperative renal dysfunction in LDLT showed that 34.9% of recipients had renal dysfunction (defined as the creatinine level ≥ 1.5 mg/dL) at 1-year post LT.³² In another study by Kim et al., 43.5% of the overall LT recipients were reported to have moderate renal dysfunction (defined as the eGFR ≥ 30 – <60 mL/min/1.73 m²).³¹ The rate of renal dysfunction post-LT varies considerably among recipients in other countries, with 8.0% in Egypt,³³ 48.0% in Spain³⁴ and 46.7% in Brazil.³⁵ This disparity in post-LT renal dysfunction rates between these studies might arise from differences in pre-LT renal function status, source of graft (LDLT or DDLT), presence of diabetes/hypertension, post-LT care, timing of kidney function monitoring, or definitions used for renal dysfunction. Thus, there is a need to standardize these definitions to enable better appraisal of studies worldwide.

The MELD scoring system has been reported as a validated tool that predicts the renal failure post LT.³⁶ In consistency with previous research,^{32,36} the pre-transplant MELD score ≥ 35 was found to be a strong predictor of both renal failure and renal dysfunction in our study. Other significant risk factors such as increased recipient age and comorbid hypertension as predictor of renal dysfunction post LT were also in line with literature.²⁹ The occurrence of renal dysfunction post LT leads to poor prognosis. We observed an increasing trend of renal dysfunction rates with longer follow-ups (44.2%, 50.2% and 54.8% at 1, 2 and 3 years, respectively). Since renal dysfunction can increase the need for renal replacement therapy and also increase the risk of cardiovascular diseases, preventive measures are warranted for better prognosis.^{37,38}

The rate of NODM in the current study was observed to be approximately 14.8%. In a meta-analysis based on 56 studies, NODM was reported in 13.4% of patients following

transplantation.³⁹ We found that pre-transplant obesity ($\text{BMI} \geq 25 \text{ kg/m}^2$), use of steroids, and anti-metabolites were associated with a higher risk of NODM post LT. These results are in alignment with recent results published by KOTRY on NODM in the same population.⁴⁰

In order to improve the outcomes after LT, appropriate surgical technique, donor-recipient matching and optimization of immunosuppressants is essential. Also, there is a need for further research on innovative management strategies such as machine perfusion, liver support device and tissue bioengineering.⁴¹

Our study has several limitations. First, this study was a retrospective we analyzed registry database and cautious interpretation is required since the severity of each patient in database is different and interventional adjustment was not possible. Second, the KOTRY is a relatively new registry with short follow-up period, so long-term outcomes, including median survival time, could not be assessed. Third, it was difficult to compare the effects of medications because the patient population using these medications was heterogeneous. The use of immunosuppressants as the risk factor for the occurrence of post LT outcomes could not be ascertained because patients' status were different for choosing an immunosuppressant agent which may lead to different outcomes. Lastly, in case of HCC recurrence, the use of mTOR inhibitors has the advantage of lowering the post-transplant malignancy compared with other immunosuppressants; however, the study showed that the use of an mTOR inhibitor at 6 months of transplantation poses a higher risk. Given that the order of the incidents could not be confirmed, the use of immunosuppressant was not considered as one of the risk factors.

Despite of these limitations, this study is a good indicator of the outcomes after transplantation in Korean LT recipients. Following liver transplantation, recipients of deceased donor reported

higher rates of graft loss, rejection and death and the most frequent complications were renal dysfunction, infection, and surgical complications. Risk factors associated with poor outcomes warrant special attention to improve patient prognosis.

ACKNOWLEDGMENTS

This research was supported by a fund by Research of Korea Centers for Disease Control and Prevention (2014-ER6301-00, 2014-ER6301-01, 2014-ER6301-02, 2017-ER6301-00, 2017-ER6301-01, 2017-ER6301-02) and Novartis Korea Ltd., South Korea. Raju Gautam (Novartis Healthcare Pvt. Ltd, Hyderabad, India) provided medical writing assistance for the manuscript.

AUTHOR CONTRIBUTIONS

JM Kim, DG Kim, MS Kim and S Hwang had full access to all aspects of the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. JM Kim and MS Kim participated in the research design. DG Kim, S Hwang and MS Kim participated in the data acquisition. JM Kim and DG Kim participated in the statistical analysis. JH Kim contributed to study design, study execution, contributed to manuscript revision, and administrative support. KS Lee supported data analysis and study contributed to manuscript revision. The other authors participated in the performance of the research. MS Kim and S Hwang supervised the study process.

CONFLICT OF INTEREST

JM Kim, DG Kim and MS Kim received grants from Novartis Korea Ltd., South Korea for the conduct of this study. KS Lee employees of Novartis Korea Ltd., South Korea. JH Kim was an employee of Novartis Korea Ltd., South Korea at the time of conduct of this study, and left.

Authors declare that they have no other conflicts of interest regarding the content of this manuscript.

REFERENCES

1. Wong MCS, Huang JLW, George J, et al. The changing epidemiology of liver diseases in the Asia-Pacific region. *Nature reviews Gastroenterology & hepatology* 2019;16:57-73.
2. Statistics Korea. Causes of Death Statistics in 2017 [Accessed 2019 May 24]. Available from: <http://www.kostat.go.kr>. 2018.
3. Lee SG, Moon DB, Hwang S, et al. Liver transplantation in Korea: past, present, and future. *Transplantation proceedings* 2015;47:705-8.
4. Lee JG, Lee KW, Kwon CHD, et al. Donor safety in living donor liver transplantation: The Korean organ transplantation registry study. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2017;23:999-1006.
5. Park GC, Song GW, Moon DB, Lee SG. A review of current status of living donor liver transplantation. *Hepatobiliary surgery and nutrition* 2016;5:107-17.
6. Korean Network for Organ Sharing. Organ Transplantation Statistics. 2019. (Accessed August, 2020, at <http://konos.go.kr>.)
7. Moreno R, Berenguer M. Post-liver transplantation medical complications. *Annals of hepatology* 2006;5:77-85.
8. Adam R, Karam V, Delvart V, et al. Evolution of indications and results of liver transplantation in Europe. A report from the European Liver Transplant Registry (ELTR). *Journal of hepatology* 2012;57:675-88.

9. Bachir NM, Larson AM. Adult liver transplantation in the United States. *The American journal of the medical sciences* 2012;343:462-9.
10. Durand F. How to improve long-term outcome after liver transplantation? *Liver international : official journal of the International Association for the Study of the Liver* 2018;38 Suppl 1:134-8.
11. Durand F, Levitsky J, Cauchy F, Gilgenkrantz H, Soubrane O, Francoz C. Age and liver transplantation. *Journal of hepatology* 2019;70:745-58.
12. Sarkar M, Watt KD, Terrault N, Berenguer M. Outcomes in liver transplantation: does sex matter? *Journal of hepatology* 2015;62:946-55.
13. Yang J, Jeong JC, Lee J, et al. Design and Methods of the Korean Organ Transplantation Registry. *Transplantation direct* 2017;3:e191.
14. Lee J, Lee JG, Jung I, Joo DJ, Kim SI, Kim MS. Development of a Korean Liver Allocation System using Model for End Stage Liver Disease Scores: A Nationwide, Multicenter study. *Scientific reports* 2019;9:7495.
15. Gil E, Kim JM, Jeon K, et al. Recipient Age and Mortality After Liver Transplantation: A Population-based Cohort Study. *Transplantation* 2018;102:2025-32.
16. Humar A, Ganesh S, Jorgensen D, et al. Adult Living Donor Versus Deceased Donor Liver Transplant (LDLT Versus DDLT) at a Single Center: Time to Change Our Paradigm for Liver Transplant. *Annals of surgery* 2019;270:444-51.
17. Azzam A. Risk factors of hepatic graft failure, morbidity and mortality after living donor liver transplantation (LDLT): Review Article. *Clin Res Trials* 2018;4:1-11.

18. Zhu B, Wang J, Li H, Chen X, Zeng Y. Living or deceased organ donors in liver transplantation for hepatocellular carcinoma: a systematic review and meta-analysis. *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2019;21:133-47.
19. Jadlowiec CC, Taner T. Liver transplantation: Current status and challenges. *World journal of gastroenterology* 2016;22:4438-45.
20. Yoon YI, Lee SG. Living Donor Liver Transplantation for Hepatocellular Carcinoma: An Asian Perspective. *Digestive diseases and sciences* 2019;64:993-1000.
21. Bäckman L, Gibbs J, Levy M, et al. Causes of late graft loss after liver transplantation. *Transplantation* 1993;55:1078-82.
22. Olthoff KM, Abecassis MM, Emond JC, et al. Outcomes of adult living donor liver transplantation: comparison of the Adult-to-adult Living Donor Liver Transplantation Cohort Study and the national experience. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2011;17:789-97.
23. Gwiasda J, Schrem H, Klempnauer J, Kaltenborn A. Identifying independent risk factors for graft loss after primary liver transplantation. *Langenbeck's archives of surgery* 2017;402:757-66.
24. Sharma R, Kashyap R, Jain A, et al. Surgical complications following liver transplantation in patients with portal vein thrombosis--a single-center perspective. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract* 2010;14:520-7.
25. Sandhu L, Sandroussi C, Guba M, et al. Living donor liver transplantation versus deceased donor liver transplantation for hepatocellular carcinoma: comparable survival and

recurrence. Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society 2012;18:315-22.

26. Park HW, Hwang S, Ahn CS, et al. De novo malignancies after liver transplantation: incidence comparison with the Korean cancer registry. Transplantation proceedings 2012;44:802-5.

27. Hong SK, Lee KW, Kim HS, Yoon KC, Yi NJ, Suh KS. Living donor liver transplantation for hepatocellular carcinoma in Seoul National University. Hepatobiliary surgery and nutrition 2016;5:453-60.

28. Lee S, Ahn C, Ha T, et al. Liver transplantation for hepatocellular carcinoma: Korean experience. Journal of hepato-biliary-pancreatic sciences 2010;17:539-47.

29. Bahirwani R, Reddy KR. Outcomes after liver transplantation: chronic kidney disease. Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society 2009;15 Suppl 2:S70-4.

30. Wiesen P, Massion PB, Joris J, Detry O, Damas P. Incidence and risk factors for early renal dysfunction after liver transplantation. World journal of transplantation 2016;6:220-32.

31. Kim SG, Kim HJ, Lee JP, et al. Incidence and risk factors of renal dysfunction after liver transplantation in Korea. Transplantation proceedings 2004;36:2318-20.

32. Lee SK, Park JB, Kim SJ, et al. Early postoperative renal dysfunction in the adult living donor liver transplantation. Transplantation proceedings 2007;39:1517-9.

33. Abdel-Khalek EE, Alrefaey AK, Yassen AM, et al. Renal Dysfunction after Living-Donor Liver Transplantation: Experience with 500 Cases. Journal of transplantation 2018;2018:5910372.

34. Cabezuelo JB, Ramirez P, Rios A, et al. Risk factors of acute renal failure after liver transplantation. *Kidney international* 2006;69:1073-80.
35. Barreto AG, Daher EF, Silva Junior GB, et al. Risk factors for acute kidney injury and 30-day mortality after liver transplantation. *Annals of hepatology* 2015;14:688-94.
36. Machicao VI, Srinivas TR, Hemming AW, et al. Impact of implementation of the MELD scoring system on the prevalence and incidence of chronic renal disease following liver transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* 2006;12:754-61.
37. Gonwa TA, Mai ML, Melton LB, et al. Renal replacement therapy and orthotopic liver transplantation: the role of continuous veno-venous hemodialysis. *Transplantation* 2001;71:1424-8.
38. Weiner DE, Tighiouart H, Elsayed EF, et al. The relationship between nontraditional risk factors and outcomes in individuals with stage 3 to 4 CKD. *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2008;51:212-23.
39. Heisel O, Heisel R, Balshaw R, Keown P. New onset diabetes mellitus in patients receiving calcineurin inhibitors: a systematic review and meta-analysis. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* 2004;4:583-95.
40. Kim JM, Hwang S, Lee KW, et al. New-onset diabetes after adult liver transplantation in the Korean Organ Transplantation Registry (KOTRY) study. *Hepatobiliary surgery and nutrition* 2020;9:425-39.

41. Bodzin AS, Baker TB. Liver Transplantation Today: Where We Are Now and Where We Are Going. Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society 2018;24:1470-5.

Accepted

Table 1: Demographics and clinical characteristics of liver transplant recipients overall and by donor types


Variables	Missing	Total (N=2563)	LDLT (N=1956)	DDLT (N=607)
Age, yr	0	53.9±8.9	54.3±8.4	52.6±10.3
Sex, male	0	1850 (72.2)	1441 (73.7)	409 (67.4)
BMI ≥ 25 kg/m ²	2	937 (36.6)	725 (37.1)	212 (34.9)
Underlying liver disease	0			
Hepatitis B		1433 (55.9)	1184 (60.5)	249 (41.0)
Alcoholic		622 (24.3)	401 (20.5)	221 (36.4)
Hepatitis C		179 (7.0)	133 (6.8)	46 (7.6)
Cryptogenic		138 (5.4)	111 (5.7)	27 (4.4)
Others		191 (7.5)	127 (6.5)	64 (10.5)
Acute hepatitis	0	109 (4.3)	52 (2.7)	57 (9.4)
Liver cancer		1262 (49.2)		
HCC	0	1229 (48.0)	1056 (54.0)	173 (28.5)
CCC		11 (0.4)	—	—
Combined HCC + CC		20 (0.8)	—	—
Other malignancy		2 (0.1)	—	—
Pretransplant HCC status	3			
Within Milan criteria		790 (30.8)	677 (34.6)	113 (18.6)
Beyond Milan criteria		212 (8.3)	176 (9.0)	36 (5.9)
No viable tumor		224 (8.7)	201 (10.3)	23 (3.8)
Incidental liver cancer		0 (0.0)	0 (0.0)	0 (0.0)
Hypertension	0	454 (17.7)	363 (18.6)	91 (15.0)
Diabetes mellitus	0	629 (24.5)	480 (24.5)	149 (24.5)
MELD score (N=2562)	1			
<15		1319 (51.5)	1234 (63.1)	85 (14.0)
15–34		996 (38.9)	662 (33.9)	334 (55.0)
≥35		247 (9.6)	59 (3.0)	188 (31.0)
CTP classification	0			
A (≤6)		742 (29.0)	723 (37.0)	19 (3.1)
B (7–9)		860 (33.6)	769 (39.3)	91 (15.0)
C (≥10)		961 (37.5)	464 (23.7)	497 (81.9)

Variables	Missing	Total (N=2563)	LDLT (N=1956)	DDLT (N=607)
KONOS status	0			
Score 1		58 (2.2)	9 (0.5)	49 (8.1)
Score 2		163 (6.4)	35 (1.8)	128 (21.1)
Score 3		150 (5.9)	58 (3.0)	92 (15.2)
Score 4		416 (16.2)	262 (13.4)	154 (25.4)
Score 5		1776 (69.3)	1592 (81.4)	184 (30.3)
ABO incompatible	0	408 (15.9)	404 (20.7)	4 (0.7)
Graft type	0			
Whole liver		577 (22.5)	0 (0.00)	577 (95.1)
Right liver graft		87 (3.4)	85 (4.3)	2 (0.3)
Left liver graft		1899 (74.1)	1871 (95.7)	28 (4.6)
Graft-recipient weight ratio for LDLT < 0.8	33	–	159 (8.3)	–
Type of donor operation in LDLT (N=1956)	0			
Minimal invasive surgery		–	388 (19.8)	–
Conventional (open)		–	1568 (80.2)	–
Use of steroids		2288 (89.3)	1767(90.3)	521(85.8)
Use of anti-metabolites		1892 (73.8)	1485(75.9)	407(67.1)
Use of mTOR inhibitors		282 (11.0)	212(10.8)	70(11.5)

Data are presented as number (%) or mean±SD.

BMI, body mass index; CTP, Child-Turcotte-Pugh; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; CCC, cholangiocarcinoma; KONOS, Korean Network for Organ Sharing; LDLT, living donor liver transplantation; LT, liver transplantation; MELD, Model for End-Stage Liver Disease; SD, standard deviation.


Figure 1: The incidence of post-transplantation outcomes among the recipients of liver transplantation by type of donors



N is the total number of recipients in each group; n is the number of recipients with a particular outcome. In DDLT group, N=568 (for rejection) and N=254 (for HCC recurrence). In LDLT group, N=1942 for rejection and N=1245 for HCC recurrence.

DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; LDLT, living donor liver transplantation; NODM, new onset of diabetes mellitus.


Figure 2. Causes of graft loss in adult patients with liver transplantation (N = 195)



N is the total number of recipients with graft loss; n is the number of recipients with a particular reason. HCC, hepatocellular carcinoma; IVC, inferior vena cava.

Accepted

Figure 3: Kaplan-Meier survival curve of post-transplantation outcomes in adult patients with liver transplantation



DDLT, deceased donor liver transplantation; LDLT, living donor liver transplantation; HCC, hepatocellular carcinoma; NODM, new onset of diabetes mellitus.

Supplementary material

Supplementary Table 1. Multiple Cox regression analysis for risk factors influencing occurrence of death after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Death (N=2367)				
Recipients' age	1.43 (1.20, 1.70)	<.001	1.38 (1.15, 1.67)	<.001
Donors' age	1.30 (1.18, 1.43)	<.001	—	—
Male recipient	0.88 (0.64, 1.22)	0.445	1.39 (0.93, 2.08)	0.108
Male donor	0.88 (0.65, 1.20)	0.425	—	—
Recipients' BMI ≥ 25 (kg/m ²)	0.87 (0.63, 1.19)	0.376	0.81 (0.57, 1.15)	0.242
Donors' BMI ≥ 25 (kg/m ²)	0.97 (0.69, 1.36)	0.848	—	—
LDLT vs DDLT	0.29 (0.22, 0.40)	<.001	0.22 (0.09, 0.53)	<.001
Hypertension	1.45 (1.01, 2.07)	0.043	1.29 (0.85, 1.95)	0.229
Diabetes mellitus	1.37 (0.99, 1.89)	0.060	1.01 (0.69, 1.47)	0.958
MELD score: ≥ 35	3.39 (2.37, 4.86)	<.001	1.42 (0.89, 2.25)	0.140
HCC	0.81 (0.60, 1.10)	0.179	—	—
Acute hepatitis	2.22 (1.28, 3.83)	0.004	1.53 (0.78, 3.00)	0.212
ABO incompatible	0.67 (0.41, 1.09)	0.105	1.44 (0.83, 2.48)	0.193
Use of steroids	0.28 (0.20, 0.38)	<.001	0.47 (0.31, 0.71)	<.001
Use of anti-metabolites	0.33 (0.24, 0.44)	<.001	0.54 (0.38, 0.78)	<.001
Use of mTOR inhibitors	0.81 (0.49, 1.31)	0.386	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 2. Multiple Cox regression analysis for risk factors influencing occurrence of graft loss after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Graft loss (N=2368)				
Recipients' age	1.03 (1.01, 1.04)	0.002	1.24 (1.05, 1.46)	0.012
Donors' age	1.02 (1.02, 1.03)	<.001	1.04 (0.92, 1.17)	0.536
Male recipient	0.86 (0.64, 1.17)	0.344	—	—
Male donor	0.89 (0.67, 1.19)	0.425	—	—
Recipients' BMI ≥ 25 (kg/m ²)	0.92 (0.68, 1.25)	0.611	—	—
Donors' BMI ≥ 25 (kg/m ²)	1.01 (0.72, 1.39)	0.984	—	—
LDLT vs DDLT	0.33 (0.25, 0.43)	<.001	0.22 (0.09, 0.54)	0.001
Hypertension	1.35 (0.96, 1.90)	0.081	—	—
Diabetes mellitus	1.23 (0.90, 1.67)	0.190	—	—
MELD score: ≥ 35	2.89 (2.00, 4.19)	<.001	—	—
HCC	0.81 (0.61, 1.08)	0.145	1.66 (1.13, 2.46)	0.011
Acute hepatitis	2.45 (1.42, 4.20)	0.001	1.90 (1.02, 3.55)	0.044
ABO incompatible	1.42 (0.91, 2.21)	0.123	1.40 (0.86, 2.29)	0.177
Use of steroids	0.32 (0.23, 0.44)	<.001	0.57 (0.39, 0.84)	0.005
Use of anti-metabolites	0.36 (0.27, 0.48)	<.001	0.56 (0.40, 0.78)	<.001
Use of mTOR inhibitors	0.77 (0.48, 1.22)	0.264	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 3. Multiple Cox regression analysis for risk factors influencing occurrence of HCC recurrence after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
HCC recurrence (N=1492)				
Recipients' age	0.99 (0.97, 1.02)	0.430	0.94 (0.74, 1.19)	0.610
Donors' age	1.00 (0.99, 1.01)	0.911	—	—
Male recipient	2.53 (1.32, 4.86)	0.005	—	—
Male donor	0.74 (0.50, 1.09)	0.130	—	—
Recipients' BMI ≥ 25 (kg/m ²)	0.53 (0.47, 1.23)	0.472	—	—
Donors' BMI ≥ 25 (kg/m ²)	0.98 (0.82, 1.56)	0.614	—	—
LDLT vs DDLT	1.18 (0.70, 2.00)	0.527	—	—
Hypertension	1.38 (0.87, 2.19)	0.167	—	—
Diabetes mellitus	0.92 (0.58, 1.46)	0.732	—	—
MELD score: ≥ 35	1.62 (0.53, 2.82)	0.326	1.58 (0.64, 3.92)	0.321
HCC	19.81 (2.76, 142.02)	0.003	—	—
Acute hepatitis	1.52 (0.72, 2.96)	0.612	—	—
Pretransplant HCC Beyond Milan criteria vs Within Milan criteria	3.19 (1.84-4.52)	<.001	3.06 (2.00, 4.69)	<.001
ABO incompatible	0.60 (0.38, 0.95)	0.029	—	—
Use of steroids	0.86 (0.45, 1.65)	0.642	—	—
Use of anti-metabolites	0.89 (0.57, 1.38)	0.595	—	—
Use of mTOR inhibitors	4.74 (3.20, 7.02)	<.001	3.42 (2.25, 5.20)	<.001

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 4. Multiple Cox regression analysis for risk factors influencing occurrence of rejection after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Rejection (N=2326)				
Recipients' age	1.00 (0.98, 1.01)	0.783	0.90 (0.78, 1.05)	0.172
Donors' age	1.02 (1.01, 1.03)	0.001	1.068 (0.96, 1.20)	0.249
Male recipient	0.89 (0.66, 1.19)	0.425	—	—
Male donor	1.04 (0.78, 1.38)	0.815	—	—
Recipients' BMI ≥ 25 (kg/m ²)	1.03 (0.77, 1.39)	0.830	—	—
Donors' BMI ≥ 25 (kg/m ²)	1.02 (0.74, 1.40)	0.911	—	—
LDLT vs DDLT	0.62 (0.46, 0.83)	0.001	0.66 (0.45, 0.99)	0.042
Hypertension	1.14 (0.80, 1.62)	0.463	—	—
Diabetes mellitus	1.03 (0.75, 1.41)	0.865	—	—
MELD score: ≥ 35	0.67 (0.38, 1.18)	0.164	—	—
HCC	0.88 (0.67, 1.15)	0.349	—	—
Acute hepatitis	0.69 (0.30, 1.58)	0.375	—	—
ABO incompatible	0.70 (0.50, 0.98)	0.036	1.79 (1.24, 2.59)	0.002
Use of steroids	1.89 (1.06, 3.39)	0.032	—	—
Use of anti-metabolites	1.35 (0.96, 1.90)	0.086	—	—
Use of mTOR inhibitors	1.51 (1.05, 2.17)	0.025	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 5. Multiple Cox regression analysis for risk factors influencing occurrence of renal failure after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Renal failure (N=2561)				
Recipients' age	1.03 (1.01, 1.05)	0.010	1.40 (1.15, 1.70)	<.001
Donors' age	1.03 (1.02, 1.04)	<.001	—	—
Male recipient	0.77 (0.54, 1.10)	0.151	0.84 (0.59, 1.19)	0.315
Male donor	0.80 (0.57, 1.13)	0.205	—	—
Recipients' BMI ≥ 25 (kg/m ²)	1.23 (0.86, 1.75)	0.251	1.18 (0.83, 1.69)	0.349
Donors' BMI ≥ 25 (kg/m ²)	1.21 (0.84, 1.77)	0.311	—	—
LDLT vs DDLT	0.20 (0.14, 0.28)	<.001	0.44 (0.29, 0.66)	<.001
Hypertension	1.53 (1.03, 2.27)	0.033	1.23 (0.81, 1.88)	0.338
Diabetes mellitus	1.47 (1.03, 2.09)	0.036	1.42 (0.97, 2.08)	0.072
MELD score: ≥ 35	8.49 (5.86, 12.31)	<.001	5.07 (2.12, 12.13)	<.001
HCC	0.51 (0.36, 0.73)	<.001	—	—
Acute hepatitis	0.88 (0.35, 2.19)	0.782	—	—
ABO incompatible	1.87 (1.03, 3.38)	0.039	—	—
Use of steroids	0.36 (0.25, 0.54)	<.001	0.46 (0.31, 0.69)	<.001
Use of anti-metabolites	0.39 (0.28, 0.55)	<.001	0.53 (0.37, 0.76)	<.001
Use of mTOR inhibitors	1.32 (0.83, 2.08)	0.240	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 6. Multiple Cox regression analysis for risk factors influencing occurrence of renal dysfunction after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Renal dysfunction (N=2556)				
Recipients' age	1.03 (1.02, 1.03)	<.001	1.28 (1.19, 1.38)	<.001
Donors' age	1.01 (1.00, 1.01)	0.001	—	—
Male recipient	0.71 (0.62, 0.81)	<.001	1.03 (0.90, 1.18)	0.679
Male donor	1.04 (0.91, 1.19)	0.541	—	—
Recipients' BMI ≥ 25 (kg/m ²)	1.05 (0.89, 1.25)	0.512	—	—
Donors' BMI ≥ 25 (kg/m ²)	1.12 (0.93, 1.34)	0.225	—	—
LDLT vs DDLT	0.62 (0.54, 0.71)	<.001	0.79 (0.67, 0.93)	0.004
Hypertension	1.34 (1.14, 1.56)	<.001	1.25 (1.06, 1.47)	0.007
Diabetes mellitus	1.32 (1.15, 1.52)	<.001	1.20 (1.04, 1.39)	0.011
MELD score: ≥ 35	2.26 (1.73, 2.95)	<.001	1.81 (1.47, 2.23)	<.001
HCC	0.70 (0.61, 0.79)	<.001	—	—
Acute hepatitis	1.27 (0.86, 1.89)	0.227	1.06 (0.78, 1.44)	0.695
ABO incompatible	1.46 (1.20, 1.78)	<.001	—	—
Use of steroids	0.77 (0.64, 0.93)	0.007	0.82 (0.67, 0.99)	0.040
Use of anti-metabolites	0.78 (0.68, 0.89)	<.001	—	—
Use of mTOR inhibitors	0.92 (0.76, 1.12)	0.412	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 7. Multiple Cox regression analysis for risk factors influencing occurrence of infections after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Infections (N=2563)				
Recipients' age	1.01 (1.00, 1.02)	0.071	1.11 (1.03, 1.19)	0.009
Donors' age	1.02 (1.01, 1.02)	<.001	—	—
Male recipient	0.79 (0.68, 0.92)	0.002	—	—
Male donor	0.98 (0.85, 1.13)	0.738	—	—
Recipients' BMI ≥ 25 (kg/m ²)	0.94 (0.79, 1.12)	0.499	—	—
Donors' BMI ≥ 25 (kg/m ²)	0.88 (0.73, 1.07)	0.197	—	—
LDLT vs DDLT	0.49 (0.43, 0.57)	<.001	—	—
Hypertension	0.94 (0.78, 1.13)	0.499	1.08 (0.85, 1.37)	0.527
Diabetes mellitus	1.09 (0.94, 1.28)	0.257	1.09 (0.93, 1.28)	0.268
MELD score: ≥ 35	2.49 (1.91, 3.25)	<.001	1.86 (1.50, 2.30)	<.001
HCC	0.74 (0.64, 0.85)	<.001	—	—
Acute hepatitis	1.48 (0.99, 2.21)	0.051	0.83 (0.59, 1.17)	0.294
ABO incompatible	1.15 (0.94, 1.40)	0.168	1.10 (0.90, 1.35)	0.370
Use of steroids	0.38 (0.32, 0.46)	<.001	—	—
Use of anti-metabolites	0.62 (0.54, 0.72)	<.001	—	—
Use of mTOR inhibitors	1.40 (1.15, 1.69)	<.001	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 8. Multiple Cox regression analysis for risk factors influencing occurrence of surgical complications after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
Surgical complications (N=2561)				
Recipients' age	1.00 (0.99, 1.01)	0.689	1.05 (0.96, 1.15)	0.280
Donors' age	1.01 (1.00, 1.01)	0.059	—	—
Male recipient	0.95 (0.79, 1.13)	0.537	—	—
Male donor	1.03 (0.87, 1.21)	0.771	—	—
Recipients' BMI ≥ 25 (kg/m ²)	1.32 (1.09, 2.11)	0.031	1.06 (0.89, 1.25)	0.533
Donors' BMI ≥ 25 (kg/m ²)	0.73 (0.51, 1.42)	0.846	—	—
LDLT vs DDLT	0.94 (0.78, 1.14)	0.539		
Hypertension	0.91 (0.74, 1.14)	0.417	—	—
Diabetes mellitus	0.80 (0.66, 0.98)	0.028	0.79 (0.65, 0.97)	0.021
MELD score: ≥ 35	1.72 (1.25, 1.87)	0.005	1.46 (1.11, 1.92)	0.007
HCC	0.93 (0.79, 1.09)	0.352	0.95 (0.80, 1.13)	0.550
Acute hepatitis	1.45 (0.94, 1.87)	0.650	—	—
ABO incompatible	0.91 (0.73, 1.13)	0.388	—	—
Use of steroids	0.86 (0.67, 1.11)	0.246	—	—
Use of anti-metabolites	0.91 (0.76, 1.09)	0.284	—	—
Use of mTOR inhibitors	0.97 (0.76, 1.24)	0.802	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.

Supplementary Table 9. Multiple Cox regression analysis for risk factors influencing occurrence of NODM after liver transplantation

Variables	Multiple Cox regression			
	Univariate		Multivariate	
	HR (95% CI)	p-value	HR (95% CI)	p-value
NODM (N=2554)				
Recipients' age	1.01 (1.00, 1.03)	0.032	1.10 (0.98, 1.23)	0.100
Donors' age	0.99 (0.99, 1.00)	0.034	—	—
Male recipient	1.08 (0.86, 1.35)	0.536	1.21 (0.98, 1.51)	0.082
Male donor	1.27 (1.02, 1.58)	0.031	—	—
Recipients' BMI ≥ 25 (kg/m ²)	1.37 (1.09, 1.70)	0.006	1.39 (1.13, 1.71)	0.002
Donors' BMI ≥ 25 (kg/m ²)	0.76 (0.59, 0.99)	0.039	—	—
LDLT vs DDLT	1.30 (1.01, 1.67)	0.042		
Hypertension	1.02 (0.79, 1.33)	0.877	0.91 (0.69, 1.19)	0.490
Diabetes mellitus	—	—	—	—
MELD score: ≥ 35	0.78 (0.53, 1.16)	0.225	—	—
HCC	0.88 (0.72, 1.08)	0.217	—	—
Acute hepatitis	0.53 (0.26, 1.05)	0.068	—	—
ABO incompatible	0.84 (0.64, 1.09)	0.185	—	—
Use of steroids	3.03 (1.81, 5.08)	<.001	3.24 (1.90, 5.54)	<.001
Use of anti-metabolites	0.74 (0.59, 0.91)	0.005	0.68 (0.54, 0.84)	<.001
Use of mTOR inhibitors	0.75 (0.53, 1.06)	0.102	—	—

BMI, body-mass index; CI, confidence interval; DDLT, deceased donor liver transplantation; HCC, hepatocellular carcinoma; HR, hazards ratio; LDLT, living donor liver transplantation; MELD, Model for End-Stage Liver Disease; mTOR, mammalian target of rapamycin.