




Genome-Wide Transcriptional Response to Varying RpoS Levels in *Escherichia coli* K-12

Garrett T. Wong,^{a,*} Richard P. Bonocora,^b Alicia N. Schep,^{a,*} Suzannah M. Beeler,^{a,*} Anna J. Lee Fong,^{a,*} Lauren M. Shull,^{a,*} Lakshmi E. Batachari,^a Moira Dillon,^a Ciaran Evans,^{c,*} Carla J. Becker,^a Eliot C. Bush,^a Johanna Hardin,^c Joseph T. Wade,^{b,d}  Daniel M. Stoebel^a

Department of Biology, Harvey Mudd College, Claremont, California, USA^a; Wadsworth Center, New York State Department of Health, Albany, New York, USA^b; Department of Mathematics, Pomona College, Claremont, California, USA^c; Department of Biomedical Sciences, School of Public Health, University at Albany, SUNY, Albany, New York, USA^d

ABSTRACT The alternative sigma factor RpoS is a central regulator of many stress responses in *Escherichia coli*. The level of functional RpoS differs depending on the stress. The effect of these differing concentrations of RpoS on global transcriptional responses remains unclear. We investigated the effect of RpoS concentration on the transcriptome during stationary phase in rich media. We found that 23% of genes in the *E. coli* genome are regulated by RpoS, and we identified many RpoS-transcribed genes and promoters. We observed three distinct classes of response to RpoS by genes in the regulon: genes whose expression changes linearly with increasing RpoS level, genes whose expression changes dramatically with the production of only a little RpoS ("sensitive" genes), and genes whose expression changes very little with the production of a little RpoS ("insensitive"). We show that sequences outside the core promoter region determine whether an RpoS-regulated gene is sensitive or insensitive. Moreover, we show that sensitive and insensitive genes are enriched for specific functional classes and that the sensitivity of a gene to RpoS corresponds to the timing of induction as cells enter stationary phase. Thus, promoter sensitivity to RpoS is a mechanism to coordinate specific cellular processes with growth phase and may also contribute to the diversity of stress responses directed by RpoS.

IMPORTANCE The sigma factor RpoS is a global regulator that controls the response to many stresses in *Escherichia coli*. Different stresses result in different levels of RpoS production, but the consequences of this variation are unknown. We describe how changing the level of RpoS does not influence all RpoS-regulated genes equally. The cause of this variation is likely the action of transcription factors that bind the promoters of the genes. We show that the sensitivity of a gene to RpoS levels explains the timing of expression as cells enter stationary phase and that genes with different RpoS sensitivities are enriched for specific functional groups. Thus, promoter sensitivity to RpoS is a mechanism that coordinates specific cellular processes in response to stresses.

KEYWORDS RpoS, promoters, σ^S , stress response, transcriptional regulation, transcriptome

Genome-wide measurements of RNA levels have revolutionized our understanding of how cells organize their patterns of transcription. These studies have given us snapshots of how patterns of gene expression change in response to changes in the external environment. They have also allowed us to define the regulons controlled by specific transcription factors (TFs). A major weakness of the vast majority of these

Received 21 October 2016 Accepted 12 January 2017

Accepted manuscript posted online 23 January 2017

Citation Wong GT, Bonocora RP, Schep AN, Beeler SM, Lee Fong AJ, Shull LM, Batachari LE, Dillon M, Evans C, Becker CJ, Bush EC, Hardin J, Wade JT, Stoebel DM. 2017. Genome-wide transcriptional response to varying RpoS levels in *Escherichia coli* K-12. J Bacteriol 199:e00755-16. <https://doi.org/10.1128/JB.00755-16>.

Editor Thomas J. Silhavy, Princeton University

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Joseph T. Wade, joseph.wade@health.ny.gov, or Daniel M. Stoebel, stoebel@g.hmc.edu.

* Present address: Garrett T. Wong, Departments of Neuroscience, Genetics, and Genome Sciences, Icahn School of Medicine at Mount Sinai, New York, New York, USA; Alicia N. Schep, Department of Genetics, Stanford University, Stanford, California, USA; Suzannah M. Beeler, Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, USA; Anna J. Lee Fong, a-tek, Washington, DC, USA; Lauren M. Shull, Sackler School of Graduate Biomedical Sciences, Tufts University, Boston, Massachusetts, USA; Ciaran Evans, Department of Statistics, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA.

studies is that they explore the function of a regulatory protein only by comparing expression of target genes in a wild-type strain to levels in a gene knockout strain, or to a mutant with a single diminished or increased level of activity. While some genetic regulatory networks are certainly switch-like (1, 2) and can be fully characterized by only two levels of activity, many other regulatory proteins vary continuously in their abundance and/or activity. How regulons respond to a range of regulator levels is a largely unstudied question (3). Should we expect all genes in a regulon to increase or decrease expression by the same relative amount following a change in abundance of a regulatory protein? Or should we expect genes to respond in different ways? These questions motivated this study.

A paradigmatic example of a bacterial regulatory protein whose abundance and activity vary continuously in response to different conditions is the alternative sigma factor RpoS in *Escherichia coli*. Transcription by RNA polymerase containing RpoS is responsible for the general stress response (4–6). Under conditions of optimal growth in the laboratory (such as exponential-phase growth in rich medium at 37°C), RpoS levels are nearly undetectable in the model strain *E. coli* K-12. As conditions become poorer for growth, either because cells begin to starve for various nutrients or because they face physical challenges, such as low temperature or elevated osmolarity, RpoS levels rise (4–6). RpoS coordinates the transcription of genes that are critical for the response to these stresses.

RpoS expression is not an all-or-none phenomenon. For example, RpoS levels rise continuously as cells transition from exponential growth to stationary phase (7). Moreover, starvation for different nutrients upregulates RpoS to different levels (8, 9). This level of control over RpoS levels is accomplished by regulating transcription, translation, and protein degradation (4, 7, 10), allowing for careful control over protein levels. In addition to regulation of protein abundance, RpoS activity can also be directly modulated by a number of factors, such as Crl (11).

Not only do RpoS levels vary across conditions for a single strain, but different strains of *E. coli* also differ in their patterns of expression of RpoS. For example, naturally occurring strains can differ in the amount of RpoS they produce during exponential phase (12) or stationary phase (13). All studies that have measured RpoS levels in naturally occurring strains of *E. coli* have detected variations (13–17), though the extent and cause of the variations in RpoS between strains is still a matter of some controversy (16, 17).

Microarray studies (18–20) have shown that RpoS controls the expression of at least 500 genes (over 10% of the genome) either directly or indirectly, but the set of RpoS-regulated genes differs across environmental conditions. For most RpoS-controlled genes, it is not known whether the gene is directly transcribed by RpoS or regulated indirectly as a consequence of RpoS transcribing other genes. Previous studies have not investigated the impact of changes in RpoS levels on the RpoS regulon, or whether quantitative differences in RpoS levels between environmental conditions influence the observed differences in what genes are RpoS regulated. It is clear that *E. coli* has a complicated regulatory network that fine-tunes RpoS levels under different conditions, but we do not yet know the consequences of this regulation.

In this study, we tested the hypothesis that members of the RpoS regulon vary in their responses to RpoS levels. Using a combination of chromatin immunoprecipitation sequencing (ChIP-seq) and transcriptome sequencing (RNA-seq), we identified RpoS-regulated genes, and we showed that genes vary in their sensitivity to RpoS levels in a manner dependent on sequences outside the core promoter region. Sensitivity of genes to RpoS levels corresponds to the order in which genes are induced during the transition into stationary phase, and genes with different levels of sensitivity are enriched for specific functional groups. Thus, the levels of sensitivity of genes to RpoS control the physiological response to different stress conditions.

TABLE 1 Biological processes enriched in the RpoS regulon

		No. of genes			
			In regulon		
GO ID no.	GO term	In genome	Observed	Expected	<i>P</i> value
GO:0009063	Cellular amino acid catabolic process	69	38	18.5	3.5e−07
GO:0006099	Tricarboxylic acid cycle	20	15	5.35	8.3e−06
GO:0006096	Glycolytic process	18	14	4.81	8.3e−06
GO:0006094	Gluconeogenesis	13	9	3.48	0.0016
GO:0009441	Glycolate metabolic process	6	5	1.6	0.0063
GO:0006090	Pyruvate metabolic process	21	17	5.61	0.0180
GO:0006970	Response to osmotic stress	28	13	7.49	0.0190
GO:0006542	Glutamine biosynthetic process	10	6	2.67	0.0270
GO:0009060	Aerobic respiration	65	33	17.38	0.0285
GO:0006006	Glucose metabolic process	36	20	9.62	0.0317
GO:0016052	Carbohydrate catabolic process	276	94	73.79	0.0326
GO:0009101	Glycoprotein biosynthetic process	13	7	3.48	0.0342
GO:0019395	Fatty acid oxidation	30	13	8.02	0.0354
GO:0000162	Tryptophan biosynthetic process	11	6	2.94	0.0463
GO:0009246	Enterobacterial common antigen biosynthesis	11	6	2.94	0.0463
GO:0042398	Cellular modified amino acid biosynthesis	17	7	4.55	0.0472
GO:0009239	Enterobactin biosynthetic process	6	4	1.6	0.0472
GO:0000270	Peptidoglycan metabolic process	51	14	13.64	0.0475

RESULTS

The RpoS regulon in late stationary phase. To understand the role of the RpoS protein in late stationary phase, we used RNA-seq to compare the transcriptome of wild-type and $\Delta rpoS$ mutant cells. We observed differential expression of 1,044 genes (23% of genes) between these two conditions ($P < 0.05$). Of the 1,044 genes whose expression is influenced by RpoS, 605 are upregulated and 439 are downregulated (see Table S4 in the supplemental material).

Influencing transcription of 23% of the genome could have many potential phenotypic effects. To better understand the function of these genes, we examined which kinds of gene functions, as described by the Gene Ontology Consortium's GO database, are more abundant in the regulon than expected by chance. GO enrichment analysis indicated that the RpoS regulon includes many genes involved in metabolic processes (Table 1); 17 of 18 significantly enriched GO terms were metabolic terms. This metabolic reorganization includes the upregulation of genes encoding glycolytic enzymes and pathways for metabolism of L-arginine to glutamine, or from L-arginine to putrescine and then into succinate. RpoS also drives the downregulation of genes involved in the tricarboxylic acid (TCA) cycle. These patterns of metabolic regulation are very similar to those identified in late stationary phase in *Salmonella enterica* (21). The only significant GO term not explicitly linked to metabolism (GO:0006970, response to osmotic stress) also includes metabolic genes, such as *otsA* and *otsB*, that are involved in trehalose biosynthesis.

Central metabolism is not the only phenotype similarly regulated by RpoS in both *S. enterica* and *E. coli*. Other similarities include transcription of genes involved in antioxidant activities, iron regulation and Fe-S cluster assembly, upregulation of proteases, and downregulation of porins. As with *S. enterica*, RpoS in *E. coli* influences the expression of many genes encoding other regulatory proteins, including *csrA*, *arcA*, *cra*, *fur*, *ihfA*, *hupA*, and *hupB*. These proteins regulate phenotypes, including carbon storage (CsrA), central carbon metabolism (ArcA and Cra), and iron homeostasis (Fur) and also play a central role in structuring the nucleoid (IHf and HU). Not all of the regulation is identical between *E. coli* and *S. enterica*, however. For example, Lévi-Meyrueis et al. (21) noted that RpoS appears to direct switching between many pairs of isozymes, where one isozyme is expressed under conditions when RpoS is abundant and its partner isozyme is expressed when RpoS levels are low. While some pairs show a similar pattern in *E. coli* (such as *tktA* and *tktB* and *acnA* and *acnB*), others (such as *fumA* and *fumC*) do




FIG 1 Analysis of RpoS ChIP-seq data. (A) Sequence logo (86) of the enriched sequence motif associated with RpoS-bound regions (MEME E value, 2×10^{-40}). The sequence logo was generated using WebLogo (87). Numbers below the motif refer to positions relative to the transcription start, based on known sequence preferences for RpoS (22). (B) Histogram showing the frequency distribution of distances between identified motifs and ChIP-seq peak centers. The dashed line indicates the peak center position. (C) Sequence logo (86) of the enriched sequence motif associated with TSSs located within 20 bp of RpoS ChIP-seq peak centers (MEME E value, 3×10^{-56}). Note that only a subset of the motif identified by MEME is shown, but positions outside the shown region all have information content values below 0.2. The sequence logo was generated using WebLogo (87). Numbers below the motif refer to positions relative to the transcription start, based on known sequence preferences for RpoS (22).

not show this pattern. The reason why these enzymes involved in central carbon metabolism might show this pattern is not clear.

Genome-wide binding profile of RpoS. While RNA-seq identified which genes are regulated by RpoS, it cannot distinguish between direct and indirect effects. To determine sites where RpoS binds (and hence likely plays a direct role in transcription), we used ChIP-seq to map the association of RpoS across the *E. coli* chromosome during stationary-phase growth in minimal medium. To facilitate ChIP, RpoS was C-terminally sequential peptide affinity (SPA) tagged at its native locus. We reasoned that RpoS would only be identified with promoter regions, since it is likely released from elongating RNA polymerase complexes. We identified 284 peaks of RpoS ChIP-seq signal covering 260 genomic regions (peaks within 100 bp of each other were merged). A total of 217 of the RpoS-bound regions are intergenic, and 67 are located within genes. We reasoned that annotated genes that are transcribed by RpoS would be positioned close to an RpoS-bound region. Consistent with this, 213 RpoS-bound regions are ≤ 300 bp upstream of an annotated gene start. These 213 regions include 27 that are intragenic. In 79 cases, we observed an RpoS-bound region ≤ 300 bp from the starts of two divergently transcribed genes.

We used MEME tools to search for enriched sequence motifs within the RpoS-bound regions. We detected a highly enriched motif in 107 regions (Fig. 1A); this motif closely resembled the known -10 hexamer recognized by RpoS (CTAYACT, where the central YA are not as conserved as the other positions [22]). Moreover, occurrences of the motif are positionally enriched with respect to the ChIP-seq peak center (Fig. 1B), indicating that the data have a high spatial resolution. Note that the motifs tend to be located just upstream of the peak centers (Fig. 1B), as we observed previously for the *E. coli* flagellar sigma factor FliA (23). This presumably reflects the fact that the footprint of initiating RNA polymerase associated with RpoS is not centered on the -10 hexamer.

The identification of RpoS binding sites allowed us to better understand the role of RpoS in both positive and negative regulation. While similar proportions of the RpoS regulon are positively and negatively regulated by RpoS, it is not clear if this is true at the level of direct regulation. Only 19 genes within 300 bp of a binding site are negatively regulated by RpoS. The finding of 19 genes (of the 286 such genes) is fewer than we would expect by chance if binding sites were randomly distributed around the genome, given that 439 of 4,513 genes in the genome are negatively regulated. On the other hand, 111 of 286 genes within 300 bp are positively regulated, a highly significant

effect (Fisher's exact test, $P < 10^{-16}$). Thus, the binding profile of RpoS is consistent with direct positive regulation of many genes but provides no evidence of direct negative regulation.

Identification of RpoS-transcribed genes and promoters. We combined the ChIP-seq and RNA-seq data to identify genes that are directly transcribed by RNA polymerase containing RpoS. From these data, we identified 123 RpoS-transcribed genes in 99 transcripts (see Table S5 in the supplemental material) and compared these findings to those of other published analyses (18–20, 24–28). In some cases, we identified RpoS-bound regions upstream of genes that were not detected as being RpoS regulated by RNA-seq. These genes may have promoters that bind transcriptionally inactive RNA polymerase (29, 30). (Inactive polymerase could be bound but physically blocked from transcription elongation by the presence of a bound repressor [31] or could be poised, waiting for a signal to transition to transcription elongation [32, 33].) Alternatively, the disparity between RpoS binding and regulation could be explained by differences in growth conditions between the ChIP-seq and RNA-seq experiments or by the possibility that the C-terminal SPA tag affects the function of the C terminus of the protein in the response to transcription activators (29, 33).

The high spatial resolution of sigma factor ChIP-seq can facilitate the identification of specific promoters (23, 34) when combined with nucleotide resolution transcription start site (TSS) maps. Using published TSS data for *E. coli* under stationary-phase conditions similar to those used for the ChIP-seq experiment (25), we determined all pairwise distances between RpoS ChIP-seq peaks and stationary-phase TSSs. We observed a strong enrichment for peak TSS distances of ≤ 20 bp (Fig. S2). We inferred that these 112 TSSs are RpoS transcribed. Consistent with this, the putative RpoS-transcribed TSSs are associated with -10 hexamers that have features expected of RpoS promoters (Fig. 1C). In some cases, RpoS promoters were identified ≤ 300 bp upstream of genes that were not RpoS regulated in the RNA-seq data set. We presumed that these are RpoS-transcribed genes that otherwise escaped detection because of differences in the growth conditions used for ChIP-seq and RNA-seq analyses.

The transcriptome at three different RpoS levels. This first view of the RpoS regulon considers RpoS either present or absent. RpoS levels vary continuously across environmental stresses (7), so we sought to better understand how the level of RpoS in the cell influences transcription of target genes. To do this, we placed the *rpoS* gene under the control of the arabinose-inducible promoter P_{araB} . This promoter was integrated just upstream of the native *rpoS* gene, placing transcription under the control of arabinose concentration and removing the 5' region that regulates translation of the native mRNA (4).

To measure the resulting arabinose-induced expression of RpoS, we employed quantitative Western blotting. RpoS levels increased with increasing arabinose concentration, from undetectable RpoS levels to levels similar to those in wild-type cells (Fig. 2A). To confirm that expression was graded and not an all-or-none response in this system (35), we used flow cytometry to measure expression in individual cells. We transformed the arabinose-inducible RpoS strain with the plasmid pDMS123 (36), which contains the RpoS-dependent *otsBA* promoter fused to *gfp*, the gene for green fluorescent protein (GFP). As expected, *gfp* expression increased with increasing arabinose concentrations, and at each expression level the population was unimodal (Fig. 2B).

We measured the transcriptome in cells with RpoS levels that were 26% of wild-type levels; this was achieved by the addition of $10^{-4}\%$ arabinose to cells with our arabinose-inducible *rpoS* strain. Of the genes that were differentially expressed between 26% RpoS and either 0% ($\Delta rpoS$ mutant) or 100% (wild type), 95% were also differentially expressed between 0% and 100% ($P < 0.05$) (Fig. 3).

Nearly all genes that were significantly differentially expressed had monotonically increasing or decreasing patterns of expression between the three levels of RpoS. Only two genes (*ytfR* and *ytfT*) had an expression level at 26% RpoS that was significantly higher than expression at both 100% and 0% RpoS. The only two genes with expression




FIG 2 Arabinose control of RpoS protein levels in strain DMS2564. (A) Increasing the arabinose concentration results in greater RpoS expression in strain DMS2564, as measured by Western blotting. $n = 4$ to 5. Error bars represent standard errors of the mean results. wt, wild type. (B) Flow cytometric measurement of an *otsB-gfp* fusion under 0%, 10⁻⁴%, and 10⁻¹% arabinose (dotted line, dashed line, and solid line, respectively).

lower under the 26% RpoS condition than under either 100% or 0% were *nlpD* and *pcm*. These genes lie immediately upstream of *rpoS*. *nlpD* was removed from the genome during the construction of the arabinose-inducible RpoS strain. The *pcm* gene was still present, but the level of transcription was lowered by the genetic modification.

Classifying quantitative responses to RpoS levels. To explore how genes respond to changing levels of RpoS, we developed a new metric, sensitivity. Our null expectation was that gene expression would increase linearly with increasing RpoS concentration. We observed many genes in our RNA-seq data set (such as *osmY*) whose expression at intermediate RpoS levels fell on, or close to, a line drawn between the 0% and 100% RpoS conditions (Fig. 4A). We refer to these genes as linear in their response to increasing RpoS levels. Other genes (such as *astA*) were transcribed more at 26% than would be expected based on their expression levels at 0% and 100% (Fig. 4B). We refer to these genes as sensitive, because only a small amount of RpoS resulted in relatively high levels of transcription. In contrast, the expression of some genes (like *gadC*) at intermediate RpoS levels was less than expected based on expression at 0% and 100% RpoS; such genes are referred to as insensitive (Fig. 4C).

We identified 910 linear, 102 sensitive, and 32 insensitive genes. Ninety-six percent of sensitive genes and 88% of insensitive genes are positively regulated by RpoS. In contrast, only 53% of linear genes are positively regulated by RpoS, a significant difference (chi-square test, $P < 0.001$).

To determine whether sensitive or insensitive genes are associated with specific physiological responses to increasing RpoS levels, we again used GO enrichment. We tested the null hypothesis that the functions of these genes are a random sample from the entire RpoS regulon (not the whole genome). The GO terms significantly enriched in the sensitive class are response to osmotic stress, cellular amino acid catabolic




FIG 3 Area-proportional Venn diagram showing the number of genes differentially expressed under each of the three conditions.




FIG 4 Examples of three classes of patterns of RpoS response. *osmY* expression is linear with the level of RpoS, *astA* is sensitive to the level of RpoS, and *gadC* is insensitive to the level of RpoS. Dots represent normalized expression levels from individual RNA-seq samples. The lines are drawn between the mean at 0% and the mean at 100%. wt, wild type.

process, and fatty acid oxidation (Table 2). Several genes encoding regulators were among the sensitive genes, including *arcA*, which encodes a global regulator of respiratory metabolism, and *rssB*, which encodes the adaptor protein required for degradation of RpoS by ClpXP.

GO enrichment was less useful for understanding possible functions of the insensitive gene set. Three GO terms were enriched (Table 3), but only a few genes with these annotations were present in the insensitive gene set, and their enrichment probably reflected the relatively small number of insensitive genes. More strikingly, the insensitive genes included nearly all of the genes required for acid resistance system 2: the structural genes *gadA*, *gadB*, and, *gadC* and the regulator of this system, *gadE* (37, 38). In addition, the genes *yhiM*, *yhiD*, *hdeA*, *hdeB*, *hdeD*, *mdtE*, and *mdtF*, all of which have been described as having roles in acid resistance (37, 38), were insensitive.

We used reverse transcription coupled to quantitative PCR (qPCR) to confirm the expression patterns of two insensitive genes (*gadC* and *gadE*) and three sensitive genes (*prpR*, *prpD*, and *astA*). All genes were positively regulated by RpoS (Fig. 5), and the median expression at 26% RpoS was consistent with RNA-seq expression patterns for all genes. We used a bootstrapping approach to assess if expression at 26% was significantly above or below the linear expectation at 26% RpoS. The expression of *gadC* was significantly insensitive ($P < 10^{-4}$), and *prpR* expression was significantly sensitive ($P < 10^{-4}$). *astA* expression was marginally significantly sensitive ($P = 0.06$ for sensitivity), while *gadE* and *prpD* were not significantly different from the linear expectation ($P = 0.10$ and $P = 0.56$, respectively).

Control of sensitivity of expression. What makes one promoter sensitive to RpoS levels and another promoter insensitive to RpoS levels? We hypothesized three possible mechanisms. First, chromosomal location could determine the response to RpoS levels, as is known to occur in the context of total transcription levels (39). Second, it is possible that the DNA sequence of the core promoter drives the response. Finally, it is possible that the binding of transcription factors upstream of the core promoter influences the response to RpoS levels. To test these hypotheses, we cloned the promoters (including all upstream transcription factor binding sites annotated in EcoCyc [40]) of four operons into the *lacZ* fusion plasmid pLFX (41). The four promoters were the sensitive *astCADBE* promoter and the insensitive *gadA*, *gadBC*, and *hdeAB*-

TABLE 2 Biological processes enriched in the sensitive genes

		No. of genes			
			In sensitive set		
GO ID no.	GO term	In genome	Observed	Expected	P value
GO:0006970	Response to osmotic stress	13	4	0.79	0.0055
GO:0009063	Cellular amino acid catabolic process	38	6	2.31	0.0210
GO:0019395	Fatty acid oxidation	13	3	0.79	0.0385

TABLE 3 Biological processes enriched in the insensitive genes

GO ID no.	GO term	No. of genes		P value
		In genome	In insensitive set	
GO:0006412	Translation	21	3	0.018
GO:0006164	Purine nucleotide biosynthetic process	10	2	0.030
GO:0006805	Xenobiotic metabolic process	13	2	0.049

yhiD. Plasmid pLFX recombines into the lambda attachment site, placing the fusion in a novel genomic context. While we did not detect binding of RpoS upstream of *astC*, *gadA*, *gadB*, or *hdeA* by ChIP-seq, this was likely due to the difference in growth conditions, since these genes have been previously shown to be directly transcribed by RpoS (42–44).

The patterns of transcription of all four fusions were the same as observed for the respective genes in the RNA-seq data (Fig. 6A to D). *astC* transcription was sensitive to RpoS levels (one-sample *t* test, $P = 0.04$), while *gadA*, *gadB*, and *hdeA* transcription levels were all insensitive (one-sample *t* test, $P = 2 \times 10^{-6}$, $P = 10^{-5}$, $P = 0.04$, respectively). Since all reporters were placed at the same genomic locus, this result suggests that genomic location is not the determinant of the response to RpoS levels.

A second potential mechanism to explain the difference between sensitive and insensitive genes is interactions between RpoS and the core promoter sequence. For example, specific nucleotides (or combinations of nucleotides) might tend to confer sensitive or insensitive patterns of transcription. The majorities of both sensitive and insensitive genes were not associated with RpoS-bound regions in the ChIP-seq experiment, suggesting that they are indirectly regulated by RpoS (Table 4). The fact that




FIG 5 Results of testing RNA-seq expression patterns by using qPCR. Expression patterns were measured for *gadC* and *gadE* (A and B), which were deemed insensitive in the RNA-seq data, and *astA*, *prpR*, and *prpD* (C to E), which were deemed sensitive in the RNA-seq data. In all cases, the level of transcription at 26% RpoS was on the side of the line predicted by the RNA-seq data, though not always significantly. *gadC* was significantly insensitive ($P < 10^{-4}$), *gadE* was not significantly different from linear ($P = 0.10$), *astA* was marginally significantly sensitive ($P = 0.06$), *prpR* was significantly sensitive ($P < 10^{-4}$), and *prpD* was not significant ($P = 0.56$). $n = 6$. Error bars indicate the standard errors of the means.




FIG 6 Expression patterns of whole promoters and core promoters only. Expression patterns were measured for the whole upstream regulatory region (A to D) and only the core promoter (E to H) of *astC*, *gadA*, *gadB*, and *hdeA*. As expected from the RNA-seq data, the whole *astC* fusion was sensitive ($P = 0.04$; t test), and the *gadA*, *gadB*, and *hdeA* fusions were all insensitive ($P = 2 \times 10^{-6}$, $P = 10^{-5}$, and $P = 0.04$, all by one-sample t test). The P values were adjusted for multiple comparisons by using Holm's sequential adjustment method (85). The core promoters (E to H) had much lower levels of maximal transcription than the full-length promoters. All four had much-altered patterns of sensitivity, although only *gadA* and *gadB* patterns were significantly different from the whole promoter pattern ($P = 0.08$ [*astC*], $P = 0.01$ [*gadA*], $P = 10^{-6}$ [*gadB*], and $P = 0.21$ [*hdeA*], all by two-sample t test with Holm's correction). $n = 6$ to 8 . Error bars indicate standard errors of the mean results.

most sensitive and insensitive genes are not bound by RpoS argues against the hypothesis of direct RpoS-DNA interactions driving sensitivity. To see if specific sequence motifs are consistently associated with sensitivity, we used the discriminative motif search feature of DREME (45) to identify motifs that differed between sensitive and linear, or insensitive and linear, regulatory sequences. There were 28 ChIP-seq peaks associated with an operon with at least one sensitive gene, and 4 ChIP-seq peaks were associated with an operon with at least one insensitive gene. We found no motifs that distinguished these sets of sequences, although the small number of sequences would have restricted the power of such a test.

To directly test the hypothesis that the core promoter region of RpoS-transcribed genes is responsible for determining the sensitivity to RpoS levels, we cloned this short region from the *astC*, *gadA*, *gadB*, and *hdeA* promoters into pLFX and recombined the plasmids into the chromosome. These core promoters had absolute levels of transcription that were much lower than that for the entire promoter that included upstream transcription factor binding sites (Fig. 6E to H). The core promoter sequence for *astC* alone was somewhat less sensitive to RpoS levels than was the full-length construct

TABLE 4 Contingency table for the sensitivity of a gene and whether its promoter is bound by RpoS

RpoS interaction	No. of genes with transcription pattern		
	Sensitive	Linear	Insensitive
Bound by RpoS	31	87	5
Not bound by RpoS	71	823	27




FIG 7 Expression profiles of genes when cells enter stationary phase. (A) Relative expression levels of RpoS-regulated genes were obtained from a published study (47) for four time points following entry into stationary phase. All expression values were normalized to those from the final time point for each gene. The graph shows the range of relative expression values for insensitive, linear, and sensitive genes for each time point. Boxes represent the 25th to 75th percentile ranges, and horizontal lines indicate the median values. (B) Results of an experiment similar to that depicted in panel A but for a $\Delta rpoS$ strain of *E. coli*.

(two-sample *t* test, $P = 0.08$), although not significantly so, probably due to the variability of expression from the full-length *astC* reporter. The responses of the *gadA* and *gadB* core promoters differed significantly from their full-length promoters (two-sample *t* test, $P = 0.01$ and $P < 10^{-6}$, respectively). The *hdeA* core promoter (Fig. 6H) is not RpoS dependent, as it showed a decline in expression of approximately 10% in the presence of RpoS. This is consistent with the previous finding that the ability of RpoD (but not RpoS) to transcribe *hdeA* is repressed by the H-NS protein bound upstream of the promoter (46). The core promoter construct lacks the native H-NS binding sites upstream, and so the selectivity is apparently lost. Thus, the core promoters do not replicate the RpoS sensitivity of their whole promoter sequences. In addition, the three RpoS-dependent core promoters (Fig. 6E to G) do not differ from each other in sensitivity (analysis of variance [ANOVA], $P = 0.32$). We conclude that the core promoter is not responsible for RpoS sensitivity.

Experiments with the *lacZ* fusions suggested that neither genomic location nor the core promoter sequence influences the sensitivity of a promoter. The remaining possible mechanism is the binding of specific TFs. If this is the case, we would expect that sensitive and insensitive genes are enriched for binding by different TFs. We looked for such enrichment and found that the sensitive genes were enriched for binding by ArgR, Nac, and NtrC (false-discovery rate [FDR], <0.05). The insensitive genes were enriched for binding by ArcA, FlhZ, GadE, GadW, GadX, H-NS, PhoP, RcsB, and TorR. This very large set of regulators occurs largely because these proteins are all annotated as regulating some or all of the operons involved in AR2: *gadA*, *gadBC*, *gadE*, *hdeAB-yhiD*, and *hdeD*. The actions of GadE, GadW, and GadX occur primarily at these loci, while the other regulators have many additional known binding sites that are not near the promoters of insensitive genes. This specific enrichment for TFs highlights proteins that may be responsible for the sensitive or insensitive patterns of expression.

Sensitivity to RpoS determines the timing of induction during entry into stationary phase. We hypothesized that the degree of sensitivity to RpoS impacts the timing of expression under conditions when levels of active RpoS increase, such as during entry into stationary phase. A previous study used RNA-seq to monitor the transcriptome over a time course of growth, including four time points during stationary phase (47). We analyzed these data to determine whether insensitive, linear, and sensitive genes showed differences in the timing of induction. We selected only those RpoS-induced genes whose transcription increased upon entry into stationary phase. (In total, there were 250 such linear genes, 90 sensitive genes, and 19 insensitive genes.) We then determined the pattern of expression for each such gene over four time points beginning at the onset of stationary phase. Although there was considerable variability in the expression patterns of genes, as a group the three classes showed clear differences in the timing of induction (Fig. 7A). Specifically, sensitive genes were induced most rapidly, followed closely by linear genes. In both cases, expression

peaked early in stationary phase and then decreased between 30 and 180 min after entering stationary phase. In contrast, insensitive genes showed relatively little change in expression until the final time point, 180 min into stationary phase. To determine the importance of RpoS on the patterns of gene expression, we repeated the above analysis using data generated from a $\Delta rpoS$ strain. As expected, the large difference in timing between the groups of genes was greatly diminished (Fig. 7B). We conclude that the sensitivity of a gene is associated with the timing of expression during stationary phase in an RpoS-dependent manner.

DISCUSSION

Expanding the known set of RpoS-transcribed genes and promoters. The *E. coli* RpoS regulon has been widely investigated using targeted and genome-scale approaches. Most genome-scale studies have focused on genes whose expression is altered in the absence of RpoS (18–20, 27, 28). Hence, these studies cannot distinguish between genes that are transcribed by RpoS and those that are indirectly regulated. ChIP-seq affords a high-resolution view of RpoS binding. By combining ChIP-seq with RNA-seq, we identified 123 RpoS-transcribed genes with high confidence, considerably expanding the known RpoS regulon. Previous studies have suggested a role for RpoS in direct repression of some target genes (25, 29, 48). While we observed negative regulation of 439 genes by RpoS, there were fewer of these repressed genes associated with ChIP-seq peaks than expected by chance, suggesting that direct negative regulation by RpoS is rare.

Only two previous studies have used ChIP methods to map RpoS binding genome-wide in *E. coli*. The first study used ChIP-chip to identify 868 RpoS-bound regions (25), many more than were identified in our study but at considerably lower resolution (median peak length of 324 bp for RpoS ChIP-chip). The second study used ChIP-seq but identified relatively few RpoS-bound regions (26). Of the 63 RpoS-bound regions identified in the second study, 41 were shared with those from our study.

The high resolution of ChIP-seq allowed us to identify specific promoter sequences recognized by RpoS. By combining ChIP-seq data with a TSS map, we identified many high-confidence RpoS promoters. These promoters are strongly enriched for the presence of a -10 hexamer, with sequence preferences consistent with several of the previously described features of RpoS promoters (22). Specifically, we observed a preference for a C at position -8 (within the -10 hexamer), a C at position -13 (immediately upstream of the -10 hexamer), and a TAA at positions -6 to -4 (immediately downstream of the -10 hexamer). Previous studies have suggested that RpoS promoters often contain a -35 hexamer (49), although the spacing relative to the -10 hexamer is considerably more variable than for σ^{70} promoters. However, we did not detect enrichment of a -35 hexamer-like sequence among RpoS promoters, suggesting that the requirement for this element is weak.

Sensitivity to RpoS affects timing of expression and groups of functionally related genes. As is true for many transcription factors, RpoS levels vary continuously across a wide range of conditions. Our data show that genes differ in the sensitivity of their response to RpoS levels. Moreover, whether a gene is sensitive or insensitive to RpoS levels is associated with its function, suggesting a physiological rationale for sensitivity. For example, genes that are insensitive include many of those involved in the glutamate-dependent acid resistance 2 system (AR2). These genes are of particular interest because AR2 allows *E. coli* to survive a pH of 2, an important trait for its ability to pass through the stomach and colonize the gastrointestinal tract (50, 51). To some extent, the shared sensitivity of functionally related genes can be explained by operon structure, i.e., cotranscription of multiple functionally related genes from a single promoter. However, the phenomenon of shared sensitivity for functionally related genes extends beyond operons. For example, the insensitive genes involved in AR2 are transcribed from at least five different promoters (40).

The fact that functionally related genes often have similar patterns of sensitivity to RpoS suggests that sensitivity can serve as a mechanism to control the timing of gene

expression and hence to coordinate specific cellular processes as part of a response to environmental stresses. Consistent with this idea, we have shown that the sensitivity of genes to RpoS levels correlates with the timing of their expression. RpoS sensitivity may drive similar patterns of expression in response to other stresses. Different environmental stresses are known to upregulate RpoS to varied levels (8), suggesting that some insensitive genes may only be expressed under certain stresses. In addition to the effects on gene expression, sensitivity to RpoS may also impact the effects of mutations in *rpoS*, which have been seen to evolve in the lab (52, 53). We expect mutations attenuating RpoS have the strongest effect on insensitive genes and the weakest effect on sensitive genes.

Possible mechanisms of RpoS sensitivity. While the connection between RpoS sensitivity and the timing of gene expression is clear, the molecular basis of sensitivity is less so. Our data indicate that the genomic location of these operons does not determine the expression pattern. Moreover, several lines of evidence suggest that direct interactions between RpoS and the core promoter are also not responsible for determining sensitivity. First, analysis of the ChIP-seq data for the sensitive and insensitive genes found no motif that distinguishes between them. Second, core promoters from both sensitive and insensitive genes did not replicate the pattern of expression of the full-length promoters. Third, the core promoters of a sensitive operon (*astC*) and two insensitive operons (*gadA* and *gadB*) had indistinguishable patterns of sensitivity, suggesting that what was excluded from those constructs (i.e., binding sites of regulatory proteins) determines the shape of the relationship.

Given our finding that core promoter sequences cannot explain the difference in sensitivity between promoters, we suggest that sensitivity is largely due to the action of specific regulatory proteins bound upstream. If this hypothesis is correct, it could also explain the physiological coherence of these groups. For example, many insensitive genes are involved in the AR2 phenotype and are also regulated by GadX, GadW, and GadE (54–56). If one or more of these three regulators is directly responsible for the insensitive pattern of expression, then this could help to explain the physiological coherence of the insensitive group. The sensitive genes, being a larger group, have no obvious single regulator, although relatively little is known about regulators that function in stationary phase.

It is also possible that the physical properties of promoters play a role in this process, either alone or in concert with transcription factor binding. For example, the supercoiling state of promoters influences levels of transcription (57, 58) and has been implicated in regulating RpoS-dependent transcription (59, 60). It is possible that differential responses of promoters to supercoiling levels, either due to interactions directly with RpoS or to changes in the ability of transcription factors to bind (61), play a role in determining the sensitivity of a promoter. If this type of regulation plays a role, it must be due to the structure of the whole promoter itself, rather than supercoiling differences conferred by genomic location (62). We know this because the full promoters cloned into *lacZ* fusions were able to replicate both sensitive and insensitive patterns of transcription, even when moved to the same chromosomal location.

RpoS responds to a wide variety of environmental cues and regulates genes responsible for many different kinds of responses. This work has demonstrated that one facet of that response, the level of RpoS produced, has varied effects across the entirety of the regulon. The level of RpoS produced in a stress response, together with the action of other transcription factors, may help to tune the RpoS-dependent stress response in ways appropriate for individual stresses.

MATERIALS AND METHODS

Culture conditions. Cells were grown in 5 ml of LB broth (1% tryptone, 0.5% yeast extract, 1% NaCl) in 150- by 18-mm tubes, positioned vertically and shaking at 225 rpm at 37°C in a water bath, unless otherwise specified in the text. When required, antibiotics were used at the following final concentrations: ampicillin at 100 µg/ml (for plasmids) or 25 µg/ml (for chromosomal integration); chloramphenicol at 20 µg/ml; kanamycin at 50 µg/ml.

Strains and genetic manipulations. Strains and plasmids used in this study are listed in Table S1 in the supplemental material. The wild-type genetic background for all experiments except for ChIP-seq was that of strain BW27786, a strain designed to give a graded transcriptional response to increasing arabinose concentrations (35). To create a strain of this background lacking *rpoS*, the $\Delta rpoS746::kan$ allele of JW5437 (63) was moved by P1 transduction into BW27786, creating strain DMS2545.

The arabinose-inducible RpoS strain was created by PCR amplifying the *kan* gene and the P_{araB} promoter of plasmid pAH150 (64) using primers ParaBRpoSRecomb-F and ParaBRpoSRecomb-R (Table S2). This PCR product was then integrated into the *nlpD* gene (i.e., 5' of *rpoS*) in a MG1655 background using plasmid pKD46 (65) and P1 transduced into BW27786, creating strain DMS2564. This strain thus lacked both the native transcriptional and translational control of RpoS.

For flow cytometry experiments, plasmid pDMS123 (36), which contains a *otsB-gfp* transcriptional fusion, was transformed into DMS2564 by the method described by Groth et al. (66).

ChIP-seq experiments used strain RPB104, an unmarked derivative of MG1655 that expresses a C-terminally SPA-tagged derivative of RpoS from its native locus. This strain was constructed by P1 transduction of Kan^r-linked *rpoS*-SPA from a previously described strain (67). The Kan^r cassette was removed using the pCP20 plasmid, which encodes Flp recombinase (65).

Construction and chromosomal integration of *lacZ* fusions. The *gadB*- and *astC-lacZ* fusion plasmids were built by using standard cloning methods. The *gadC* and *astA* promoter regions with transcription factor binding sites were PCR amplified with primers gadCpromoter+/− and astApromoter+/− (Table S2), which included KpnI and EcoRI restriction sites for cloning. Cloning of core promoter regions was performed by annealing oligonucleotides designed to contain the whole RpoS binding region, as predicted by Fraley et al. (68) and Castanie-Cornet and Foster (69). Oligonucleotides were annealed by heating 1 μ M forward and reverse primer for 1 min at 100°C with 5 mM MgCl₂ and 7 mM Tris-HCl and then cooling slowly to room temperature. Inserts and plasmid (pLFX) were digested with EcoRI-HF and KpnI-HF (New England BioLabs), ligated with T7 ligase (New England BioLabs), and cloned into strain BW23473. Transformants were minipreped, and inserts were verified by Sanger sequencing.

The *gadA*- and *hdeA-lacZ* fusion plasmids were built using Gibson assembly (70) with the NEBuilder HiFi assembly kit (New England BioLabs). The *gadA* and *hdeA* promoter regions were PCR amplified with primers hdeAHiFi+/− and gadAHiFi+/−. The core promoter was cloned with the single long oligonucleotide hdeAcoreHiFi or gadAcoreHiFi, as predicted by Arqvist et al. (42) and De Biase et al. (71). PCR products or oligonucleotides were mixed with pLFX digested with KpnI-HF and EcoRI-HF and assembled according to the manufacturer's instructions. Mixtures were cloned into strain BW23473, transformants were minipreped, and inserts were verified by Sanger sequencing.

lacZ fusion plasmids were integrated into strain DMS2564 with helper plasmid pPFINT (41). Single-copy integrants were confirmed using the PCR assay of Haldimann and Wanner (64).

Quantitative Western blotting. Quantitative Western blotting was used to measure RpoS levels. Cells were inoculated from frozen cultures into 5 ml of LB and grown overnight at 37°C, shaken at 225 rpm. Five microliters of this overnight culture was diluted into 5 ml of LB with the appropriate concentration of arabinose and grown for 20 h. A 100- μ l aliquot of the overnight culture to be assayed was centrifuged and resuspended in 1 \times Laemmli sample buffer (Sigma-Aldrich) and boiled for 5 min. Samples were diluted 1:10 in 1 \times Laemmli buffer, and 10 μ l was electrophoresed on a 10% polyacrylamide gel (Bio-Rad) in Tris-glycine running buffer (25 mM Tris base, 250 mM glycine, 0.5% SDS) at 100 V for 90 min at room temperature. Proteins were transferred to an Immobilon-FL polyvinylidene difluoride membrane by electrophoresis at 100 V for 45 min at 4°C in transfer buffer (48 mM Tris base, 39 mM glycine, 20% methanol, and 0.0375% SDS). Membranes were blocked by overnight incubation in Odyssey blocking buffer (Li-Cor) at 4°C.

The blocked membrane was probed with affinity-purified monoclonal antibodies to RpoS (clone 1RS1) and RpoD (clone 2G10) (NeoClone) at a final concentration of 0.4 μ g/ml in Odyssey blocking buffer plus 0.2% Tween 20 for 1 h at room temperature. The membrane was washed four times for 5 min each with 15 ml of 1 \times Tris-buffered saline with Tween 20. A fluorescent secondary antibody (IRDye 800CW goat anti-mouse antibody; Li-Cor) was diluted 1:10,000 in a solution of Odyssey blocking buffer plus 0.2% Tween 20 and 0.01% SDS and incubated in the dark for 1 h at room temperature. The membrane was washed as before, dried for 2 h between sheets of Whatmann 3MM blotting paper, and imaged on a LiCor Clx fluorescent imager.

Band intensity was estimated using the Image Studio 2.1 package (LiCor). RpoS levels were divided by RpoD levels to normalize for differences in total protein levels. The ratio of RpoS to RpoD is biologically meaningful because this ratio, rather than the RpoS level alone, dictates levels of transcription from RpoS-dependent promoters due to sigma factor competition (72).

RNA-seq experiments and analysis. Cells were inoculated from frozen cultures into 5 ml of LB and grown overnight. Five-microliter aliquots of this overnight culture were diluted into 5 ml of LB and grown for 20 h. (For the intermediate 26% RpoS condition, a final concentration of 10^{−4}% arabinose was also added.) RNA was purified from 200 μ l of overnight culture by pelleting and resuspending in 500 μ l of TRIzol at 65°C, followed by purification on a column (Direct-Zol; Zymo Research). Samples received two 30-min DNase treatments using Turbo DNA-free (Ambion) following the manufacturer's instructions. RNA samples were then purified on a column (RNA Clean and Concentrator; Zymo Research). Samples were stored at −80°C until use. Three samples were prepped from each culture and pooled to generate sufficient RNA. Two biological replicates were prepared for each strain or condition of interest. rRNA depletion, cDNA synthesis, library preparation, and sequencing were performed by a commercial

provider (Otogenetics, Norcross, GA). Paired-end, 100-bp sequences were generated for 7 to 15 million reads per sample.

Before reads were mapped, the first 10 bp of each read were trimmed using the FASTX toolkit 0.0.13. Reads were mapped to the NCBI K-12 reference genome (NC_000913.2 *Escherichia coli* strain K-12 substrain MG1655) by using the BWA program v 0.7.5a (73). The number of read pairs mapped to each gene was counted with HTSeq 0.6.1 (74). Differential expression analysis was performed with DESeq v2.13 (75). All *P* values were first FDR adjusted using the procedure of Benjamini and Hochberg (76). *P* values were then further Bonferroni adjusted for the three comparisons between pairs of RpoS levels. All differential expression *P* values reported in this paper reflect both the FDR and Bonferroni adjustments.

To determine if a gene differed significantly from the null expectation of linearity, we calculated the probability of the observed read count value at 26% RpoS if the true expression level was given by the linear prediction between the endpoints. Note that to calculate both the expected read count and the probability of the observed value under the null hypothesis (i.e., the *P* value), our model required estimating both the DESeq size factor (for scaling) and the dispersion (a variance factor) for each of the samples. The negative binomial probability model (routinely used to measure count data [75]) was used, with the size factors and dispersion estimated from DESeq (75) to calculate the probability of the observed read count at 26% RpoS.

GO term analysis was performed using the topGO package (77) together with the org.Eck12.eg.db annotation package (78) in R 3.1.0. Enrichment was assessed using the weight01 algorithm (77) together with Fisher's exact test. The GO hierarchy was pruned to include only nodes with at least five associated genes, as significance tests can be unstable for GO terms with fewer genes (77).

A Venn diagram of the number of significant genes under each condition was prepared with EulerAPE 3.0.0 (79).

Analysis of transcription factor binding site enrichment used data from RegulonDB (24). We divided RpoS-regulated transcriptional units in RegulonDB into sets: sensitive, insensitive, and linear. Then, for each transcription factor in the database, we determined how many of the transcription units it regulates fell into each set. We compared this with the number of regulated transcription units within the whole RpoS regulon. A case of enrichment is where a transcription factor regulates a disproportionately large number of units in a particular set (e.g., the sensitive set) compared with what would be expected based on the RpoS regulon as a whole. We identified such cases using a one-tailed hypergeometric test. All *P* values were then FDR adjusted using the procedure of Benjamini and Hochberg (76). Transcription factors with an FDR-adjusted *P* value of <0.05 that regulated at least two transcription units in the gene set were considered enriched.

ChIP-seq analysis. *E. coli* strain RPB104 (MG1655 with C-terminally SPA-tagged *rpoS*) was grown overnight in M9 minimal medium with 0.4% glycerol at 30°C, then subcultured 1:100 in the same medium, and grown for 60 h to saturation (optical density at 600 nm of ~3). ChIP-seq using the M2 monoclonal anti-FLAG antibody was performed as described previously (80). Regions of enrichment (peaks) were identified as described previously (23). Relative enrichment is reported as the fold above threshold (FAT) score.

We used MEME-ChIP (version 4.11.2; default parameters) to analyze enriched regions identified by ChIP-seq (81). Regions within 100 bp were merged. The reported sequence motif was identified by MEME (version 4.11.2) (82), run within the MEME-ChIP environment.

We used MEME (82) within the MEME-ChIP environment (version 4.11.2), with default parameters except that only the given strand was analyzed, to identify enriched sequence motifs in regions surrounding TSSs associated with RpoS ChIP-seq peaks. We analyzed sequences from -45 to +5 relative to each TSS.

We directly identified RpoS-transcribed genes by requiring that (i) the gene start was within 300 bp of a ChIP-seq peak, (ii) the gene was positively regulated by RpoS, as determined by RNA-seq, (iii) no other positively regulated gene started within 300 bp of the ChIP-seq peak, and (iv) there was no associated sequence motif or TSS (identified in reference 25) that would be consistent with transcription in the opposite orientation. For regulated genes, we determined whether other genes in the same operon were also regulated, using a published operon list (47) for *E. coli*. Peaks were associated with transcription start sites from a previous study (25) by identifying transcription start sites within 20 bp of a peak.

qPCR. RNA was isolated as for RNA-seq, except that samples underwent three 30-min DNase treatments using Turbo DNA-free (Ambion). cDNA was made from the RNA samples using SuperScript Vilo master mix (Invitrogen) and stored at -20°C for use in real-time qPCRs.

qPCR was performed using Power SYBR green master mix (Invitrogen), 2 µl of cDNA, and primers at 300 nM. Cycling was performed at 95°C for 10 min followed by 40 cycles of 95°C for 15 s, 58°C for 15 s, and 68°C for 30 s. Three control genes (*ftsZ*, *pgm*, and *hemL*) were used in addition to genes of interest. These genes were selected using the approach of Vandesompele et al. (83). Details of this selection process, including the other seven genes tested, are available in Text S1, Table S3, and Fig. S1.

For each gene, a standard curve was made using the following amounts of genomic DNA: 200 ng, 20 ng, 2 ng, 200 pg, 20 pg, and 2 pg. Genomic DNA was extracted from overnight cultures using a Puregene kit (Qiagen), following the manufacturer's instructions. Expression levels for each gene were interpolated from the standard curve. Expression levels of experimental genes were divided by the geometric means of the levels for the three control genes.

Statistical assessment of sensitivity was performed by a bootstrapping approach. Bootstrapping, rather than a parametric approach, was appropriate because the data did not conform to parametric assumptions. In this approach, the unit of resampling was the RNA isolated from the 0%, 26%, and 100%

RpoS conditions on an individual day. For each resampled data set, the median expression at 0% and 100% RpoS was calculated. The linear fit from the 0% RpoS median and the 100% RpoS median was then used to predict the level of expression at the intermediate level of RpoS. Repeating this resampling of the data 10,000 times yielded a 95% confidence interval for 26% RpoS. The observed median for 26% was then compared to the confidence interval to test for significance.

Beta-galactosidase activity. Beta-galactosidase activity was measured using the method of Miller (84). With *lacZ* fusion strains, the level of sensitivity was quantified, rather than categorized only as sensitive, linear, or insensitive. To quantify the level of sensitivity, for each replicate we calculated the distance between the observed expression at the intermediate RpoS concentration and the expected level based on a linear pattern, standardized by the difference in expression between high and low RpoS conditions. Testing if sensitivities were different from zero was performed with a one-sample *t* test, with *P* values adjusted for multiple comparisons using Holm's sequential adjustment method (85). Testing if two sensitivities were different relied on a two-sample *t* test with the same method of adjustment for multiple comparisons.

Analysis of published RNA-seq data. We analyzed published RNA-seq data for wild-type and the $\Delta rpoS$ mutant *E. coli* strain over a time course of growth into stationary phase (47). Using normalized genome coverage information extracted from wiggle files, we calculated relative abundance for all genes at each of the four stationary-phase time points in the growth curves (the last four time points for each strain). We arbitrarily selected a threshold coverage value of 500 and excluded any genes scoring below this threshold at all four time points in wild-type cells. This reduced variability associated with low expression levels. We excluded any genes for which coverage at the final time point was 0, since this would have prevented normalization. We also excluded any gene for which the first stationary-phase time point had the highest expression value of the four time points for wild-type cells, since RpoS-dependent expression of these genes is likely to be masked by other factors. We then selected genes whose expression we had found to be induced by RpoS and separated these genes into insensitive, linear, and sensitive classes. We calculated expression levels for each of these genes relative to expression at the final time point.

Accession number(s). RNA-seq data were deposited in the GEO database with accession number GSE87856. ChIP-seq data were deposited in ArrayExpress with accession number E-MTAB-5339.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/JB.00755-16>.

TEXT S1, PDF file, 0.5 MB.

DATA SET S1, XLSX file, 0.2 MB.

DATA SET S2, XLSX file, 0.03 MB.

ACKNOWLEDGMENTS

We thank Rachael Kretsch for experimental help and Robert Drewell, Xuelin Wu, Jae Hur, Keith Derbyshire, and Todd Gray for helpful discussions.

This work was supported by HHMI Undergraduate Science Education award 52007544 to Harvey Mudd College and by the NIH Director's New Innovator Award Program grant 1DP2OD007188 (to J.T.W.).

REFERENCES

- Novick A, Weiner M. 1957. Enzyme induction as an all-or-none phenomenon. *Proc Natl Acad Sci U S A* 43:553–566. <https://doi.org/10.1073/pnas.43.7.553>.
- Ptashne M. 2004. Phage lambda revisited, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Brinsmade SR, Alexander EL, Livny J, Stettner AI, Segrè D, Rhee KY, Sonenshein AL. 2014. Hierarchical expression of genes controlled by the *Bacillus subtilis* global regulatory protein CodY. *Proc Natl Acad Sci U S A* 111:8227–8232. <https://doi.org/10.1073/pnas.1321308111>.
- Battesti A, Majdalani N, Gottesman S. 2011. The RpoS-mediated general stress response in *Escherichia coli*. *Annu Rev Microbiol* 65:189–213. <https://doi.org/10.1146/annurev-micro-090110-102946>.
- Hengge R. 16 December 2011. Stationary-phase gene regulation in *Escherichia coli*. *EcoSal Plus* <https://doi.org/10.1128/ecosalplus.5.6.3>.
- Schellhorn HE. 2014. Elucidating the function of the RpoS regulon. *Future Microbiol* 9:497–507. <https://doi.org/10.2217/fmb.14.9>.
- Lange R, Hengge-Aronis R. 1994. The cellular concentration of the sigma S subunit of RNA polymerase in *Escherichia coli* is controlled at the levels of transcription, translation, and protein stability. *Genes Dev* 8:1600–1612. <https://doi.org/10.1101/gad.8.13.1600>.
- Mandel MJ, Silhavy TJ. 2005. Starvation for different nutrients in *Escherichia coli* results in differential modulation of RpoS levels and stability. *J Bacteriol* 187:434–442. <https://doi.org/10.1128/JB.187.2.434-442.2005>.
- Zafar MA, Carabetta VJ, Mandel MJ, Silhavy TJ. 2014. Transcriptional occlusion caused by overlapping promoters. *Proc Natl Acad Sci U S A* 111:1557–1561. <https://doi.org/10.1073/pnas.1323413111>.
- Hengge R. 2009. Proteolysis of σ^S (RpoS) and the general stress response in *Escherichia coli*. *Res Microbiol* 160:667–676. <https://doi.org/10.1016/j.resmic.2009.08.014>.
- Pratt LA, Silhavy TJ. 1998. Crl stimulates RpoS activity during stationary phase. *Mol Microbiol* 29:1225–1236. <https://doi.org/10.1046/j.1365-2958.1998.01007.x>.
- Hryckowian AJ, Battesti A, Lemke JJ, Meyer ZC, Welch RA. 2014. IraL is an RssB anti-adaptor that stabilizes RpoS during logarithmic phase growth in *Escherichia coli* and *Shigella*. *mBio* 5:e01043-14. <https://doi.org/10.1128/mBio.01043-14>.
- Chiang SM, Dong T, Edge TA, Schellhorn HE. 2011. Phenotypic diversity caused by differential RpoS activity among environmental *Escherichia coli* isolates. *Appl Environ Microbiol* 77:7915–7923. <https://doi.org/10.1128/AEM.05274-11>.
- Bhagwat AA, Tan J, Sharma M, Kothary M, Low S, Tall BD, Bhagwat M. 2006. Functional heterogeneity of RpoS in stress tolerance of enteroh-

- morrhagic *Escherichia coli* strains. *Appl Environ Microbiol* 72:4978–4986. <https://doi.org/10.1128/AEM.02842-05>.
15. Ferenci T, Galbiati HF, Betteridge T, Phan K, Spira B. 2011. The constancy of global regulation across a species: the concentrations of ppGpp and RpoS are strain-specific in *Escherichia coli*. *BMC Microbiol* 11:62. <https://doi.org/10.1186/1471-2180-11-62>.
 16. Snyder E, Gordon DM, Stoebel DM. 2012. *Escherichia coli* lacking RpoS are rare in natural populations of non-pathogens. *G3* 2:1341–1344. <https://doi.org/10.1534/g3.112.003855>.
 17. Bleibtreu A, Clermont O, Darlu P, Glodt J, Branger C, Picard B, Denamur E. 2014. The *rpoS* gene is predominantly inactivated during laboratory storage and undergoes source-sink evolution in *Escherichia coli* species. *J Bacteriol* 196:4276–4284. <https://doi.org/10.1128/JB.01972-14>.
 18. Patten CL, Kirchhof MG, Schertzberg MR, Morton RA, Schellhorn HE. 2004. Microarray analysis of RpoS-mediated gene expression in *Escherichia coli* K-12. *Mol Genet Genomics* 272:580–591. <https://doi.org/10.1007/s00438-004-1089-2>.
 19. Weber H, Polen T, Heuveling J, Wendisch VF, Hengge R. 2005. Genome-wide analysis of the general stress response network in *Escherichia coli*: σ^S -dependent genes, promoters, and σ factor selectivity. *J Bacteriol* 187:1591–1603. <https://doi.org/10.1128/JB.187.5.1591-1603.2005>.
 20. Dong T, Schellhorn HE. 2009. Control of RpoS in global gene expression of *Escherichia coli* in minimal media. *Mol Genet Genomics* 281:19–33. <https://doi.org/10.1007/s00438-008-0389-3>.
 21. Lévi-Meyrueis C, Monteil V, Sismeiro O, Dillies M-A, Monot M, Jagla B, Coppée J-Y, Dupuy B, Norel F. 2014. Expanding the RpoS/ σ^S -network by RNA sequencing and identification of σ^S -controlled small RNAs in *Salmonella*. *PLoS One* 9:e96918. <https://doi.org/10.1371/journal.pone.0096918>.
 22. Typas A, Becker G, Hengge R. 2007. The molecular basis of selective promoter activation by the σ^S subunit of RNA polymerase. *Mol Microbiol* 63:1296–1306. <https://doi.org/10.1111/j.1365-2958.2007.05601.x>.
 23. Fitzgerald DM, Bonocora RP, Wade JT. 2014. Comprehensive mapping of the *Escherichia coli* flagellar regulatory network. *PLoS Genet* 10:e1004649. <https://doi.org/10.1371/journal.pgen.1004649>.
 24. Salgado H, Peralta-Gil M, Gama-Castro S, Santos-Zavaleta A, Muñoz-Rascado L, García-Sotelo JS, Weiss V, Solano-Lira H, Martínez-Flores I, Medina-Rivera A, Salgado-Orsorio G, Alquicira-Hernández S, Alquicira-Hernández K, López-Fuentes A, Porrón-Sotelo L, Huerta AM, Bonavides-Martínez C, Balderas-Martínez YI, Pannier L, Olvera M, Labastida A, Jiménez-Jacinto V, Vega-Alvarado L, Del Moral-Chávez V, Hernández-Alvarez A, Moret E, Collado-Vides J. 2013. RegulonDB v8.0: omics data sets, evolutionary conservation, regulatory phrases, cross-validated gold standards and more. *Nucleic Acids Res* 41:D203–D213. <https://doi.org/10.1093/nar/gks1201>.
 25. Cho B-K, Kim D, Knight EM, Zengler K, Palsson BO. 2014. Genome-scale reconstruction of the sigma factor network in *Escherichia coli*: topology and functional states. *BMC Biol* 12:4. <https://doi.org/10.1186/1741-7007-12-4>.
 26. Peano C, Wolf J, Demol J, Rossi E, Petiti L, De Bellis G, Geiselmann J, Egli T, Lacour S, Landini P. 2015. Characterization of the *Escherichia coli* σ^S core regulon by chromatin immunoprecipitation-sequencing (ChIP-seq) analysis. *Sci Rep* 5:10469. <https://doi.org/10.1038/srep10469>.
 27. Lacour S, Landini P. 2004. σ^S -dependent gene expression at the onset of stationary phase in *Escherichia coli*: function of σ^S -dependent genes and identification of their promoter sequences. *J Bacteriol* 186:7186–7195. <https://doi.org/10.1128/JB.186.21.7186-7195.2004>.
 28. Dong T, Kirchhof MG, Schellhorn HE. 2008. RpoS regulation of gene expression during exponential growth of *Escherichia coli* K12. *Mol Genet Genomics* 279:267–277. <https://doi.org/10.1007/s00438-007-0311-4>.
 29. Rosenthal AZ, Kim Y, Gralla JD. 2008. Poising of *Escherichia coli* RNA polymerase and its release from the σ^{38} C-terminal tail for *osmY* transcription. *J Mol Biol* 376:938–949. <https://doi.org/10.1016/j.jmb.2007.12.037>.
 30. Reppas NB, Wade JT, Church GM, Struhl K. 2006. The transition between transcriptional initiation and elongation in *E. coli* is highly variable and often rate limiting. *Mol Cell* 24:747–757. <https://doi.org/10.1016/j.molcel.2006.10.030>.
 31. Schröder O, Wagner R. 2000. The bacterial DNA-binding protein H-NS represses ribosomal RNA transcription by trapping RNA polymerase in the initiation complex1. *J Mol Biol* 298:737–748. <https://doi.org/10.1006/jmbi.2000.3708>.
 32. Rosenthal AZ, Hu M, Gralla JD. 2006. Osmolyte-induced transcription: –35 region elements and recognition by σ^{38} (rpoS). *Mol Microbiol* 59:1052–1061. <https://doi.org/10.1111/j.1365-2958.2005.04999.x>.
 33. Huo Y-X, Rosenthal AZ, Gralla JD. 2008. General stress response signalling: unwrapping transcription complexes by DNA relaxation via the σ^{38} C-terminal domain. *Mol Microbiol* 70:369–378. <https://doi.org/10.1111/j.1365-2958.2008.06412.x>.
 34. Bonocora RP, Smith C, Lapierre P, Wade JT. 2015. Genome-scale mapping of *Escherichia coli* σ^{54} reveals widespread, conserved intragenic binding. *PLoS Genet* 11:e1005552. <https://doi.org/10.1371/journal.pgen.1005552>.
 35. Khlebnikov A, Datsenko KA, Skaug T, Wanner BL, Keasling JD. 2001. Homogeneous expression of the P_{BAD} promoter in *Escherichia coli* by constitutive expression of the low-affinity high-capacity AraE transporter. *Microbiology* 147:3241–3247. <https://doi.org/10.1099/00221287-147-12-3241>.
 36. Stoebel DM, Hokamp K, Last MS, Dorman CJ. 2009. Compensatory evolution of gene regulation in response to stress by *Escherichia coli* lacking RpoS. *PLoS Genet* 5:e1000671. <https://doi.org/10.1371/journal.pgen.1000671>.
 37. Mates AK, Sayed AK, Foster JW. 2007. Products of the *Escherichia coli* acid fitness island attenuate metabolite stress at extremely low pH and mediate a cell density-dependent acid resistance. *J Bacteriol* 189:2759–2768. <https://doi.org/10.1128/JB.01490-06>.
 38. Kanjee U, Houry WA. 2013. Mechanisms of acid resistance in *Escherichia coli*. *Annu Rev Microbiol* 67:65–81. <https://doi.org/10.1146/annurev-micro-092412-155708>.
 39. Bryant JA, Sellars LE, Busby SJW, Lee DJ. 2014. Chromosome position effects on gene expression in *Escherichia coli* K-12. *Nucleic Acids Res* 42:11383–11392. <https://doi.org/10.1093/nar/gku828>.
 40. Keseler IM, Mackie A, Peralta-Gil M, Santos-Zavaleta A, Gama-Castro S, Bonavides-Martínez C, Fulcher C, Huerta AM, Kothari A, Krummenacker M, Latendresse M, Muñoz-Rascado L, Ong Q, Paley S, Schröder I, Shearer AG, Subhraveti P, Travers M, Weerasinghe D, Weiss V, Collado-Vides J, Gunsalus RP, Paulsen I, Karp PD. 2013. EcoCyc: fusing model organism databases with systems biology. *Nucleic Acids Res* 41:D605–D612. <https://doi.org/10.1093/nar/gks1027>.
 41. Edwards AN, Patterson-Fortin LM, Vakulskas CA, Mercante JW, Potrykus K, Vinella D, Camacho MI, Fields JA, Thompson SA, Georgellis D, Cashel M, Babitzke P, Romeo T. 2011. Circuitry linking the Csr and stringent response global regulatory systems. *Mol Microbiol* 80:1561–1580. <https://doi.org/10.1111/j.1365-2958.2011.07663.x>.
 42. Arqvist A, Olsén A, Normark S. 1994. Sigma S-dependent growth-phase induction of the *csgBA* promoter in *Escherichia coli* can be achieved in vivo by sigma 70 in the absence of the nucleoid-associated protein H-NS. *Mol Microbiol* 13:1021–1032. <https://doi.org/10.1111/j.1365-2958.1994.tb00493.x>.
 43. Kiupakis AK, Reitzer L. 2002. ArgR-independent induction and ArgR-dependent superinduction of the *astCADBE* operon in *Escherichia coli*. *J Bacteriol* 184:2940–2950. <https://doi.org/10.1128/JB.184.11.2940-2950.2002>.
 44. Waterman SR, Small PLC. 2003. Transcriptional expression of *Escherichia coli* glutamate-dependent acid resistance genes *gadA* and *gadBC* in an *hns rpoS* mutant. *J Bacteriol* 185:4644–4647. <https://doi.org/10.1128/JB.185.15.4644-4647.2003>.
 45. Bailey TL. 2011. DREME: motif discovery in transcription factor ChIP-seq data. *Bioinformatics* 27:1653–1659. <https://doi.org/10.1093/bioinformatics/btr261>.
 46. Shin M, Song M, Rhee JH, Hong Y, Kim Y-J, Seok Y-J, Ha K-S, Jung S-H, Choy HE. 2005. DNA looping-mediated repression by histone-like protein H-NS: specific requirement of $E\sigma^{70}$ as a cofactor for looping. *Genes Dev* 19:2388–2398. <https://doi.org/10.1101/gad.1316305>.
 47. Conway T, Creecy JP, Maddox SM, Grissom JE, Conkle TL, Shadid TM, Teramoto J, San Miguel P, Shimada T, Ishihama A, Mori H, Wanner BL. 2014. Unprecedented high-resolution view of bacterial operon architecture revealed by RNA sequencing. *mBio* 5:e01442-14. <https://doi.org/10.1128/mBio.01442-14>.
 48. Lévi-Meyrueis C, Monteil V, Sismeiro O, Dillies M-A, Kolb A, Monot M, Dupuy B, Duarte SS, Jagla B, Coppée J-Y, Beraud M, Norel F. 2015. Repressor activity of the RpoS/ σ^S -dependent RNA polymerase requires DNA binding. *Nucleic Acids Res* 43:1456–1468. <https://doi.org/10.1093/nar/gku1379>.
 49. Typas A, Hengge R. 2006. Role of the spacer between the –35 and –10 regions in σ^S promoter selectivity in *Escherichia coli*. *Mol Microbiol* 59:1037–1051. <https://doi.org/10.1111/j.1365-2958.2005.04998.x>.

50. Richard HT, Foster JW. 2003. Acid resistance in *Escherichia coli*. *Adv Appl Microbiol* 52:167–186. [https://doi.org/10.1016/S0065-2164\(03\)01007-4](https://doi.org/10.1016/S0065-2164(03)01007-4).
51. Lund P, Tramonti A, Biase DD. 2014. Coping with low pH: molecular strategies in neutralophilic bacteria. *FEMS Microbiol Rev* 38:1091–1125. <https://doi.org/10.1111/1574-6976.12076>.
52. Zambrano MM, Siegle DA, Almirón M, Tormo A, Kolter R. 1993. Microbial competition: *Escherichia coli* mutants that take over stationary phase cultures. *Science* 259:1757–1760. <https://doi.org/10.1126/science.7681219>.
53. Spira B, de Almeida Toledo R, Maharjan RP, Ferenci T. 2011. The uncertain consequences of transferring bacterial strains between laboratories: *rpoS* instability as an example. *BMC Microbiol* 11:248. <https://doi.org/10.1186/1471-2180-11-248>.
54. Hommais F, Krin E, Coppée J-Y, Lacroix C, Yeramian E, Danchin A, Bertin P. 2004. GadE (YhiE): a novel activator involved in the response to acid environment in *Escherichia coli*. *Microbiology (Reading)* 150:61–72. <https://doi.org/10.1099/mic.0.26659-0>.
55. Tramonti A, De Canio M, Delany I, Scarlato V, De Biase D. 2006. Mechanisms of transcription activation exerted by GadX and GadW at the *gadA* and *gadBC* gene promoters of the glutamate-based acid resistance system in *Escherichia coli*. *J Bacteriol* 188:8118–8127. <https://doi.org/10.1128/JB.01044-06>.
56. Seo SW, Kim D, O'Brien EJ, Szubin R, Palsson BO. 2015. Decoding genome-wide GadEWX-transcriptional regulatory networks reveals multifaceted cellular responses to acid stress in *Escherichia coli*. *Nat Commun* 6:7970. <https://doi.org/10.1038/ncomms8970>.
57. Travers A, Muskhelishvili G. 2005. DNA supercoiling: a global transcriptional regulator for enterobacterial growth? *Nat Rev Microbiol* 3:157–169. <https://doi.org/10.1038/nrmicro1088>.
58. Dorman CJ. 2006. DNA supercoiling and bacterial gene expression. *Sci Prog* 89:151–166. <https://doi.org/10.3184/003685006783238317>.
59. Conter A, Menchon C, Gutierrez C. 1997. Role of DNA supercoiling and RpoS sigma factor in the osmotic and growth phase-dependent induction of the gene *osmE* of *Escherichia coli* K12. *J Mol Biol* 273:75–83. <https://doi.org/10.1006/jmbi.1997.1308>.
60. Bordes P, Conter A, Morales V, Bouvier J, Kolb A, Gutierrez C. 2003. DNA supercoiling contributes to disconnect σ^S accumulation from σ^S -dependent transcription in *Escherichia coli*. *Mol Microbiol* 48:561–571. <https://doi.org/10.1046/j.1365-2958.2003.03461.x>.
61. Cameron ADS, Dorman CJ. 2012. A fundamental regulatory mechanism operating through OmpR and DNA topology controls expression of *Salmonella* pathogenicity islands SPI-1 and SPI-2. *PLoS Genet* 8:e1002615. <https://doi.org/10.1371/journal.pgen.1002615>.
62. Dorman CJ. 2013. Genome architecture and global gene regulation in bacteria: making progress towards a unified model? *Nat Rev Microbiol* 11:349–355. <https://doi.org/10.1038/nrmicro3007>.
63. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol* 2:2006.0008.
64. Haldimann A, Wanner BL. 2001. Conditional-replication, integration, excision, and retrieval plasmid-host systems for gene structure-function studies of bacteria. *J Bacteriol* 183:6384–6393. <https://doi.org/10.1128/JB.183.21.6384-6393.2001>.
65. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* 97:6640–6645. <https://doi.org/10.1073/pnas.120163297>.
66. Groth D, Reszka R, Schenk JA. 1996. Polyethylene glycol-mediated transformation of *Escherichia coli* is increased by room temperature incubation. *Anal Biochem* 240:302–304. <https://doi.org/10.1006/abio.1996.0362>.
67. Butland G, Peregrín-Alvarez JM, Li J, Yang W, Yang X, Canadien V, Starostine A, Richards D, Beattie B, Krogan N, Davey M, Parkinson J, Greenblatt J, Emili A. 2005. Interaction network containing conserved and essential protein complexes in *Escherichia coli*. *Nature* 433:531–537. <https://doi.org/10.1038/nature03239>.
68. Fraley CD, Kim JH, McCann MP, Martin A. 1998. The *Escherichia coli* starvation gene *cstC* is involved in amino acid catabolism. *J Bacteriol* 180:4287–4290.
69. Castanie-Cornet M-P, Foster JW. 2001. *Escherichia coli* acid resistance: cAMP receptor protein and a 20 bp *cis*-acting sequence control pH and stationary phase expression of the *gadA* and *gadBC* glutamate decarboxylase genes. *Microbiology* 147:709–715. <https://doi.org/10.1099/002221287-147-3-709>.
70. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 6:343–345. <https://doi.org/10.1038/nmeth.1318>.
71. De Biase D, Tramonti A, Bossa F, Visca P. 1999. The response to stationary-phase stress conditions in *Escherichia coli*: role and regulation of the glutamic acid decarboxylase system. *Mol Microbiol* 32:1198–1211. <https://doi.org/10.1046/j.1365-2958.1999.01430.x>.
72. Farewell A, Kvint K, Nyström T. 1998. Negative regulation by RpoS: a case of sigma factor competition. *Mol Microbiol* 29:1039–1051. <https://doi.org/10.1046/j.1365-2958.1998.00990.x>.
73. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
74. Anders S, Pyl PT, Huber W. 2015. HTSeq: a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166–169. <https://doi.org/10.1093/bioinformatics/btu638>.
75. Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol* 11:R106. <https://doi.org/10.1186/gb-2010-11-10-r106>.
76. Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B Methodol* 57:289–300.
77. Alexa A, Rahnenfuhrer J. 2010. topGO: enrichment analysis for gene ontology. R package version 2.22.0. R Project for Statistical Computing, Vienna, Austria.
78. Carlson M. 2014. Genome wide annotation for *E. coli* strain K12. R package version 3.0.0, org.EcK12.eg.db. R Project for Statistical Computing, Vienna, Austria.
79. Micallef L, Rodgers P. 2014. eulerAPE: drawing area-proportional 3-venn diagrams using ellipses. *PLoS One* 9:e101717. <https://doi.org/10.1371/journal.pone.0101717>.
80. Singh SS, Singh N, Bonocora RP, Fitzgerald DM, Wade JT, Grainger DC. 2014. Widespread suppression of intragenic transcription initiation by H-NS. *Genes Dev* 28:214–219. <https://doi.org/10.1101/gad.234336.113>.
81. Machanick P, Bailey TL. 2011. MEME-ChIP: motif analysis of large DNA datasets. *Bioinformatics* 27:1696–1697. <https://doi.org/10.1093/bioinformatics/btr189>.
82. Bailey TL, Elkan C. 1994. Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proc Int Conf Intell Syst Mol Biol* 2:28–36.
83. Vandesompele J, Preter KD, Pattyn F, Poppe B, Roy NV, Paepe AD, Speleman F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3:research0034.
84. Miller JH. 1992. A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
85. Holm S. 1979. A simple sequentially rejective multiple test procedure. *Scand J Stat* 6:65–70.
86. Schneider TD, Stephens RM. 1990. Sequence logos: a new way to display consensus sequences. *Nucleic Acids Res* 18:6097–6100. <https://doi.org/10.1093/nar/18.20.6097>.
87. Crooks GE, Hon G, Chandonia J-M, Brenner SE. 2004. WebLogo: a sequence logo generator. *Genome Res* 14:1188–1190. <https://doi.org/10.1101/gr.849004>.