

1 UNIVERSITY OF CHINESE ACADEMY OF SCIENCES

2 DOCTORAL THESIS

3

4 **The measurement of Higgs**
5 **production cross section in the**
6 **four-lepton final state at CMS**

7

8 *Author:*

Tahir JAVAID

Supervisor:

Prof. Guoming CHEN

Co-supervisor:

Prof. Mingshui CHEN

9 *A thesis submitted in fulfillment of the requirements*
10 *for the degree of Doctor of Science*

11 *in the*

12 **Institute of High Energy Physics**
13 **Chinese Academy of Sciences**



Contents

16	List of Figures	v
17	List of Tables	ix
18	1 Physics Objects studies with the CMS detector	1
19	1.1 Physics Objects	1
20	1.1.1 Reconstruction of Track and Vertex	1
21	1.1.2 Electrons	3
22	Electron Reconstruction	3
23	Electron Identification and Isolation	3
24	Electron Impact Parameter Selection	4
25	Electron Energy Calibrations	7
26	Electron Efficiency Measurements	7
27	1.1.3 Muons	11
28	Muon Reconstruction and Identification	11
29	Muon Isolation	12
30	Muon Impact Parameter Selection	12
31	Muon Energy Calibrations	12
32	Efficiency Measurements of muons	12
33	1.1.4 Photons for FSR recovery	18
34	1.1.5 Jets	19
35	Jet Identification	19
36	Jet Energy Corrections	19
37	B-tagging	19
38	Validation on data	20
39	1.1.6 Object Summary	22

List of Figures

41	1.1	Reconstruction efficiencies of electron in data versus p_T (left), η (right)	
42		for electrons having $p_T > 20$ GeV (top) and $p_T < 20$ GeV (bottom)	
43		with corresponding data/MC scale factors as provided by the EGM	
44		POG.	4
45	1.2	Output of the multiclassifier discriminant for prompt electrons matched	
46		to truth electrons from Z decay (blue) and for fakes (red). Events are	
47		all taken from DY events MC sample.	5
48	1.3	The receiver operating characteristic curves (background efficiency	
49		vs signal efficiency) of the MVA's trained on the 2017 simulation and	
50		applied on the 2018 simulation. The training combines identification	
51		and isolation variables. Performance are shown for electrons with	
52		$5 < p_T < 10$ GeV (left), $p_T > 10$ GeV (right) and $ \eta < 0.8$ (top),	
53		$0.8 < \eta < 1.479$ (middle) and $ \eta > 1.479$ (bottom). V1 and V2	
54		versions of training are compared, exploiting TMVA and xgboost	
55		training libraries respectively.	6
56	1.4	(a) electron energy scale measured in the $Z \rightarrow ee$ control region for	
57		all electrons, (b) for both electrons in barrel, (c) for one electron in	
58		barrel and other in endcaps, and for both of the electrons in endcaps	
59		(d). In these figures, the results based on fits using Crystall-ball are	
60		shown.	8
61	1.5	Electron selection efficiencies vs p_T measured using the Tag-and-	
62		Probe technique, non-gap electrons (left) and gap electrons (right),	
63		together with the corresponding data/MC ratio bottom part of each	
64		plot.	9
65	1.6	Electron selection efficiencies vs η , non-gap electrons (left) and gap	
66		electrons (right), together with the corresponding data/MC ratio at	
67		the bottom part of each plot.	10

68	1.7	(a): muon energy scale measured in the $Z \rightarrow \mu\mu$ control region for	
69		all muons, for both of muons from barrel region (b), for one muon	
70		from barrel, one from endcaps (c) and for both muons in the end-	
71		caps (d). The results in the figures are extracted from fits using the	
72		Crystall-ball function.	13
73	1.8	Reconstruction and identification efficiencies of muons at low p_T ,	
74		measured with the $T\&P$ method on events, in bins of p_T in barrel	
75		(left) and endcaps (center), and in terms of η for $p_T > 7$ GeV (right).	
76		In upper panel of each diagram, the larger uncertainty bars include	
77		also the systematical uncertainties, while the smaller ones are purely	
78		statistical. In the lower panel showing the ratio of two efficiencies,	
79		black uncertainty bars are for statistical uncertainty, the orange rect-	
80		angles for the systematical uncertainty and the violet rectangles in-	
81		clude both uncertainties.	15
82	1.9	Efficiency of the muon impact parameter requirements, measured	
83		with the $T\&P$ method on Z events, in bins of p_T in barrel (left) and	
84		endcaps (center), and in terms of η for all p_T (right). In upper panel,	
85		the larger uncertainty bars include also the systematical uncertain-	
86		ties, while the smaller ones are purely statistical. In the lower panel	
87		showing the ratio of two efficiencies, black error bars are for statis-	
88		tical uncertainty, the orange rectangles for the systematical uncer-	
89		tainty and the violet rectangles include both uncertainties.	16
90	1.10	Efficiency of muon isolation requirement, measured with the tag&probe	
91		method on Z events, as function of p_T in the barrel (left) and endcaps	
92		(right). In the upper panel, the larger error bars include also the sys-	
93		tematical uncertainties, while the smaller ones are purely statistical.	
94		In the lower panel showing the ratio of the two efficiencies, the black	
95		error bars are for the statistical uncertainty, the orange rectangles for	
96		the systematical uncertainty and the violet rectangles include both	
97		uncertainties.	17
98	1.11	Left: Overall scale factors for muons, as function of p_T and η . Right:	
99		Uncertainties on overall scale factors muons shown in two dimen-	
100		sions of p_T and η	17

101	1.12 Comparison of data with MC for leading jet η in $Z \rightarrow \ell\ell + \text{jets}$. MC	
102	is normalized to data. Jet ID and Jet PUID are applied. MC samples	
103	include DY and $t\bar{t}$. Data/MC ratio plots are shown in the bottom	
104	of each plots, together with the uncertainties (shaded histograms)	
105	from Jet Energy Corrections.	21

List of Tables

107	1.1	Overview of variables used at inputs to identification classifier. The	
108		Variables not used in the run I MVA are specified with ($\cdot$).	5
109	1.2	Minimum value of BDT score for electron to pass identification, to-	
110		gether with the corresponding signal and the background efficiencies.	7
111	1.3	The requirements for a muon to pass the Tracker High- p_T ID. Note	
112		that these are equivalent to the Muon POG High- p_T ID with the	
113		global track requirements removed.	11
114	1.4	Chart of selection of physics objects for the $H \rightarrow ZZ \rightarrow 4\ell$ analysis	
115		used on the 2018 data.	23

Chapter 1

Physics Objects studies with the CMS detector

1.1 Physics Objects

As described in section ??, that events organized in streams of HLT decisions are eventually transferred to the storage system (Tier-0) of CMS for its permanent storage. That data is in RAW format and needs to be fully reconstructed from this storage system. With potential reconstruction procedure, this RAW data can be associated to physical particles such as simple or general tracks, muons, photons, electrons and jets etc. These reconstructed particles then help to explore *e.g.* Higgs decay to four lepton channel for its properties measurement like spin, parity, cross sections, mass and etc. The reconstruction of the SM Higgs boson in the decay chain $H \rightarrow ZZ \rightarrow 4\ell$ requires very efficient lepton reconstruction and identification in order to be sensitive to a low mass Higgs. In this kinematics region, where at least p_T of one of the leptons lies within the range $5 - 15 \text{ GeV}$, there is need to attain an optimal reconstruction efficiency keeping misidentified lepton rate as low as possible. Further, the precise measurements on Higgs are lepton kinematics dependent and thus needs high precision momentum measurement.

1.1.1 Reconstruction of Track and Vertex

Reconstructing a track is procedure of exploiting hits in the detector parts used to estimate the momentum and/or position of a particle producing the hits. It is a challenging task and in the CMS experiment there a dedicated software known as Combinatorial Track Finder (CTF). It uses combinatorial Kalman filter approach for the recognition of patterns and fitting the track.

CTF algorithm repeats in multiple steps to search the track on the basis of its relatively larger p_T and closeness to the interaction point [Chatrchyan:2014fea].

This step is called iterative tracking in which each iteration is done in following steps:

- **Seeding:** It is to provide with the help of 2 or 3 hits initial track candidates (seeds).
- **Tracking Finding:** It is to search for the additional hits based on the seed trajectories based on the Kalman filter approach.
- **Estimate of trajectory:** It is to use the Kalman-filter and smoother to estimate the best parameters of each trajectory.
- **Track Selection:** Based on the quality of the trajectories, those tracks which do not originate from some charged particle are discarded.

It is repeated for 6 times which are done to figure out the tracks originating from outside of the luminous region of the proton proton collisions.

The completely reconstructed tracks give the information of the proton-proton interaction vertices. Reconstruction of the primary-vertex is done in three steps:

- **Selection of Tracks:** It is done by setting requirements on Significance of impact parameter (SIP) from beam spot, number of hits in ITS and normalized χ^2
- **Track Clustering:** It is done using information of transverse component of the impact parameter with respect to the beam spot. The clustering algorithm is capable to reconstruct any number of proton proton interaction in the bunch crossing. This performed using deterministic annealing algorithm (DA). This algorithm clusters the tracks into groups which can be associated to a single vertex.
- **Fitting of the track:** It is performed over the tracks in each cluster using "adaptive vertex filter" to get the position of the primary vertex (the best vertex parameters). It needs a procedure to link a primary vertex with the hard interaction. Since in hard interaction particles are produced at high p_T , the vertices can be associated with sum of p_T^2 of the reconstructed tracks. The

vertex with highest sum of p_T^2 can be associated to the hard interaction from the multiplicity of primary vertices.

1.1.2 Electrons

Electron Reconstruction

Preselection of the electron candidates is made through loose cuts on track-cluster matching observables. This done to maintain highest possible selection efficiency of electrons and to reject QCD background significantly. For the analysis, electrons are subjected to have $p_T^e > 7$ GeV within $|\eta^e| < 2.5$, and satisfy a loose primary vertex constraints on longitudinal and transvers components of impact parameter *viz.* $d_{xy} < 0.5$ cm and $d_z < 1$ cm respectively. Such electrons are called **loose electrons**. Details of electron reconstruction can be explored in Ref. [ElectronLegacy].

The data-MC discrepancy is corrected using scale factors as is done for the electron selection with data efficiencies measured through the same tag-and-probe technique outlined later (see Section 1.1.2). These studies for reconstructions are carried out by the EGM POG and the results are only summarised here.

The electron reconstruction scale factors are shown Fig. 1.1 and are applied as a function of electron p_T and super cluster η .

Electron Identification and Isolation

With respect to previous analyses, an important improvement brought in this analysis is usage of a new multivariate discriminant for electron selection in all data taking periods.

Identification and isolation of the reconstructed electrons is now made using Gradient Boosted Decision Tree (GBDT) which is a multivariate classifier algorithm. It exploits inputs from electromagnetic cluster, cluster-track matching, variables which exclusively depend on tracking measurements and as well as particle flow isolation sums. The list of observables used can be seen in the Table 1.1.

For both signal and the background, the classifier is trained on 2017 MC samples of Drell-Yan(DY) plus jets. Fig. 1.2 shows the output of the multiclassifier discriminant for prompt electrons from DY events and fakes from jets in DY events.




FIGURE 1.1: Reconstruction efficiencies of electron in data versus p_T (left), η (right) for electrons having $p_T > 20$ GeV (top) and $p_T < 20$ GeV (bottom) with corresponding data/MC scale factors as provided by the EGM POG.

The impact of the transition from the TMVA(v1) to the xgboost(v2) training library is shown in Fig. 1.3. The working points shown are adopted to get the same signal efficiency as the cut on MVA ID and a cut on the PF isolation, in each p_T and η bin.

Table 1.2 lists the cut values applied to the BDT score for the chosen working point. For the analysis, loose electrons have to pass this MVA identification and isolation working point.

Electron Impact Parameter Selection

For lepton to be consistent with some common primary vertex, we require that they have an associated track that corresponds a small impact parameter from the

observable type	observable name
6*cluster shape	RMS value of spectrum of energy-crystal number along η and φ ; $\sigma_{i\eta i\eta}$, $\sigma_{i\varphi i\varphi}$ width of super cluster along η and ϕ ratio of hadronic energy behind the supercluster of electron to supercluster energy, H/E circularity $(E_{5\times 5} - E_{5\times 1})/E_{5\times 5}$ sum of the seed and adjacent crystal over the super cluster energy R_9 for the endcap training bins: pre-shower energy fraction E_{PS}/E_{raw}
2*track-cluster matching	energy-momentum agreement E_{tot}/p_{in} , E_{ele}/p_{out} , $1/E_{tot} - 1/p_{in}$ position matching $\Delta\eta_{in}$, $\Delta\varphi_{in}$, $\Delta\eta_{seed}$
5*tracking	fractional loss of momentum loss $f_{brem} = 1 - p_{out}/p_{in}$ number of hits of KF and the GSF track N_{KF} , N_{GSF} ($\cdot$) reduced χ^2 of KF and the GSF track χ^2_{KF} , χ^2_{GSF} number of expected innre hits (missing) ($\cdot$) probability transformation in conversion vertex fit χ^2 ($\cdot$)
3*isolation	Particle Flow photon isolation sum ($\cdot$) Particle Flow charged hadrons isolation sum ($\cdot$) Particle Flow neutral hadrons isolation sum ($\cdot$)
1*For PU-resilience	Mean energy density in the event: ρ ($\cdot$)

TABLE 1.1: Overview of variables used at inputs to identification classifier. The Variables not used in the run I MVA are specified with ($\cdot$).




FIGURE 1.2: Output of the multiclassifier discriminant for prompt electrons matched to truth electrons from Z decay (blue) and for fakes (red). Events are all taken from DY events MC sample.




FIGURE 1.3: The receiver operating characteristic curves (background efficiency vs signal efficiency) of the MVA's trained on the 2017 simulation and applied on the 2018 simulation. The training combines identification and isolation variables. Performance are shown for electrons with $5 < p_T < 10 \text{ GeV}$ (left), $p_T > 10 \text{ GeV}$ (right) and $|\eta| < 0.8$ (top), $0.8 < |\eta| < 1.479$ (middle) and $|\eta| > 1.479$ (bottom). V1 and V2 versions of training are compared, exploiting TMVA and xgboost training libraries respectively.

211 event primary vertex. We use the significance of impact parameter to the vertex
 212 of an event, i.e. $|\text{SIP}_{3D}| = \frac{|\text{IP}|}{\sigma_{\text{IP}}}$, where IP being three dimensional lepton impact
 213 parameter w.r.t the primary interaction vertex, and the σ_{IP} being the uncertainty

	$ \eta < 0.8$		
	Cut on BDT score	Signal eff.	Background eff.
$5 < p_T < 10 \text{ GeV}$	1.264	81.04%	4.4%
$p_T > 10 \text{ GeV}$	2.364	97.1%	2.9%
	$0.8 < \eta < 1.479$		
	Cut on BDT score	Signal eff.	Background eff.
$5 < p_T < 10 \text{ GeV}$	1.178	79.3%	4.6%
$p_T > 10 \text{ GeV}$	2.078	96.3%	3.6%
	$ \eta > 1.479$		
	Cut on BDT score	Signal eff.	Background eff.
$5 < p_T < 10 \text{ GeV}$	1.330	72.97%	3.6%
$p_T > 10 \text{ GeV}$	1.080	95.7%	6.7%

TABLE 1.2: Minimum value of BDT score for electron to pass identification, together with the corresponding signal and the background efficiencies.

associated to this. Hence, a lepton satisfying $|\text{SIP}_{3D}| < 4$ is regarded as "primary lepton".

Electron Energy Calibrations

Electrons in data are corrected for features in ECAL energy scale in bins of p_T and $|\eta|$. Corrections are calculated on a $Z \rightarrow ee$ sample to align the dielectron mass spectrum in data to that in MC, and to minimize its width.

The resolution of $Z \rightarrow ee$ mass in the Monte Carlo is made to match with data by applying a pseudorandom Gaussian smearing to electron energies, with varying parameters of Gaussian in p_T and $|\eta|$ bins. This has effect of convoluting the electron energy spectrum with a Gaussian.

The electron energy scale is measured in data by fitting a Crystall-ball function to di-electron spectrum of mass around the Z peak in the $Z \rightarrow ee$ control region. The energy scale for the full 2018 dataset is shown in Fig. 1.4(a) and agrees decently with the MC with the preliminary corrections released by EGAMMA POG.

Electron Efficiency Measurements

The Tag-and-Probe study was performed on the EGM dataset using the golden JSON of 58.83, 41.4 and 35.9 fb⁻¹ respectively for 2018, 2017 and 2016. More details on this technique can be found in Ref. [AN-15-277].

Tag electrons need to satisfy the following quality requirements:

- trigger matched to HLT_Ele32_WPTight_Gsf_L1DoubleEG_v*
- $p_T > 30 \text{ GeV}$, super cluster (SC) $\eta < 2.17$




FIGURE 1.4: (a) electron energy scale measured in the $Z \rightarrow ee$ control region for all electrons, (b) for both electrons in barrel, (c) for one electron in barrel and other in endcaps, and for both of the electrons in endcaps (d). In these figures, the results based on fits using Crystall-ball are shown.

235 For the bin between 7 and 20 GeV, additional criteria are required: the tag has
 236 to pass a cut on the MVA ID > 0.92 , $\sqrt{(2 * PFMET * p_T(tag) * (1 - \cos(\phi_{MET} -$
 237 $\phi_{tag}))) < 45 \text{ GeV}$.

238 Probe electrons only need to be reconstructed as GsfElectron while the FSR
 239 recovery algorithm is not applied in efficiency measurement.

The nominal MC efficiencies are evaluated from the LO MadGraph Drell-Yan, while the NLO systematics use the MadGraph_AMCatNLO sample.

For the efficiency measurements a template fit is used. The m_{ee} signal shape of the passing and failing probes is taken from MC and convoluted with a Gaussian. The data is then fitted with the convoluted MC template and a CMSShape (an Error-function with a one-sided exponential tail). This change follows from the usage of the T&P tool developed by the EGM POG.

The measurement of electron selection efficiency is made as a function of the probe electron p_T and its SC η . For electrons falling in the ECAL gaps, it is measured separately. Figure 1.5 and ?? shows the p_T and η turn-on curves measured in data.




FIGURE 1.5: Electron selection efficiencies vs p_T measured using the Tag-and-Probe technique, non-gap electrons (left) and gap electrons (right), together with the corresponding data/MC ratio bottom part of each plot.

The EGM recommendations about evaluation of Tag-and-Probe uncertainties on efficiency measurements are followed. Specifically, we consider

- Variation of the signal shape from a MC shape to an analytic shape (Crystal Ball) fitted to the MC
- Use of analytic function (Crystal Ball) to fit the signal as alternative of MC template




FIGURE 1.6: Electron selection efficiencies vs η , non-gap electrons (left) and gap electrons (right), together with the corresponding data/MC ratio at the bottom part of each plot.

- Using an NLO MC sample for signal templates

Assessment of total uncertainty for measurement of the scale factors is taken as quadratic sum of statistical uncertainties (returned from the fit) and the aforementioned systematic uncertainties.

1.1.3 Muons

Muon Reconstruction and Identification

We define **loose muons** as the muons that satisfy $p_T > 5$, $|\eta| < 2.4$, $d_{xy} < 0.5$ cm, $d_z < 1$ cm, where d_{xy} and d_z are defined w.r.t. the PV and using the 'muonBestTrack'. Muons have to be reconstructed by either the Global Muon or Tracker Muon algorithm. Standalone Tracks of muon reconstructed only in the muon system are rejected. Muons with `muonBestTrackType==2` (standalone) are discarded even if they are marked as global or tracker muons. More details on the reconstruction of muons can be seen in Ref. [AN-15-277].

Loose muons with p_T below 200 GeV are considered identified muons if they also pass the PF muon ID (note that the naming convention used for these IDs differs from the muon POG naming scheme, in which the "tight ID" used here is called the "loose ID"). Loose muons with p_T above 200 GeV are considered identified muons if they pass the PF ID or the Tracker High- p_T ID, the definition of which is shown in Table 1.3. This relaxed definition is used to increase signal efficiency for the high-mass search. When a very heavy resonance decays to two bosons, both bosons will be very boosted. In the lab frame, the leptons originating from decay of a highly boosted Z will be nearly collinear, and the PF ID loses efficiency for muons separated by approximately $\Delta R < 0.4$, which roughly corresponds to muons originating from bosons with $p_T > 500$ GeV.

TABLE 1.3: The requirements for a muon to pass the Tracker High- p_T ID. Note that these are equivalent to the Muon POG High- p_T ID with the global track requirements removed.

Plain-text description	Technical description
Muon station matching	Matching of muon to the segments in two muon stations at least NB: It means that muon is an arbitrated tracker muon.
Good p_T measurement	$\frac{p_T}{\sigma_{p_T}} < 0.3$
Vertex compatibility ($x - y$)	$d_{xy} < 2$ mm
Vertex compatibility (z)	$d_z < 5$ mm
Pixel hits	One pixel hit at least
Tracker hits	Hits in at least six tracker layers

It may happen that a single muon is reconstructed incorrectly as two muons or even more. To cope with this situation, additional "ghost-cleaning" is performed

as:

- Those tracker Muons which are not Global Muons, are needed to be arbitrated.
- If two muons are sharing their segments upto 50% or more, lower quality muon is removed.

Muon Isolation

Muon isolation used is based on Particle-Flow. So-called $\Delta\beta$ correction is used in order to subtract the pileup contribution for muons, whereby $\Delta\beta = \frac{1}{2} \sum_{\text{PU}}^{\text{charged had.}} p_T$ gives an estimate of the energy deposit of neutral particles (i.e. photons and hadrons) from the pile-up vertices. Relative isolation for muons is then defined as:

$$\text{RelPFiso} = \frac{\sum^{\text{charged had.}} p_T + \max(\sum^{\text{neutral had.}} + \sum^{\text{photon}} - \Delta\beta, 0)}{p_T^{\text{lepton}}} \quad (1.1)$$

The isolation working point for muons was optimized in Ref. [AN-15-277] and the working point was chosen to be the same as electrons, namely $\text{RelPFiso}(\Delta R = 0.3) < 0.35$.

Muon Impact Parameter Selection

Selection on muon significance of impact parameter(SIP) is used same as for the electrons which is described in Section 1.1.2:

- $|\text{SIP}_{3\text{D}} = \frac{\text{IP}}{\sigma_{\text{IP}}}| < 4$

Muon Energy Calibrations

Similar to electrons the momentum scale of muon is measured in data by fitting a Crystall-ball function to the di-muon invariant mass spectrum around the Z peak in the $Z \rightarrow \mu\mu$ control region. From Fig. 1.7 shows a very good comparison of data with simulation.

Efficiency Measurements of muons

Efficiencies of muons are measured using same method used for electron measurements i.e. Tag-and-Probe (T&P) method performed on $Z \rightarrow \mu\mu$ and $\rightarrow \mu\mu$




FIGURE 1.7: (a): muon energy scale measured in the $Z \rightarrow \mu\mu$ control region for all muons, for both of muons from barrel region (b), for one muon from barrel, one from endcaps (c) and for both muons in the endcaps (d). The results in the figures are extracted from fits using the Crystall-ball function.

308 events in bins of p_T and η . More details on the methodology can be found in Ref.
 309 [CMS_AN_2015-277]. Measurements are extracted using 2018 RunA,B,C,D data
 310 while the measurements corresponding to 2016 and 2017 datasets have already
 311 been reported in Ref. [CMS_AN_2016-442] and Ref. [CMS_AN_2017-342] respec-
 312 tively. For the measurement of muon reconstruction and identification efficiency at

high p_T ($p_T > 20 \text{ GeV}$), and impact parameter requirements and isolation efficiencies for all p_T , Z sample is used. to measure the muon reconstruction and identification efficiency at high p_T , and the efficiency of the isolation and impact parameter requirements at all p_T . To measure the reconstruction and identification efficiency at low p_T , sample is used as it benefits from a better purity in that kinematic regime. In this case, events are collected using `HLT_Mu7p5_Track2_Jpsi_v*` when probing the reconstruction and identification efficiency in the muon system, and using the `HLT_Mu7p5_L2Mu2_Jpsi_v*` when probing the tracking efficiency.

Reconstruction and identification Results for muon reconstruction and identification efficiencies for high p_T muons ($p_T > 20 \text{ GeV}$) have been derived by the Muon POG. The probe in this measurement are tracks of inner tracker, and the passing probes are those which are also regarded as a global or tracker muon and passing the Muon POG Loose muon identification. Results for low p_T muons were derived using events, with the same definitions of probe and passing probes. The systematic uncertainties on the measurements are estimated by including several variations e.g. varying the analytical signal and background shape models used to fit distribution of dimuon invariant mass. Furthermore, number of bins in dimuon distribution and mass window are also increased and decreased to include systematic effects. Details on the procedure can be found in Ref. [AN-15-277]. The efficiency and scale factors used for low p_T muons are the ones derived using single muon prompt-reco dataset.

The data and simulation efficiencies and their ratios in bins are shown in Fig. 1.8.

Impact parameter requirements The measurement is performed using Z events. Events are selected with `HLT_IsoMu27_v*` or `HLT_Mu50_v*` triggers. For this measurement, the probe is a muon passing the POG Loose identification criteria, and it is considered a passing probe if satisfies the SIP3D, dxy, dz cuts of this analysis. The results are shown in the Fig. 1.9.

Isolation requirements Isolation efficiency is measured using events from the Z decay for any p_T . The events are selected with either of `HLT_IsoMu27_v*` or `HLT_Mu50_v*` triggers. The isolation of the muons are calculated after recovery




FIGURE 1.8: Reconstruction and identification efficiencies of muons at low p_T , measured with the $T\&P$ method on events, in bins of p_T in barrel (left) and endcaps (center), and in terms of η for $p_T > 7$ GeV (right). In upper panel of each diagram, the larger uncertainty bars include also the systematical uncertainties, while the smaller ones are purely statistical. In the lower panel showing the ratio of two efficiencies, black uncertainty bars are for statistical uncertainty, the orange rectangles for the systematical uncertainty and the violet rectangles include both uncertainties.




FIGURE 1.9: Efficiency of the muon impact parameter requirements, measured with the $T\&P$ method on Z events, in bins of p_T in barrel (left) and endcaps (center), and in terms of η for all p_T (right). In upper panel, the larger uncertainty bars include also the systematical uncertainties, while the smaller ones are purely statistical. In the lower panel showing the ratio of two efficiencies, black error bars are for statistical uncertainty, the orange rectangles for the systematical uncertainty and the violet rectangles include both uncertainties.

of the FSR photons and subtracting their contribution to the isolation cone of the muons. There are details described of this method which can be found in Ref. [AN-16-217].

The results are shown in Fig. 1.10.

Tracking Tracking efficiency (i.e. efficiency to associate a track to some muon in inner detector) is almost 100% and hence muon Physics Object Group no more recommends tracking scale factors in the analyses.

Overall results of muon Scale Factors The product of all scale factors for reconstruction, identification, impact parameter and isolation requirements is shown in two dimensional display in Fig. 1.11. The figure also shows overall uncertainty on scale factors.




FIGURE 1.10: Efficiency of muon isolation requirement, measured with the tag&probe method on Zevents, as function of p_T in the barrel (left) and endcaps (right). In the upper panel, the larger error bars include also the systematical uncertainties, while the smaller ones are purely statistical. In the lower panel showing the ratio of the two efficiencies, the black error bars are for the statistical uncertainty, the orange rectangles for the systematical uncertainty and the violet rectangles include both uncertainties.




FIGURE 1.11: Left: Overall scale factors for muons, as function of p_T and η . Right: Uncertainties on overall scale factors muons shown in two dimensions of p_T and η .

1.1.4 Photons for FSR recovery

The FSR recovery algorithm was considerably simplified with respect to what was done in Run I, while maintaining similar performance. The selection of FSR photons is now only done per-lepton and no longer depends on any Z mass criterion, thus much simplifying the subsequent ZZ candidate building and selection. As regards the association of photons with leptons, the rectangular cuts on $\Delta R(\gamma, l)$ and $E_{T,\gamma}$ have been replaced by a cut on $\Delta R(\gamma, l)/E_{T,\gamma}^2$.

Initially, photons are collected using particle-flow algorithm. Further selection and their association with a lepton is done as follows:

1. PF photons are preselected by requiring $p_{T,\gamma} > 2 \text{ GeV}$, $|\eta^\gamma| < 2.4$, and a relative Particle-flow isolation smaller than 1.8 (computed within a cone of radius $R = 0.3$). On charged hadrons and neutral hadrons or photons 0.2 GeV and 0.5 GeV thresholds are applied with a size of veto cone 0.0001 and 0.01 respectively, also considering contribution from pileup vertices (with the same radius and threshold as per charged isolation) .
2. Supercluster (SC) veto: All photons matching (direct association with PF candidates) with some electron that passes ID and SIP requirements are removed.
3. Association of photons is made with lepton which is closest and passes ID and SIP requirements.
4. Photons not satisfying the requirements $\Delta R(\gamma, l)/E_{T,\gamma}^2 < 0.012$, and $\Delta R(\gamma, l) < 0.5$ are discarded.
5. In case if a photon gets associated with same lepton, the one with less $\Delta R(\gamma, l)/E_{T,\gamma}^2$ is opted.
6. For each selected FSR photon, we exclude that photon from sum of isolation of all leptons in an event which pass both loose ID and SIP requirements. This concerns the photons which are in isolation cone and outside veto of isolation of said leptons ($\Delta R < 0.4$ and $\Delta R > 0.01$ for muons and $\Delta R < 0.4$ and $(\eta^{\text{SC}} < 1.479$ or $\Delta R > 0.08)$ for electrons).

More details on the optimization of the FSR photon selection can be found in Ref. [AN-15-277, AN-16-217].

1.1.5 Jets

Higgs production through fusion of vector bosons (VBF) and through other mechanisms normally differ as regards the jet kinematics.

Jet Identification

Jets are reconstructed through anti- k_T clustering algorithm out of particle-flow candidates, with distance parameter $R = 0.4$, after rejecting the charged hadrons that are associated with a pileup primary vertex.

To reduce instrumental background, the tight working point jet ID suggested by the JetMET Physics Object Group is applied [JetID2018]. In addition, jets from Pile-Up are rejected using the PileUp jet ID criteria suggested by the JetMET POG [JetPUID2017]. In this analysis, the jets from within $|\eta| < 4.7$ area are required to have a transverse momentum greater than 30 GeV. In addition, jets are cleaned from any of the tight leptons (passing the SIP and isolation cut computed after FSR correction) and FSR photons by a separation criterion: $\Delta R(\text{jet}, \text{lepton}/\text{photon}) > 0.4$.

Jet Energy Corrections

The capability of current calorimeter to measure the energy of jets correctly is not linear. Hence the measured energy cannot be regarded as particle's true energy and we need to estimate corrections to Jet Energy. In the analysis, standard jet energy corrections are implemented to the reconstructed jets, which consist of L1 Pileup, L2 Relative Jet Correction, L3 Absolute Jet Correction for both Monte Carlo samples and data, and also residual calibration for data [JECMC2018]. Jet Energy resolution corrections are also applied to MC.

B-tagging

In order to distinguish whether a jet is b-jet or not *DeepCSV* algorithm is used as our b-tagging algorithm. It combines the same information as the previous tagger *CSVv2* i.e. SIP, jet kinematics and secondary vertex but uses information of more tracks. Also, the b-tag output discriminator is computed with a Deep Neural Network. In the analysis, a jet is considered as a b-tagged if it meets requirement

of *medium* working point, i.e. if its `|pfDeepCSVJetTags:probb+ pfDeepCSVJetTags:probbb|` discriminator is greater than 0.4184 [BTAG2018].

Scale factors of data to simulation efficiencies for b-tagging are provided for this working point for the full dataset varied in jet p_T , η and flavour. They are applied to simulated jets by downgrading (upgrading) the b-tagging status of a fraction of the b-tagged (untagged) jets that have a scale factor smaller (larger) than one.

Validation on data

Similarly to what was done with electrons and muons, some validation studies are performed to probe agreement of data with simulated samples for jets. Events containing or $Z \rightarrow \mu^+ \mu^-$ were selecting, following the same trigger and lepton selection criteria of the main analysis, but increasing the p_T threshold on leptons to 30 GeV and asking the dilepton invariant mass window of 70-110 GeV. Basic properties of additional jets in an event, requiring $p_T > 30$ GeV and $\eta < 4.7$, were then compared. Figure 1.12 shows comparison of data with the MC in bins of pseudo-rapidity of the leading jet. Simulated samples of Drell-Yan and $\bar{\nu}$ are used. Uncertainties from the jet energy corrections are also displayed on ratio plots of data to MC.

As it is shown, Jet Energy Resolution corrections are creating the “horns” one can see in the MC for η of about 2.8. Although not satisfactory, the disagreement of data from MC we see in this region is covered by the very large uncertainty. After discussion with jet experts in the JetMET POG, it has been decided that this is the configuration we should use for this preliminary result. We also verified that it was not inducing significant migration of events in the signal MC between the categories.

The disagreement of data with MC for the MET distribution prevents us at this stage to use this variable in the analysis, in particular to better target VH events where, for instance, a Z would decay into neutrinos.




FIGURE 1.12: Comparison of data with MC for leading jet η in $Z \rightarrow \ell\ell$ +jets. MC is normalized to data. Jet ID and Jet PUID are applied. MC samples include DY and $t\bar{t}$. Data/MC ratio plots are shown in the bottom of each plots, together with the uncertainties (shaded histograms) from Jet Energy Corrections.

1.1.6 Object Summary

The requirements on all objects used in the analysis (with 2018 data) are summed up in the Table 1.4. In addition, a “ghost-cleaning” procedure is applied to the muons, as described in Sec. 1.1.3.

A lepton is declared **loose** if it passes the reconstruction, kinematics and d_{xy}/dz cuts and declared **tight** if it passes in addition the identification, isolation and SIP3D cut.

TABLE 1.4: Chart of selection of physics objects for the $H \rightarrow ZZ \rightarrow 4\ell$ analysis used on the 2018 data.

Electrons
$p_T^e > 7 \text{ GeV} \quad \text{and} \quad \eta^e < 2.5$ $d_{xy} < 0.5 \text{ cm} \quad \text{and} \quad d_z < 1 \text{ cm}$ $ SIP_{3D} < 4$ BDT ID with isolation (Fall17v2 training) with cuts from Table 1.2
Muons
Global or Tracker Muon Discard Standalone Muon tracks if reconstructed in muon system only Discard muons with muonBestTrackType==2 even if they are global or tracker muons $p_T^\mu > 5 \text{ GeV} \quad \text{and} \quad \eta^\mu < 2.4$ $d_{xy} < 0.5 \text{ cm} \quad \text{and} \quad d_z < 1 \text{ cm}$ $ SIP_{3D} < 4$ The PF muon ID if $p_T < 200 \text{ GeV}$, PF muon ID or High- p_T muon ID (Table 1.3) if $p_T > 200 \text{ GeV}$ $\mathcal{I}_{PF}^\mu < 0.35$
The FSR photons
$p_T^\gamma > 2 \text{ GeV} \quad \text{and} \quad \eta^\gamma < 2.4$ $\mathcal{I}_{PF}^\gamma < 1.8$ $\Delta R(\ell, \gamma) < 0.5 \quad \text{and} \quad \frac{\Delta R(\ell, \gamma)}{(p_T^\gamma)^2} < 0.012 \text{ GeV}^{-2}$
Jets
$p_T^{\text{jet}} > 30 \text{ GeV} \quad \text{and} \quad \eta^{\text{jet}} < 4.7$ $\Delta R(\ell/\gamma, \text{jet}) > 0.4$ Cut-based jet ID (tight WP) Jet pileup ID (tight WP) Deep CSV algorithm for b-tagging (medium WP)

Bibliography

- [1] ATLAS Collaboration. “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”. In: *Physics Letters B* 716 (2012), pp. 1–29.
- [2] CMS Collaboration. “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”. In: *Physics Letters B* 716 (2012), pp. 30–61.
- [3] ATLAS Collaboration. “Evidence for the spin-0 nature of the Higgs boson using ATLAS data”. In: *Phys. Lett. B* 726 (2013), pp. 120–144.
- [4] ATLAS and CMS Collaboration. “Combined measurement of the Higgs Boson mass in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS experiment”. In: *Physics Rev. Lett.* 115 (2015), p. 191803.
- [5] ATLAS Collaboration. “Study of the spin and parity of the Higgs Boson in diboson decays with the ATLAS detector”. In: *Eur. Phys. Phys. J. C* 75 (2015), p. 476.
- [6] CMS Collaboration. “Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV”. In: *Phys. Rev. D* 92 (2015), p. 012004.
- [7] G. Aad et al. “Measurements of fiducial and differential cross sections for Higgs boson production in the diphoton decay channel at $\sqrt{s} = 8$ TeV with ATLAS”. In: *JHEP* 2014.9 (2014), pp. 1–61.
- [8] G. Aad et al. “Measurement of the Higgs boson production cross section at 7, 8 and 13 TeV center-of-mass energies in the channel $H \rightarrow \gamma\gamma$ with the ATLAS detector”. In: *JHEP* 2014.9 (2014), pp. 1–61.
- [9] G. Aad et al. “Fiducial and differential cross sections of Higgs boson production measured in the four-lepton decay channel in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”. In: *Physics Letters B* 738 (2014), pp. 234–253.

- [10] ATLAS collaboration. “Measurements of the Higgs boson production cross section at 7, 8 and 13 TeV centre-of-mass energies and search for new physics at 13 TeV in the $H \rightarrow ZZ^* \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$ final state with the ATLAS detector”. In: *ATLAS-CONF-2015-059* (2015).
- [11] G. Aad et al. “Measurements of the Total and Differential Higgs Boson Production Cross Sections Combining the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ Decay Channels at $\sqrt{s} = 8$ TeV with the ATLAS Detector”. In: *Phys. Rev. Lett.* 115 (2014), p. 091801.
- [12] ATLAS collaboration. “Measurements of the total cross sections for Higgs boson production combining the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channels at 7, 8 and 13 TeV center-of-mass energies with the ATLAS detector”. In: *ATLAS Note ATLAS-CONF-2015-069* (2015), pp. 1–61.
- [13] V. Khachatryan et al. “Measurement of differential cross sections for Higgs boson production in the diphoton decay channel in pp collisions at $\sqrt{s} = 8$ TeV”. In: *Eur. Phys. J. C* 76 (2016), pp. 1–31.
- [14] CMS Collaboration. “Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV”. In: *Eur. Phys. J. C* 75 (2015), p. 212.
- [15] Stefano Frixione, Paolo Nason, and Carlo Oleari. “Matching NLO QCD computations with Parton Shower simulations: the POWHEG method”. In: *JHEP* 11 (2007), p. 070.
- [16] Tom Melia et al. “ ZZ , $W^\pm Z$ and W^+W^- production, including γ/Z interference, singly resonant contributions and interference for identical leptons”. In: *JHEP* 11 (2011), p. 078.
- [17] S. Alioli et al. “NLO vector-boson production matched with shower in POWHEG”. In: *JHEP* 07 (2008), p. 060.
- [18] Paolo Nason. “A new method for combining NLO QCD with shower Monte Carlo algorithms”. In: *JHEP* 11 (2008), p. 040.
- [19] Stefano N Frixione, Paolo Nason, and Carlo Oleari. “Matching NLO QCD computations with parton shower simulations: the POWHEG method”. In: *JHEP* 11 (2007), p. 070.

- [20] E. Bagnaschi et al. "Higgs production via gluon fusion in the POWHEG approach in the SM and in the MSSM". In: *JHEP* 02 (2012), p. 88.
- [21] Paolo Nason and Carlo Oleari. "NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG". In: *JHEP* 02 (2010), p. 037.
- [22] Heribertus B. Hartanto et al. "Higgs boson production in association with top quarks in the POWHEG BOX". In: *Phys. Rev. D* 91 (2015), p. 094003.
- [23] Gionata Luisoni et al. " $HW^\pm/HZ + 0$ and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO". In: *JHEP* 10 (2013), p. 083.
- [24] Yanyan Gao et al. "Spin determination of single-produced resonances at hadron colliders". In: *Phys. Rev. D* 81 (2010), p. 075022.
- [25] Richard D. Ball et al. "Parton distributions for the LHC Run II". In: *JHEP* 04 (2015), p. 040.
- [26] Sara Bolognesi et al. "On the spin and parity of a single-produced resonance at the LHC". In: *Phys. Rev. D* 86 (2012), p. 095031.
- [27] Ian Anderson et al. "Constraining anomalous HVV interactions at proton and lepton colliders". In: *Phys. Rev. D* 89 (2014), p. 035007.
- [28] Paolo Nason and Giulia Zanderighi. " W^+W^- , WZ and ZZ production in the POWHEG-BOX-V2". In: *Eur. Phys. J. C* 74 (2014), p. 2702.
- [29] John M. Campbell and R. K. Ellis. "MCFM for the Tevatron and the LHC". In: *Nucl. Phys. Proc. Suppl.* 205 (2010), p. 10.
- [30] John M. Campbell, R. Keith Ellis, and Ciaran. Williams. "Bounding the Higgs width at the LHC using full analytic results for $gg \rightarrow e^-e^+\mu^-\mu^+$ ". In: *JHEP* 04 (2014), p. 060.
- [31] J. Allison et al. "Geant4 developments and applications". In: *IEEE Trans. Nucl. Sci.* 53 (2006), p. 270.
- [32] Serguei Chatrchyan et al. "Measurement of the properties of a Higgs boson in the four-lepton final state". In: *Phys. Rev. D* 89 (2014), p. 092007.
- [33] CMS Collaboration. "Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV". In: *JHEP CMS-PAS-HIG-14-018* (2014), p. 026.

- [34] Serguei Chatrchyan et al. "Measurement of the properties of a Higgs boson in the four-lepton final state". In: *Phys.Rev. D* 89 (2014), p. 092007.
- [35] Nikolas Kauer. "Inadequacy of zero-width approximation for a light Higgs boson signal". In: *Mod.Phys.Lett. A* 28 (2013), p. 1330015.
- [36] Nikolas Kauer and Giampiero Passarino. "Inadequacy of zero-width approximation for a light Higgs boson signal". In: *JHEP* 1208 (2012), p. 116.
- [37] Massimiliano Grazzini, Stefan Kallweit, and Dirk Rathlev. "ZZ production at the LHC: Fiducial cross sections and distributions in NNLO QCD". In: *Phys. Lett. B* 750 (2015), pp. 407–410. ISSN: 0370-2693. DOI: <http://dx.doi.org/10.1016/j.physletb.2015.09.055>. arXiv: 1507.06257 [hep-ph]. URL: <http://www.sciencedirect.com/science/article/pii/S0370269315007303>.
- [38] John M. Campbell, R. Keith Ellis, and Ciaran Williams. "Vector boson pair production at the LHC". In: *JHEP* 07 (2011), p. 018. DOI: [10.1007/JHEP07\(2011\)018](https://doi.org/10.1007/JHEP07(2011)018). arXiv: 1105.0020 [hep-ph].
- [39] Marco Bonvini et al. "Signal-background interference effects in $gg \rightarrow H \rightarrow WW$ beyond leading order". In: *Phys. Rev. D* 88 (2013), p. 034032. DOI: [10.1103/PhysRevD.88.034032](https://doi.org/10.1103/PhysRevD.88.034032). arXiv: 1304.3053 [hep-ph].
- [40] Kirill Melnikov and Matthew Dowling. "Production of two Z-bosons in gluon fusion in the heavy top quark approximation". In: *Phys. Lett. B* 744 (2015), p. 43. DOI: [10.1016/j.physletb.2015.03.030](https://doi.org/10.1016/j.physletb.2015.03.030). arXiv: 1503.01274 [hep-ph].
- [41] Chong Sheng Li et al. "Soft gluon resummation in the signal-background interference process of $gg(\rightarrow h^*) \rightarrow ZZ$ ". 2015.
- [42] Giampiero Passarino. "Higgs CAT". In: *Eur. Phys. J. C* 74 (2014), p. 2866. DOI: [10.1140/epjc/s10052-014-2866-7](https://doi.org/10.1140/epjc/s10052-014-2866-7). arXiv: 1312.2397 [hep-ph].
- [43] Stefano Catani and Massimiliano Grazzini. "An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC". In: *Phys. Rev. Lett.* 98 (2007), p. 222002. DOI: [10.1103/PhysRevLett.98.222002](https://doi.org/10.1103/PhysRevLett.98.222002). arXiv: hep-ph/0703012 [hep-ph].

- [44] Massimiliano Grazzini. “NNLO predictions for the Higgs boson signal in the $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ and $H \rightarrow ZZ \rightarrow 4\ell$ decay channels”. In: *JHEP* 02 (2008), p. 043. DOI: [10.1088/1126-6708/2008/02/043](https://doi.org/10.1088/1126-6708/2008/02/043). arXiv: [0801.3232](https://arxiv.org/abs/0801.3232) [hep-ph].
- [45] Massimiliano Grazzini and Hayk Sargsyan. “Heavy-quark mass effects in Higgs boson production at the LHC”. In: *JHEP* 1309 (2013), p. 129.
- [46] LHC Higgs Cross Section Working Group. *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*. CERN Report CERN-2013-004. 2013. DOI: [10.5170/CERN-2013-004](https://doi.org/10.5170/CERN-2013-004). arXiv: [1307.1347](https://arxiv.org/abs/1307.1347) [hep-ph].
- [47] Marco Bonvini et al. “Signal-background interference effects for $gg \rightarrow H \rightarrow W+W$ -beyond leading order”. In: *Phys.Rev. D* 88 (2013), p. 034032.
- [48] Glen Cowan et al. “Asymptotic formulae for likelihood-based tests of new physics”. In: *Eur. Phys. J.* 71 (2011), p. 1554.
- [49] ATLAS and LHC Higgs Combination Group CMS Collaborations. “Procedure for the LHC Higgs boson search combination in Summer 2011”. In: *ATL-PHYS-PUB/CMS NOTE 2011-11* 2011/005 (CERN, 2011).
- [50] Harrison B. Prosper and Louis Lyons. “Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding, CERN, Geneva, Switzerland 17-20 January 2011”. In: ().
- [51] V. Khachatryan et al. “Measurement of differential cross sections for Higgs boson production in the diphoton decay channel in pp collisions at $\sqrt{s} = 8$ TeV”. In: *Eur. Phys. J. C* 76 (2016), p. 13.
- [52] D. de Florian et al. “Higgs boson production at the LHC: transverse momentum resummation effects in the H -to- 2γ , H -to- WW -to- $\ell\nu\ell\nu$ and H -to- ZZ -to- 4ℓ decay modes”. In: *JHEP* 1206 (2012), p. 132.
- [53] Charalampos Anastasiou et al. “High precision determination of the gluon fusion Higgs boson cross-section at the LHC”. In: *JHEP* 2016.5 (2016), p. 1. DOI: [10.1007/JHEP05\(2016\)058](https://doi.org/10.1007/JHEP05(2016)058). arXiv: [1602.00695](https://arxiv.org/abs/1602.00695) [hep-ph].
- [54] “Measurement of differential and integrated fiducial cross sections for Higgs boson production in the four-lepton decay channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV”. In: (2015). arXiv: [1512.08377](https://arxiv.org/abs/1512.08377) [hep-ex].

-
- 601 [55] LHC Higgs Cross Section Working Group et al. “Handbook of LHC Higgs
602 Cross Sections: 3. Higgs Properties”. In: *JHEP* CERN-2013-004 (2013), p. 026.