
LECTURE 1

Author

Tom Jeong

January 8, 2025

Contents

1	D-integral domains	2
2	Characteristic of a Ring	2
3	vector Space	3


1 D-integral domains

$Q(D)$ -fields of fraction, smallest field containing D . Define

universal property $\theta: D \rightarrow K$ θ injective, K field (draw down $i: D \rightarrow Q(D)$)

sending a to $(a, 1)$

and draw $\phi: Q(D) \rightarrow K$ and say $\theta = \phi \circ i$



- $i: D \rightarrow Q(D)$ sends $a \mapsto (a, 1)$
- θ is injective
- K is a field
- $\theta = \phi \circ i$

2 Characteristic of a Ring

Let R be a unital commutative ring. The characteristic of R , denoted $\text{char}(R)$, is defined as follows:

Definition 2.1. The characteristic of a ring R is the smallest positive integer n such that

$$n \cdot 1_R = \underbrace{1_R + 1_R + \cdots + 1_R}_{n \text{ times}} = 0_R$$

If no such positive integer exists, we say $\text{char}(R) = 0$.

Proposition 2.1. For a unital commutative ring R , exactly one of the following holds:

1. $\text{char}(R) = 0$: In this case, the additive subgroup generated by 1_R is infinite.
2. $\text{char}(R) = n > 0$: In this case, n is the smallest positive integer such that $n \cdot 1_R = 0_R$.

If $k \cdot 1_R = m \cdot 1_R$ for some integers k, m , then:

$$(k - m) \cdot 1_R = 0_R$$

This means that if $\text{char}(R) = n > 0$, then n divides $k - m$.

example:

1. $\text{char}(\mathbb{Z}) = 0$
2. $\text{char}(\mathbb{Q}) = 0$

3. $\text{char}(\mathbb{F}_p) = p$ for any prime field
4. For any field K , $\text{char}(K)$ is either 0 or a prime number

Proposition 2.2. If R is a domain (i.e., has no zero divisors), then $\text{char}(R)$ is either 0 or prime.

Proof.

$s = \text{char}(K)$ and $s = ab$ where $a, b < s$

$(a \cdot 1)(b \cdot 1) = (a \cdot b) \cdot 1 = 0 \rightarrow a \cdot 1 = 0$ or $b \cdot 1 = 0$ but $a, b < s$ □

K -field:: $\text{char}(K) = 0 \rightarrow \mathbb{Q} \subseteq K$

$\text{char}(K) = p \rightarrow \mathbb{Z}_p$ or $\mathbb{F}_p \subseteq K$

\mathbb{Q} or \mathbb{Z}_p are called prime subfields of K .

3 vector Space

Proposition 3.1. Let's say a field is inside another field, $F \subseteq K$ then K is an F -vector space

So vector space over F (F field) if

1. $s_1 s_2 \in S \rightarrow s_1 + s_2 \in S$
2. $c \cdot s_1 \in S, c \in F$
3. $c(s_1 + s_2) = cs_1 + cs_2$

$\mathbb{R} \subseteq \mathbb{C}$

$a + bi$

1, i is a basis of \mathbb{C} over \mathbb{R}

$\mathbb{C} \cong \mathbb{R}^2$ but then

$\mathbb{Q} \subseteq \mathbb{R}$.. \mathbb{R} is an infinite-dimensional vector space over \mathbb{Q}

K is a field extension of F

Proposition 3.2 (freshman's dream).

if $\text{char}(K) = p, K$ field then, $(x + y)^p = x^p + y^p$

Proof. binomial expansion of

$$\begin{aligned}
(x+y)^p &= x^p + \binom{p}{1}x^{p-1}y + \binom{p}{2}x^{p-2}y^2 + \cdots + \binom{p}{p-1}xy^{p-1} + y^p \\
&= x^p + \binom{p}{1}x^{p-1}y + \binom{p}{2}x^{p-2}y^2 + \cdots + \binom{p}{p-1}xy^{p-1} + y^p \\
&= x^p + y^p
\end{aligned}$$

In characteristic p , all binomial coefficients $\binom{p}{k}$ for $1 \leq k \leq p-1$ are divisible by p , hence equal to zero in the field. \square

Let K be a field of characteristic $p > 0$. The Frobenius homomorphism $\phi : K \rightarrow K$ is defined as:

$$\begin{aligned}
\phi : K &\rightarrow K \\
x &\mapsto x^p
\end{aligned}$$

Proposition 3.3 (Properties of Frobenius). The map ϕ is a ring homomorphism:

1. $\phi(x+y) = (x+y)^p = x^p + y^p = \phi(x) + \phi(y)$ (using the binomial expansion in char p)
2. $\phi(xy) = (xy)^p = x^p y^p = \phi(x)\phi(y)$
3. $\phi(1) = 1^p = 1$

Example 3.4. In \mathbb{F}_p , the Frobenius map is the identity map since:

$$a^p = a \text{ for all } a \in \mathbb{F}_p$$

This is known as Fermat's Little Theorem.