LECTURE 7 SEP 11

Cosets

Author

Tom Jeong

September 11, 2024

Contents

1 Bijective homomorphism

3

1 Bijective homomorphism

A Bijective homomorphism is called an isomorphism.

Proposition 1.1 (2.4.9).

Let $f:G\to K$ be a group homomorphism.

- 1. the image $Imf \subseteq K$ is a subgroup of K.
- 2. The kernel $Kerf \subseteq G$ is a normal subgroup of G.
- 3. f is injective if and only if $Kerf = \{e\}$

Proof continutes.

2. Normality Let N = Kerf for every g in G and $n \in N$, $gng^{-1} \in N$.

$$f(gng^{-1}) = f(g)f(n)f(g)^{-1}$$
(1)

$$= f(g)f(g)^{-1} \tag{2}$$

$$=e$$
 (3)

so
$$(4)$$

$$gng^{-1} \in Kerf = N \forall g \in G, n \in N \tag{5}$$

Hence
$$gNg^{-1} \subseteq N \ \forall g \in G$$
 (6)

3. \rightarrow Since $f(e_G) = e_K$ and f is injective. $kerf = \{e_G\}$

 \leftarrow Suppose $kerf = \{e_G\}$ and f(g) = f(h) then $f(gh^{-1}) = f(g)f(h)^{-1} = e$ so $gh^{-1} \in kerf = \{e_G\}$ so g = h so f is injective.

Theorem 1.2 (2.5.1 Isomorphism Theorem).

Leg G and K be groups and $f:G\to K$ a homomorphism with the kernel Kerf=N, then $\tilde{f}:G/N\to f(G)$ given by $\tilde{f}(gN)=f(g)$ is well defined and a group isomorphism.

Proof. Notice that $f(x) = f(y) \iff f(y)^{-1}f(x) = e_K \iff f(y^{-1}x) = e_K \iff y^{-1}x \in N \iff xN = yN \text{ so } \tilde{f} \text{ is well defined.}$

Recall lemma 2.2.6 -ii that $y^{-1}x \in N \iff xN = yN$ so \tilde{f} . Hence

$$f(x) = f(y) \iff xN = yN$$

 \rightarrow gives that \tilde{f} is injective.

 \leftarrow give that well defined

Proposition 1.3 (2.4.9).

states that Kerf=N is normal so \tilde{f} is a homomorphism since $\tilde{f}((g_1N)(g_2N))=\tilde{f}(g_1g_2N)=f(g_1g_2)=f(g_1)f(g_2)=\tilde{f}(g_1N)\tilde{f}(g_2N)$ Now \tilde{f} is surjective because f is surjective onto f(G) so \tilde{f} is an isomorphism.