


# Algebraicity of elliptic KZB equations

Tiago J. Fonseca

Unicamp, FAPESP

November 2022, Campinas

Genus 0



- Let  $X$  be a smooth manifold and  $A^\bullet(X)$  be the complex of smooth differential forms. **De Rham's** theorem:

$$H^n(X, \mathbb{R}) \cong H^n(A^\bullet(X))$$





- Line integrals cannot detect non-abelian phenomena in  $\pi_1(X)$ :

$$\int_{aba^{-1}b^{-1}} \omega = \int_a \omega + \int_b \omega - \int_a \omega - \int_b \omega = 0.$$

- Line integrals cannot detect non-abelian phenomena in  $\pi_1(X)$ :

$$\int_{aba^{-1}b^{-1}} \omega = \int_a \omega + \int_b \omega - \int_a \omega - \int_b \omega = 0.$$

- **Chen:** consider 'iterated line integrals'. Let  $c : [0, 1] \rightarrow X$  be a path and  $\omega_i \in A^1(X)$ . Define

$$\int_c \omega_1 \cdots \omega_r := \int_{0 \leq t_1 \leq \cdots \leq t_r \leq 1} f_1(t_1) \cdots f_r(t_r) dt_1 \cdots dt_r.$$

where  $c^* \omega_i = f_i(t_i) dt_i$ .

- Line integrals cannot detect non-abelian phenomena in  $\pi_1(X)$ :

$$\int_{aba^{-1}b^{-1}} \omega = \int_a \omega + \int_b \omega - \int_a \omega - \int_b \omega = 0.$$


- Chen:** consider 'iterated line integrals'. Let  $c : [0, 1] \rightarrow X$  be a path and  $\omega_i \in A^1(X)$ . Define

$$\int_c \omega_1 \cdots \omega_r := \int_{0 \leq t_1 \leq \cdots \leq t_r \leq 1} f_1(t_1) \cdots f_r(t_r) dt_1 \cdots dt_r.$$

where  $c^* \omega_i = f_i(t_i) dt_i$ .

- Example:  $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$

$$\int_{aba^{-1}b^{-1}} \frac{dz}{z} \frac{dz}{z-1} = (2\pi i)^2.$$





- Let  $x \in X$  and  $Ch(P_x X, \mathbb{R})$  be the space of iterated integrals

$$\int \omega_1 \cdots \omega_r = \left( c \mapsto \int_c \omega_1 \cdots \omega_r \right)$$

where  $c$  denotes a loop based at  $x$ .

- ▶ Let  $x \in X$  and  $Ch(P_x X, \mathbb{R})$  be the space of iterated integrals

$$\int \omega_1 \cdots \omega_r = \left( c \mapsto \int_c \omega_1 \cdots \omega_r \right)$$

where  $c$  denotes a loop based at  $x$ .

- ▶ Let  $I$  be the augmentation ideal of  $\mathbb{Z}[\pi_1(X, x)]$ . **Chen's** theorem:

$$\varinjlim_n Hom(\mathbb{Z}[\pi_1(X, x)]/I^{n+1}, \mathbb{R}) \cong H^0 Ch(P_x X, \mathbb{R}),$$

where RHS is the subspace of homotopy-invariant iterated integrals.

- ▶ Let  $x \in X$  and  $Ch(P_x X, \mathbb{R})$  be the space of iterated integrals

$$\int \omega_1 \cdots \omega_r = \left( c \mapsto \int_c \omega_1 \cdots \omega_r \right)$$

where  $c$  denotes a loop based at  $x$ .

- ▶ Let  $I$  be the augmentation ideal of  $\mathbb{Z}[\pi_1(X, x)]$ . **Chen's** theorem:

$$\varinjlim_n Hom(\mathbb{Z}[\pi_1(X, x)]/I^{n+1}, \mathbb{R}) \cong H^0 Ch(P_x X, \mathbb{R}),$$

where RHS is the subspace of homotopy-invariant iterated integrals.

- ▶ Remarks:
  - ▶ The LHS is dual to the **Mal'cev** completion of  $\pi_1(X, x)$ .
  - ▶ Can replace  $\mathbb{R}$  by  $\mathbb{C}$ .
  - ▶ There is also a version with two base points  $x, y$ .

► If  $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$ ,

$$H^0 Ch(P_X X, \mathbb{C}) \cong \mathbb{Q}\langle \omega_0, \omega_1 \rangle \otimes \mathbb{C}, \quad \omega_0 = \frac{dz}{z}, \quad \omega_1 = \frac{dz}{1-z}$$

- If  $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$ ,

$$H^0 Ch(P_X X, \mathbb{C}) \cong \mathbb{Q}\langle \omega_0, \omega_1 \rangle \otimes \mathbb{C}, \quad \omega_0 = \frac{dz}{z}, \quad \omega_1 = \frac{dz}{1-z}$$

- Multiple zeta value (MZV):

$$\zeta(n_1, \dots, n_r) = \sum_{0 < k_1 < \dots < k_r} \frac{1}{k_1^{n_1} \dots k_r^{n_r}}$$

$$\stackrel{\text{Kontsevich}}{=} \int_0^1 \omega_1 \underbrace{\omega_0 \dots \omega_0}_{n_1-1} \dots \omega_1 \underbrace{\omega_0 \dots \omega_0}_{n_r-1}$$

Relations are explained by the integral formula. Example:  
 $\zeta(3) = \zeta(1, 2)$  comes from change of variables  $z \mapsto 1 - z$ .

- ▶ If  $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$ ,

$$H^0 Ch(P_X X, \mathbb{C}) \cong \mathbb{Q}\langle \omega_0, \omega_1 \rangle \otimes \mathbb{C}, \quad \omega_0 = \frac{dz}{z}, \quad \omega_1 = \frac{dz}{1-z}$$

- ▶ Multiple zeta value (MZV):

$$\zeta(n_1, \dots, n_r) = \sum_{0 < k_1 < \dots < k_r} \frac{1}{k_1^{n_1} \dots k_r^{n_r}}$$

$$\stackrel{\text{Kontsevich}}{=} \int_0^1 \omega_1 \underbrace{\omega_0 \dots \omega_0}_{n_1-1} \dots \omega_1 \underbrace{\omega_0 \dots \omega_0}_{n_r-1}$$

Relations are explained by the integral formula. Example:  
 $\zeta(3) = \zeta(1, 2)$  comes from change of variables  $z \mapsto 1 - z$ .

- ▶ **Brown's theorem:** every MZV is a  $\mathbb{Q}$ -linear combination of MZVs of the form  $\zeta(n_1, \dots, n_r)$  with  $n_i \in \{2, 3\}$ .

► Polylogarithm:

$$Li_n(z) = \int_0^z \omega_1 \underbrace{\omega_0 \cdots \omega_0}_{n-1} = \sum_{k=1}^{\infty} \frac{z^k}{k^n}.$$

Values at  $z \in \overline{\mathbb{Q}}$  related to regulators of number fields.

► Polylogarithm:

$$Li_n(z) = \int_0^z \omega_1 \underbrace{\omega_0 \cdots \omega_0}_{n-1} = \sum_{k=1}^{\infty} \frac{z^k}{k^n}.$$

Values at  $z \in \overline{\mathbb{Q}}$  related to regulators of number fields.

► Multiple polylogarithm (MPL):

$$Li_{n_1, \dots, n_r}(z) = \int_0^z \omega_1 \omega_0^{n_1-1} \cdots \omega_1 \omega_0^{n_r-1} = \sum_{0 < k_1 < \dots < k_r} \frac{z^{k_r}}{k_1^{n_1} \cdots k_r^{n_r}}$$



- Polylogarithm:

$$Li_n(z) = \int_0^z \omega_1 \underbrace{\omega_0 \cdots \omega_0}_{n-1} = \sum_{k=1}^{\infty} \frac{z^k}{k^n}.$$

Values at  $z \in \overline{\mathbb{Q}}$  related to regulators of number fields.

- Multiple polylogarithm (MPL):

$$Li_{n_1, \dots, n_r}(z) = \int_0^z \omega_1 \omega_0^{n_1-1} \cdots \omega_1 \omega_0^{n_r-1} = \sum_{0 < k_1 < \dots < k_r} \frac{z^{k_r}}{k_1^{n_1} \cdots k_r^{n_r}}$$

- More generally, if  $w$  is a word in  $x_0, x_1$ , define

$$Li_w(z)$$

in such a way that  $Li_{n_1, \dots, n_r}(z) = Li_{x_0^{n_1-1} x_1 \cdots x_0^{n_r-1} x_1}(z)$ .

- Differential equations:

$$dLi_{x_i w}(z) = \omega_i Li_w(z), \quad i = 0, 1$$

- Differential equations:

$$dLi_{x_i w}(z) = \omega_i Li_w(z), \quad i = 0, 1$$

- Equivalently, the generating series

$$\begin{aligned} L(z) &= \sum_w Li_w(z)w \\ &= 1 + Li_{x_0}(z)x_0 + Li_{x_1}(z)x_1 + \cdots \in \mathbb{C}\langle\langle x_0, x_1 \rangle\rangle \end{aligned}$$

is a solution of

$$dL(z) = (\omega_0 x_0 + \omega_1 x_1)L(z).$$

- Differential equations:

$$dLi_{x_i w}(z) = \omega_i Li_w(z), \quad i = 0, 1$$

- Equivalently, the generating series

$$\begin{aligned} L(z) &= \sum_w Li_w(z) w \\ &= 1 + Li_{x_0}(z) x_0 + Li_{x_1}(z) x_1 + \cdots \in \mathbb{C} \langle\langle x_0, x_1 \rangle\rangle \end{aligned}$$

is a solution of

$$dL(z) = (\omega_0 x_0 + \omega_1 x_1) L(z).$$

- KZ connection (**Knizhnik–Zamolodchikov**):

$$\nabla_{KZ} = d - \omega_0 \otimes x_0 - \omega_1 \otimes x_1$$

on the trivial infinite-rank vector bundle  $\mathcal{V}_{KZ} \rightarrow \mathbb{P}^1 \setminus \{0, 1, \infty\}$   
with fibres  $\mathbb{C} \langle\langle x_0, x_1 \rangle\rangle$ .

- ▶ The KZ connection is algebraic and defined over  $\mathbb{Q}$ .

- ▶ The KZ connection is algebraic and defined over  $\mathbb{Q}$ .
- ▶ Let  $b \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ . For every **unipotent** flat connection  $(\mathcal{V}, \nabla)$  over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$  with  $v \in \mathcal{V}_b$ , there is a unique horizontal map

$$\varphi : (\mathcal{V}_{KZ}, \nabla_{KZ}) \rightarrow (\mathcal{V}, \nabla), \quad \varphi(1) = v.$$

- ▶ The KZ connection is algebraic and defined over  $\mathbb{Q}$ .
- ▶ Let  $b \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ . For every **unipotent** flat connection  $(\mathcal{V}, \nabla)$  over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$  with  $v \in \mathcal{V}_b$ , there is a unique horizontal map

$$\varphi : (\mathcal{V}_{KZ}, \nabla_{KZ}) \rightarrow (\mathcal{V}, \nabla), \quad \varphi(1) = v.$$

- ▶  $(\mathcal{V}, \nabla)$  is unipotent if there is a filtration

$$0 = (\mathcal{V}_0, \nabla_0) \subset (\mathcal{V}_1, \nabla_1) \subset \cdots \subset (\mathcal{V}_n, \nabla_n) = (\mathcal{V}, \nabla)$$

such that  $(\mathcal{V}_i, \nabla_i)/(\mathcal{V}_{i-1}, \nabla_{i-1})$  is trivial for every  $i$ .

- ▶ The KZ connection is algebraic and defined over  $\mathbb{Q}$ .
- ▶ Let  $b \in \mathbb{P}^1 \setminus \{0, 1, \infty\}$ . For every **unipotent** flat connection  $(\mathcal{V}, \nabla)$  over  $\mathbb{P}^1 \setminus \{0, 1, \infty\}$  with  $v \in \mathcal{V}_b$ , there is a unique horizontal map

$$\varphi : (\mathcal{V}_{KZ}, \nabla_{KZ}) \rightarrow (\mathcal{V}, \nabla), \quad \varphi(1) = v.$$

- ▶  $(\mathcal{V}, \nabla)$  is unipotent if there is a filtration

$$0 = (\mathcal{V}_0, \nabla_0) \subset (\mathcal{V}_1, \nabla_1) \subset \cdots \subset (\mathcal{V}_n, \nabla_n) = (\mathcal{V}, \nabla)$$

such that  $(\mathcal{V}_i, \nabla_i)/(\mathcal{V}_{i-1}, \nabla_{i-1})$  is trivial for every  $i$ .


- ▶ KZ is so nice (constant vector bundle, forms with log singularities) because

$$H^0(\mathbb{P}^1, \mathcal{O}) = \mathbb{C}, \quad H^1(\mathbb{P}^1, \mathcal{O}) = 0$$

Every unipotent vector bundle over  $\mathbb{P}^1$  is *canonically* trivial.



# Genus 1



- ▶ Let  $(E, O)$  be a complex elliptic curve.

- ▶ Let  $(E, O)$  be a complex elliptic curve.
- ▶ As a Riemann surface

$$(E, O) \cong (\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), 0) \cong (\mathbb{C}^\times / q^{\mathbb{Z}}, 1)$$

with  $\text{Im}(\tau) > 0$  and  $q = e^{2\pi i\tau}$ .

- ▶ Let  $(E, O)$  be a complex elliptic curve.
- ▶ As a Riemann surface

$$(E, O) \cong (\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), 0) \cong (\mathbb{C}^\times/q^{\mathbb{Z}}, 1)$$

with  $\text{Im}(\tau) > 0$  and  $q = e^{2\pi i\tau}$ .

- ▶ Are there analogues of MPLs defined over  $E \setminus \{O\}$ ?

- ▶ Let  $(E, O)$  be a complex elliptic curve.
- ▶ As a Riemann surface

$$(E, O) \cong (\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), 0) \cong (\mathbb{C}^\times / q^{\mathbb{Z}}, 1)$$

with  $\text{Im}(\tau) > 0$  and  $q = e^{2\pi i\tau}$ .

- ▶ Are there analogues of MPLs defined over  $E \setminus \{O\}$ ?
- ▶ **Bloch's** elliptic dilogarithm:

$$D_E(z) = \sum_{m \in \mathbb{Z}} \mathcal{L}_2(q^m z)$$

where  $\mathcal{L}_2(z) = -2i\text{Im}(Li_2(z)) + 2\log|z|\log(1 - \bar{z})$ .

- ▶ Let  $(E, O)$  be a complex elliptic curve.
- ▶ As a Riemann surface

$$(E, O) \cong (\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), 0) \cong (\mathbb{C}^\times/q^{\mathbb{Z}}, 1)$$

with  $\text{Im}(\tau) > 0$  and  $q = e^{2\pi i\tau}$ .

- ▶ Are there analogues of MPLs defined over  $E \setminus \{O\}$ ?
- ▶ **Bloch's** elliptic dilogarithm:

$$D_E(z) = \sum_{m \in \mathbb{Z}} \mathcal{L}_2(q^m z)$$

where  $\mathcal{L}_2(z) = -2i \text{Im}(Li_2(z)) + 2 \log |z| \log(1 - \bar{z})$ .

- ▶ There are general theories of elliptic MPLs as iterated integrals on elliptic curves (e.g. **Brown–Levin**), but the underlying algebraic geometry is not clear.

- ▶ The analogue of the KZ equation for a punctured elliptic curve is the *elliptic KZB equation* (**Bernard**). Taking account of the moduli of elliptic curves, we get the *universal elliptic KZB equation*.

- ▶ The analogue of the KZ equation for a punctured elliptic curve is the *elliptic KZB equation* (**Bernard**). Taking account of the moduli of elliptic curves, we get the *universal elliptic KZB equation*.
- ▶ **Calaque–Enriquez–Etingof, Levin–Racinet, Hain**: explicit formula.

9.2. **The formula.** The connection is defined by a 1-form

$$\omega \in \Omega^1(\mathbb{C} \times \mathfrak{h}, \log \Lambda) \hat{\otimes} \text{End } \mathfrak{p}.$$

via the formula

$$\nabla f = df + \omega f$$

where  $f : \mathbb{C} \times \mathfrak{h} \rightarrow \mathfrak{p}$  is a (locally defined) section of (9.1). Specifically,

$$\omega = \frac{1}{2\pi i} d\tau \otimes \mathbf{a} \frac{\partial}{\partial \mathbf{t}} + \psi + \nu$$

where

$$\psi = \sum_{m \geq 1} \left( \frac{(2\pi i)^{2m+1}}{(2m)!} G_{2m+2}(\tau) d\tau \otimes \sum_{\substack{j+k=2m+1 \\ j, k > 0}} (-1)^j [\text{ad}_{\mathbf{t}}^j(\mathbf{a}), \text{ad}_{\mathbf{t}}^k(\mathbf{a})] \frac{\partial}{\partial \mathbf{a}} \right)$$

and

$$\nu = \mathbf{t} F(\xi, \mathbf{t}, \tau) \cdot \mathbf{a} d\xi + \frac{1}{2\pi i} \left( \frac{1}{\mathbf{t}} + \mathbf{t} \frac{\partial F}{\partial \mathbf{t}}(\xi, \mathbf{t}, \tau) \right) \cdot \mathbf{a} d\tau.$$



- ▶ The analogue of the KZ equation for a punctured elliptic curve is the *elliptic KZB equation* (**Bernard**). Taking account of the moduli of elliptic curves, we get the *universal elliptic KZB equation*.
- ▶ **Calaque–Enriquez–Etingof, Levin–Racinet, Hain**: explicit formula.

9.2. **The formula.** The connection is defined by a 1-form

$$\omega \in \Omega^1(\mathbb{C} \times \mathfrak{h}, \log \Lambda) \hat{\otimes} \text{End } \mathfrak{p}.$$

via the formula

$$\nabla f = df + \omega f$$

where  $f : \mathbb{C} \times \mathfrak{h} \rightarrow \mathfrak{p}$  is a (locally defined) section of (9.1). Specifically,

$$\omega = \frac{1}{2\pi i} d\tau \otimes \mathbf{a} \frac{\partial}{\partial \mathbf{t}} + \psi + \nu$$

where

$$\psi = \sum_{m \geq 1} \left( \frac{(2\pi i)^{2m+1}}{(2m)!} G_{2m+2}(\tau) d\tau \otimes \sum_{\substack{j+k=2m+1 \\ j,k > 0}} (-1)^j [\text{ad}_{\mathbf{t}}^j(\mathbf{a}), \text{ad}_{\mathbf{t}}^k(\mathbf{a})] \frac{\partial}{\partial \mathbf{a}} \right)$$

and

$$\nu = \mathbf{t} F(\xi, \mathbf{t}, \tau) \cdot \mathbf{a} d\xi + \frac{1}{2\pi i} \left( \frac{1}{\mathbf{t}} + \mathbf{t} \frac{\partial F}{\partial \mathbf{t}}(\xi, \mathbf{t}, \tau) \right) \cdot \mathbf{a} d\tau.$$

- ▶ Problem (cf. **Luo**): is this algebraic? Defined over  $\mathbb{Q}$ ?

- ▶ Two main difficulties for a purely algebraic theory:
  1.  $H^1(E, \mathcal{O}) \neq 0$ , i.e., there are non-trivial unipotent vector bundles over  $E$ ,
  2. presence of moduli.

- ▶ Two main difficulties for a purely algebraic theory:
  1.  $H^1(E, \mathcal{O}) \neq 0$ , i.e., there are non-trivial unipotent vector bundles over  $E$ ,
  2. presence of moduli.
- ▶ Joint work in progress with **Nils Matthes**: work over the  $\mathbb{A}^1$ -bundle

$$\pi : E^{\natural} \rightarrow E$$

such that  $H^0(E^{\natural}, \mathcal{O}) = \mathbb{C}$  and  $H^1(E^{\natural}, \mathcal{O}) = 0$  (cf. **Deligne, Enriquez–Etingof**).

- ▶ Two main difficulties for a purely algebraic theory:
  1.  $H^1(E, \mathcal{O}) \neq 0$ , i.e., there are non-trivial unipotent vector bundles over  $E$ ,
  2. presence of moduli.
- ▶ Joint work in progress with **Nils Matthes**: work over the  $\mathbb{A}^1$ -bundle

$$\pi : E^{\natural} \rightarrow E$$

such that  $H^0(E^{\natural}, \mathcal{O}) = \mathbb{C}$  and  $H^1(E^{\natural}, \mathcal{O}) = 0$  (cf. **Deligne, Enriquez–Etingof**).

- ▶  $E^{\natural}$  is known in the literature as the *universal vector extension* of  $E$  (**Rosenlicht, Serre, Grothendieck**, etc.). It is the moduli space of *algebraic* flat line bundles on  $E$  and

$$\pi : E^{\natural} \rightarrow E \cong \text{Pic}^0(E), \quad [(\mathcal{L}, \nabla)] \mapsto [\mathcal{L}].$$

## Theorem (F.–Matthes; cf. Enriques–Etingof)

Set  $D = \pi^{-1}(O)$ . There is a canonical decomposition

$$\Gamma(E^{\natural}, \Omega^1(\log D)) = \Gamma(E^{\natural}, \Omega^1) \oplus K^{(1)} \oplus K^{(2)} \oplus \dots$$

where  $K^{(n)}$  are 1-dimensional subspaces uniquely determined by:

1.  $dK^{(n)} = \Gamma(E^{\natural}, \Omega^1) \wedge K^{(n-1)}$ , where  $K^{(0)} := \pi^* \Gamma(E, \Omega^1)$ ,
2.  $K^{(n)} \wedge K^{(0)} = 0$ ,
3.  $\text{Res}_D(K^{(n)})$  has degree  $n - 1$ .

## Theorem (F.–Matthes; cf. Enríquez–Etingof)

Set  $D = \pi^{-1}(O)$ . There is a canonical decomposition

$$\Gamma(E^{\natural}, \Omega^1(\log D)) = \Gamma(E^{\natural}, \Omega^1) \oplus K^{(1)} \oplus K^{(2)} \oplus \dots$$

where  $K^{(n)}$  are 1-dimensional subspaces uniquely determined by:

1.  $dK^{(n)} = \Gamma(E^{\natural}, \Omega^1) \wedge K^{(n-1)}$ , where  $K^{(0)} := \pi^* \Gamma(E, \Omega^1)$ ,
2.  $K^{(n)} \wedge K^{(0)} = 0$ ,
3.  $\text{Res}_D(K^{(n)})$  has degree  $n - 1$ .

We can find  $\nu, \omega^{(0)}, \omega^{(1)}, \omega^{(2)}, \dots$  such that

$$\Gamma(E^{\natural}, \Omega^1) = \mathbb{C}\nu \oplus \mathbb{C}\omega^{(0)}, \quad K^{(n)} = \mathbb{C}\omega^{(n)}$$

and  $d\omega^{(n)} = \nu \wedge \omega^{(n-1)}$ ,  $\omega^{(n)} \wedge \omega^{(0)} = 0$ ,  $\text{Res}_D(\omega^{(n)}) = t^{n-1}/(n-1)!$ .

Let  $\mathcal{V}_{KZB}$  be the trivial vector bundle over  $E^{\natural} \setminus D$  with fibres  $\mathbb{C}\langle\langle a, b \rangle\rangle$ , and

$$\nabla_{KZB}^{vert} = d - \nu \otimes a - \sum_{n \geq 0} \omega^{(n)} \otimes ad_a^n b$$

By the last theorem,  $(\mathcal{V}_{KZB}, \nabla_{KZB}^{vert})$  is flat. Fix  $b \in E^{\natural} \setminus D$ .

Let  $\mathcal{V}_{KZB}$  be the trivial vector bundle over  $E^{\natural} \setminus D$  with fibres  $\mathbb{C}\langle\langle a, b \rangle\rangle$ , and

$$\nabla_{KZB}^{vert} = d - \nu \otimes a - \sum_{n \geq 0} \omega^{(n)} \otimes ad_a^n b$$

By the last theorem,  $(\mathcal{V}_{KZB}, \nabla_{KZB}^{vert})$  is flat. Fix  $b \in E^{\natural} \setminus D$ .

### Theorem (F.–Matthes)

*For every unipotent flat connection  $(\mathcal{V}, \nabla)$  on  $E^{\natural} \setminus D$  with  $v \in \mathcal{V}_b$ , there is a unique horizontal map*

$$\varphi : (\mathcal{V}_{KZB}, \nabla_{KZB}^{vert}) \rightarrow (\mathcal{V}, \nabla), \quad \varphi(1) = v.$$



Let  $\mathcal{V}_{KZB}$  be the trivial vector bundle over  $E^{\natural} \setminus D$  with fibres  $\mathbb{C}\langle\langle a, b \rangle\rangle$ , and

$$\nabla_{KZB}^{vert} = d - \nu \otimes a - \sum_{n \geq 0} \omega^{(n)} \otimes ad_a^n b$$

By the last theorem,  $(\mathcal{V}_{KZB}, \nabla_{KZB}^{vert})$  is flat. Fix  $b \in E^{\natural} \setminus D$ .

### Theorem (F.–Matthes)

*For every unipotent flat connection  $(\mathcal{V}, \nabla)$  on  $E^{\natural} \setminus D$  with  $v \in \mathcal{V}_b$ , there is a unique horizontal map*

$$\varphi : (\mathcal{V}_{KZB}, \nabla_{KZB}^{vert}) \rightarrow (\mathcal{V}, \nabla), \quad \varphi(1) = v.$$

- ▶ Can be descended to  $E$ .
- ▶ If  $E/\mathbb{Q}$ ,  $(\mathcal{V}_{KZB}, \nabla_{KZB}^{vert})$  is also defined over  $\mathbb{Q}$ .



## Theorem (F.–Matthes)

*The sequence*

$$0 \longrightarrow \Omega_S^1 \longrightarrow f_* \Omega_{E^\natural}^1(\log D) \longrightarrow f_* \Omega_{E^\natural/S}^1(\log D) \longrightarrow 0$$

*is exact and canonically split.*

## Theorem (F.–Matthes)

*The sequence*

$$0 \longrightarrow \Omega_S^1 \longrightarrow f_* \Omega_{E^\natural}^1(\log D) \longrightarrow f_* \Omega_{E^\natural/S}^1(\log D) \longrightarrow 0$$

*is exact and canonically split.*

► Let  $E \rightarrow \mathbb{H}$  with  $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ .

## Theorem (F.–Matthes)

*The sequence*

$$0 \longrightarrow \Omega_S^1 \longrightarrow f_* \Omega_{E/S}^1(\log D) \longrightarrow f_* \Omega_{E/S}^1(\log D) \longrightarrow 0$$

*is exact and canonically split.*

- ▶ Let  $E \rightarrow \mathbb{H}$  with  $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ .
- ▶ Relative form in  $f_* \Omega_{E/\mathbb{H}}^1(\log D)$ :

$$\omega^{(1)} = \left( \frac{\theta'_\tau(z)}{\theta_\tau(z)} + 2\pi i r \right) dz$$

## Theorem (F.–Matthes)

The sequence

$$0 \longrightarrow \Omega_S^1 \longrightarrow f_* \Omega_{E/\mathbb{H}}^1(\log D) \longrightarrow f_* \Omega_{E/\mathbb{H}/S}^1(\log D) \longrightarrow 0$$

is exact and *canonically* split.

- ▶ Let  $E \rightarrow \mathbb{H}$  with  $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ .
- ▶ Relative form in  $f_* \Omega_{E/\mathbb{H}}^1(\log D)$ :

$$\omega^{(1)} = \left( \frac{\theta'_\tau(z)}{\theta_\tau(z)} + 2\pi i r \right) dz$$

- ▶ Canonical lift to  $f_* \Omega_E^1(\log D)$ :

$$\tilde{\omega}^{(1)} = \left( \frac{\theta'_\tau(z)}{\theta_\tau(z)} + 2\pi i r \right) dz + \frac{1}{2\pi i} \left( \frac{1}{2} \frac{\theta''_\tau(z)}{\theta_\tau(z)} - \frac{1}{6} \frac{\theta'''_\tau(0)}{\theta'_\tau(0)} - \frac{(2\pi i r)^2}{2} \right) d\tau$$

Thank you!