

On universal vector extensions of elliptic curves and elliptic KZB equations

Tiago J. Fonseca

IMECC - Unicamp

October 2024, Mathamsud conference
IMECC, Campinas

Joint work with **Nils Matthes**.

- ▶ *Towards algebraic iterated integrals for elliptic curves via the universal vectorial extension.*
RIMS Kôkyurôku, No. 2160 (2020).
- ▶ *A note on the Gauss-Manin connection for abelian schemes.*
Rend. Sem. Mat. Univ. Padova 152 (2024).
- ▶ *Elliptic KZB connections via universal vector extensions.*
To appear in Algebra & Number Theory.

- ▶ Elliptic curve (E, O) over k : smooth genus 1 curve E over k with a marked point $O \in E(k)$.

- ▶ Elliptic curve (E, O) over k : smooth genus 1 curve E over k with a marked point $O \in E(k)$.
- ▶ If $\text{char}(k) \neq 2, 3$, Weierstrass equation

$$y^2 = 4x^3 - g_2x - g_3, \quad g_2^3 \neq 27g_3^2.$$

- ▶ Elliptic curve (E, O) over k : smooth genus 1 curve E over k with a marked point $O \in E(k)$.
- ▶ If $\text{char}(k) \neq 2, 3$, Weierstrass equation

$$y^2 = 4x^3 - g_2x - g_3, \quad g_2^3 \neq 27g_3^2.$$

- ▶ If $k = \mathbb{C}$, uniformisation:

$$E^{an} = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), \quad \tau \in \mathbb{H}$$

- ▶ Elliptic curve (E, O) over k : smooth genus 1 curve E over k with a marked point $O \in E(k)$.
- ▶ If $\text{char}(k) \neq 2, 3$, Weierstrass equation

$$y^2 = 4x^3 - g_2x - g_3, \quad g_2^3 \neq 27g_3^2.$$

- ▶ If $k = \mathbb{C}$, uniformisation:

$$E^{an} = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}), \quad \tau \in \mathbb{H}$$

- ▶ Related via elliptic functions

$$z \mapsto (\wp_\tau(z), \wp'_\tau(z))$$

Coefficients $g_2(\tau), g_3(\tau)$ are modular forms.

- Moduli of elliptic curves $\mathcal{M}_{1,1}$. Uniformisation:

$$\mathcal{M}_{1,1}^{an} = SL_2(\mathbb{Z}) \backslash \mathbb{H}$$

- ▶ Moduli of elliptic curves $\mathcal{M}_{1,1}$. Uniformisation:

$$\mathcal{M}_{1,1}^{an} = SL_2(\mathbb{Z}) \backslash \mathbb{H}$$

- ▶ Moduli of genus 1 curves with 2 marked points $\mathcal{M}_{1,2}$. Same as universal family of elliptic curves minus zero section

$$\mathcal{E} \setminus \mathcal{O} = \mathcal{M}_{1,2} \rightarrow \mathcal{M}_{1,1}$$

- ▶ Moduli of elliptic curves $\mathcal{M}_{1,1}$. Uniformisation:

$$\mathcal{M}_{1,1}^{an} = SL_2(\mathbb{Z}) \backslash \mathbb{H}$$


- ▶ Moduli of genus 1 curves with 2 marked points $\mathcal{M}_{1,2}$. Same as universal family of elliptic curves minus zero section

$$\mathcal{E} \setminus O = \mathcal{M}_{1,2} \rightarrow \mathcal{M}_{1,1}$$

- ▶ Uniformisation:

$$\mathcal{E}^{an} = SL_2(\mathbb{Z}) \ltimes \mathbb{Z}^2 \backslash \mathbb{H} \times \mathbb{C}$$

A point in $\mathcal{M}_{1,2}^{an}$ is determined by two coordinates $(\tau, z) \in \mathbb{H} \times \mathbb{C}$, where τ is 'modular' (horizontal) and z is 'elliptic' (vertical).



universal family elliptic curves punctured at the identity

- ▶ **Calaque–Enriquez–Etingof '09:** universal elliptic KZB equation

- ▶ **Calaque–Enriquez–Etingof '09:** universal elliptic KZB equation
- ▶ Integrable holomorphic connection on $\mathcal{M}_{1,2}^{an}$

- ▶ **Calaque–Enriquez–Etingof '09:** universal elliptic KZB equation
- ▶ Integrable holomorphic connection on $\mathcal{M}_{1,2}^{an}$
- ▶ Constructed first over the ‘infinite-rank vector bundle’ with fibre $\mathbb{L}(a, b)^\wedge \subset \mathbb{C}\langle\langle a, b \rangle\rangle$ over $\mathbb{H} \times \mathbb{C}$

- ▶ **Calaque–Enriquez–Etingof '09:** universal elliptic KZB equation
- ▶ Integrable holomorphic connection on $\mathcal{M}_{1,2}^{an}$
- ▶ Constructed first over the 'infinite-rank vector bundle' with fibre $\mathbb{L}(a, b)^\wedge \subset \mathbb{C}\langle\langle a, b \rangle\rangle$ over $\mathbb{H} \times \mathbb{C}$
- ▶ Explicit formula:

$$\nabla_{KZB} = d - dz \otimes ad_a F_\tau(z, ad_a)b - \frac{d\tau}{2\pi i} \otimes (ad_a F'_\tau(z, ad_a)b + D_\tau)$$

where $F'_\tau(z, x) = \frac{\partial}{\partial x} F_\tau(z, x) + \frac{1}{x^2}$, and

$$D_\tau = b \frac{\partial}{\partial a} + \frac{1}{2} \sum_{n \geq 2} (2n-1) G_{2n}(\tau) \sum_{\substack{j+k=2n-1 \\ j, k > 0}} [(-ad_a)^j b, (ad_a)^k b] \frac{\partial}{\partial b}.$$

and

$$F_\tau(z, x) = \frac{\theta'_\tau(0)\theta_\tau(z+x)}{\theta_\tau(z)\theta_\tau(x)}$$

There are also 'higher level versions' (punctures at torsion points)
by **Calaque–Gonzalez** '20.

There are also 'higher level versions' (punctures at torsion points) by **Calaque–Gonzalez** '20.

- ▶ **Problem:** how to study these equations purely within algebraic geometry?
 - ▶ Are they algebraic?
 - ▶ Explicit algebraic formulae?
 - ▶ Field of definition?

There are also 'higher level versions' (punctures at torsion points) by **Calaque–Gonzalez** '20.

- ▶ **Problem**: how to study these equations purely within algebraic geometry?
 - ▶ Are they algebraic?
 - ▶ Explicit algebraic formulae?
 - ▶ Field of definition?
- ▶ Motivations come from the theory of (arithmetic) **periods and motives** corresponding to **fundamental groups**.

There are also 'higher level versions' (punctures at torsion points) by **Calaque–Gonzalez** '20.

- ▶ **Problem**: how to study these equations purely within algebraic geometry?
 - ▶ Are they algebraic?
 - ▶ Explicit algebraic formulae?
 - ▶ Field of definition?
- ▶ Motivations come from the theory of (arithmetic) **periods and motives** corresponding to **fundamental groups**.
- ▶ Our approach with **Nils Matthes**: use **universal vector extensions of elliptic curves** (cf. Deligne).

There are also 'higher level versions' (punctures at torsion points) by **Calaque–Gonzalez** '20.

- ▶ **Problem**: how to study these equations purely within algebraic geometry?
 - ▶ Are they algebraic?
 - ▶ Explicit algebraic formulae?
 - ▶ Field of definition?
- ▶ Motivations come from the theory of (arithmetic) **periods and motives** corresponding to **fundamental groups**.
- ▶ Our approach with **Nils Matthes**: use **universal vector extensions of elliptic curves** (cf. Deligne).

Let X be a smooth variety over k , with $\text{char}(k) = 0$.

Let X be a smooth variety over k , with $\text{char}(k) = 0$.

- Vector bundle with integrable connection (algebraic):

Let X be a smooth variety over k , with $\text{char}(k) = 0$.

- **Vector bundle with integrable connection** (algebraic):
 (\mathcal{V}, ∇) , where \mathcal{V} is a vector bundle over X and

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/k}^1 \otimes \mathcal{V}$$

with vanishing curvature.

Let X be a smooth variety over k , with $\text{char}(k) = 0$.

- ▶ **Vector bundle with integrable connection** (algebraic):
 (\mathcal{V}, ∇) , where \mathcal{V} is a vector bundle over X and

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/k}^1 \otimes \mathcal{V}$$

with vanishing curvature.

- ▶ Locally:

$$\mathcal{V} = \mathcal{O}_X \otimes V, \quad \nabla = d + \omega, \quad d\omega + \omega \wedge \omega = 0$$

where $\omega \in \Omega^1(X) \otimes \text{End}(V)$.

Let X be a smooth variety over k , with $\text{char}(k) = 0$.

- ▶ **Vector bundle with integrable connection** (algebraic):
 (\mathcal{V}, ∇) , where \mathcal{V} is a vector bundle over X and

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/k}^1 \otimes \mathcal{V}$$

with vanishing curvature.

- ▶ Locally:

$$\mathcal{V} = \mathcal{O}_X \otimes V, \quad \nabla = d + \omega, \quad d\omega + \omega \wedge \omega = 0$$

where $\omega \in \Omega^1(X) \otimes \text{End}(V)$.

- ▶ (\mathcal{V}, ∇) is **unipotent** if there is a filtration

$$0 = \mathcal{V}_0 \subset \mathcal{V}_1 \subset \cdots \subset \mathcal{V}_n = \mathcal{V}$$

such that $\nabla \mathcal{V}_i \subset \Omega_{X/k}^1 \otimes \mathcal{V}_{i-1}$ and $(\mathcal{V}_i/\mathcal{V}_{i-1}, \nabla) \cong (\mathcal{O}_X, d)$.

Let X be a smooth variety over k , with $\text{char}(k) = 0$.

- **Vector bundle with integrable connection** (algebraic):
 (\mathcal{V}, ∇) , where \mathcal{V} is a vector bundle over X and

$$\nabla : \mathcal{V} \rightarrow \Omega_{X/k}^1 \otimes \mathcal{V}$$

with vanishing curvature.

- Locally:

$$\mathcal{V} = \mathcal{O}_X \otimes V, \quad \nabla = d + \omega, \quad d\omega + \omega \wedge \omega = 0$$

where $\omega \in \Omega^1(X) \otimes \text{End}(V)$.

- (\mathcal{V}, ∇) is **unipotent** if there is a filtration

$$0 = \mathcal{V}_0 \subset \mathcal{V}_1 \subset \cdots \subset \mathcal{V}_n = \mathcal{V}$$

such that $\nabla \mathcal{V}_i \subset \Omega_{X/k}^1 \otimes \mathcal{V}_{i-1}$ and $(\mathcal{V}_i/\mathcal{V}_{i-1}, \nabla) \cong (\mathcal{O}_X, d)$.

Problem: classify (and describe) unipotent integrable vector bundles with connection on X .

Example

Let $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$, and

$$\mathcal{V} = \mathcal{O}_X^{\otimes 3}, \quad \nabla = d - \begin{pmatrix} 0 & \frac{dz}{z} & 0 \\ 0 & 0 & \frac{dz}{1-z} \\ 0 & 0 & 0 \end{pmatrix}$$

Example

Let $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$, and

$$\mathcal{V} = \mathcal{O}_X^{\otimes 3}, \quad \nabla = d - \begin{pmatrix} 0 & \frac{dz}{z} & 0 \\ 0 & 0 & \frac{dz}{1-z} \\ 0 & 0 & 0 \end{pmatrix}$$

Fundamental solution of $\nabla = 0$:

$$\begin{pmatrix} 1 & \log(z) & -Li_2(z) \\ 0 & 1 & \log(1-z) \\ 0 & 0 & 1 \end{pmatrix}$$

where

$$Li_2(z) = \sum_{n=0}^{\infty} \frac{z^n}{n^2} = \int_0^z \int_0^y \frac{dx}{x-1} \frac{dy}{y}.$$

Theorem ('well-known...')

Every unipotent vector bundle with integrable connection on $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$ is canonically isomorphic to

$$(\mathcal{O}_X \otimes V, d - \frac{dz}{z} \otimes X_0 - \frac{dz}{1-z} \otimes X_1)$$

where V is a finite-dimensional vector space and $X_0, X_1 \in \text{End}(V)$ are simultaneously strictly upper triangularizable.

Key properties:

$$\begin{cases} H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}) = k \\ H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}) = \text{Ext}^1(\mathcal{O}_{\mathbb{P}^1}, \mathcal{O}_{\mathbb{P}^1}) = 0. \end{cases}$$

“Deligne's good conditions”

KZ connection on $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$:

$$\mathcal{V}_{KZ} = \mathcal{O}_X \hat{\otimes} k \langle\langle x_0, x_1 \rangle\rangle, \quad \nabla_{KZ} = d - \frac{dz}{z} \otimes x_0 - \frac{dz}{1-z} \otimes x_1$$

Universal property: given $b \in X$, for every unipotent (\mathcal{V}, ∇) with $v \in \mathcal{V}_b$, there is a unique $(\mathcal{V}_{KZ}, \nabla_{KZ}) \rightarrow (\mathcal{V}, \nabla)$ such that $1 \mapsto v$.

Questions:

- ▶ Something similar for $X = E \setminus O$?
- ▶ What about moduli $\mathcal{M}_{1,1}$? Extend to $\mathcal{M}_{1,2} = \mathcal{E} \setminus O$?

KZ connection on $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$:

$$\mathcal{V}_{KZ} = \mathcal{O}_X \hat{\otimes} k \langle\langle x_0, x_1 \rangle\rangle, \quad \nabla_{KZ} = d - \frac{dz}{z} \otimes x_0 - \frac{dz}{1-z} \otimes x_1$$

Universal property: given $b \in X$, for every unipotent (\mathcal{V}, ∇) with $v \in \mathcal{V}_b$, there is a unique $(\mathcal{V}_{KZ}, \nabla_{KZ}) \rightarrow (\mathcal{V}, \nabla)$ such that $1 \mapsto v$.

Questions:

- ▶ Something similar for $X = E \setminus O$?
- ▶ What about moduli $\mathcal{M}_{1,1}$? Extend to $\mathcal{M}_{1,2} = \mathcal{E} \setminus O$?

Note: $\mathbb{P}^1 \setminus \{0, 1, \infty\} = \mathcal{M}_{0,4}$

KZ connection on $X = \mathbb{P}^1 \setminus \{0, 1, \infty\}$:

$$\mathcal{V}_{KZ} = \mathcal{O}_X \hat{\otimes} k \langle\langle x_0, x_1 \rangle\rangle, \quad \nabla_{KZ} = d - \frac{dz}{z} \otimes x_0 - \frac{dz}{1-z} \otimes x_1$$

Universal property: given $b \in X$, for every unipotent (\mathcal{V}, ∇) with $v \in \mathcal{V}_b$, there is a unique $(\mathcal{V}_{KZ}, \nabla_{KZ}) \rightarrow (\mathcal{V}, \nabla)$ such that $1 \mapsto v$.

Questions:

- ▶ Something similar for $X = E \setminus O$?
- ▶ What about moduli $\mathcal{M}_{1,1}$? Extend to $\mathcal{M}_{1,2} = \mathcal{E} \setminus O$?

Note: $\mathbb{P}^1 \setminus \{0, 1, \infty\} = \mathcal{M}_{0,4}$

Elliptic KZB connection for $k = \mathbb{C}$ (**Levin–Racinet** '07)

Elliptic KZB connection for $k = \mathbb{C}$ (Levin–Racinet '07)

- Uniformisation $p : \mathbb{C} \rightarrow E^{an}$ inducing $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}) \cong E^{an}$

Elliptic KZB connection for $k = \mathbb{C}$ (Levin–Racinet '07)

- ▶ Uniformisation $p : \mathbb{C} \rightarrow E^{an}$ inducing $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}) \cong E^{an}$
- ▶ $(\mathcal{V}_E, \nabla_E)$ over $E \setminus O$ such that

$$p^*\mathcal{V}_E \cong \mathcal{O} \hat{\otimes} \mathbb{C} \langle\langle a, b \rangle\rangle$$

and

$$\begin{aligned} p^*\nabla_E &= d - dz \otimes ad_a F_\tau(z, ad_a)b \\ &= d - dz \otimes b - \frac{\theta'_\tau(z)}{\theta_\tau(z)} dz \otimes [a, b] - \dots \end{aligned}$$

Elliptic KZB connection for $k = \mathbb{C}$ (Levin–Racinet '07)

- ▶ Uniformisation $p : \mathbb{C} \rightarrow E^{an}$ inducing $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}) \cong E^{an}$
- ▶ $(\mathcal{V}_E, \nabla_E)$ over $E \setminus O$ such that

$$p^*\mathcal{V}_E \cong \mathcal{O} \hat{\otimes} \mathbb{C} \langle\langle a, b \rangle\rangle$$

and

$$\begin{aligned} p^*\nabla_E &= d - dz \otimes ad_a F_\tau(z, ad_a)b \\ &= d - dz \otimes b - \frac{\theta'_\tau(z)}{\theta_\tau(z)} dz \otimes [a, b] - \dots \end{aligned}$$

- ▶ Analytic formulae (not algebraic). What are the genus 1 analogues of $\frac{dz}{z}$ and $\frac{dz}{1-z}$?

Elliptic KZB connection for $k = \mathbb{C}$ (Levin–Racinet '07)

- ▶ Uniformisation $p : \mathbb{C} \rightarrow E^{an}$ inducing $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}) \cong E^{an}$
- ▶ $(\mathcal{V}_E, \nabla_E)$ over $E \setminus O$ such that

$$p^*\mathcal{V}_E \cong \mathcal{O} \hat{\otimes} \mathbb{C} \langle\langle a, b \rangle\rangle$$

and

$$\begin{aligned} p^*\nabla_E &= d - dz \otimes ad_a F_\tau(z, ad_a)b \\ &= d - dz \otimes b - \frac{\theta'_\tau(z)}{\theta_\tau(z)} dz \otimes [a, b] - \dots \end{aligned}$$

- ▶ Analytic formulae (not algebraic). What are the genus 1 analogues of $\frac{dz}{z}$ and $\frac{dz}{1-z}$?
- ▶ Note: $H^1(E, \mathcal{O}_E) \neq 0$. Deligne's good conditions are *not* satisfied.

Deligne '15: “use the universal vector extension E^{\natural} of E ”

Deligne '15: “use the **universal vector extension** E^\natural of E ”

- ▶ E^\natural is a smooth commutative algebraic group of dimension 2 (not affine, neither projective).

Deligne '15: “use the **universal vector extension** E^\natural of E ”

- ▶ E^\natural is a smooth commutative algebraic group of dimension 2 (not affine, neither projective).
- ▶ Extension of commutative algebraic groups:

$$0 \rightarrow \mathbb{G}_a \rightarrow E^\natural \xrightarrow{\pi} E \rightarrow 0$$

Deligne '15: “use the **universal vector extension** E^\natural of E ”

- ▶ E^\natural is a smooth commutative algebraic group of dimension 2 (not affine, neither projective).
- ▶ Extension of commutative algebraic groups:

$$0 \rightarrow \mathbb{G}_a \rightarrow E^\natural \xrightarrow{\pi} E \rightarrow 0$$

- ▶ Moduli space of line bundles with connection (\mathcal{L}, ∇) on E

Deligne '15: “use the **universal vector extension** E^\natural of E ”

- ▶ E^\natural is a smooth commutative algebraic group of dimension 2 (not affine, neither projective).
- ▶ Extension of commutative algebraic groups:

$$0 \rightarrow \mathbb{G}_a \rightarrow E^\natural \xrightarrow{\pi} E \rightarrow 0$$

- ▶ Moduli space of line bundles with connection (\mathcal{L}, ∇) on E
- ▶ If $\text{char}(k) = 0$, then

$$\begin{cases} H^0(E^\natural, \mathcal{O}_{E^\natural}) \cong k \\ H^1(E^\natural, \mathcal{O}_{E^\natural}) = 0 \end{cases}$$

Deligne's good conditions are satisfied!

Theorem (F.–Matthes)

If $\text{char}(k) = 0$, then

$$\Gamma(E^{\natural}, \Omega^1(\log \pi^{-1} O)) = \underbrace{k\nu \oplus k\omega^{(0)}}_{\Gamma(E^{\natural}, \Omega^1)} \oplus k\omega^{(1)} \oplus \dots$$

where $\omega^{(n)}$ are uniquely determined by $\omega^{(0)} \in \pi^* \Gamma(E, \Omega^1)$ and

- ▶ $d\omega^{(n)} = \nu \wedge \omega^{(n-1)}$
- ▶ $\omega^{(n)} \wedge \omega^{(0)} = 0$
- ▶ $\text{Res}(\omega^{(n)}) = t^n / (n-1)!$

Theorem (F.–Matthes)

The pullback of $(\mathcal{V}_E, \nabla_E)$ to E^\natural is given by

$$\pi^*\mathcal{V}_E \cong \mathcal{O} \hat{\otimes} k \langle\langle a, b \rangle\rangle, \quad \pi^*\nabla_E = d - \nu \otimes a - \sum_{n \geq 0} \omega^{(n)} \otimes ad_a^n b.$$

Theorem (F.–Matthes)

The pullback of $(\mathcal{V}_E, \nabla_E)$ to E^{\natural} is given by

$$\pi^* \mathcal{V}_E \cong \mathcal{O} \hat{\otimes} k \langle\langle a, b \rangle\rangle, \quad \pi^* \nabla_E = d - \nu \otimes a - \sum_{n \geq 0} \omega^{(n)} \otimes ad_a^n b.$$


Note: if $k = \mathbb{C}$ and $E^{an} \cong \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$, then

$$E^{\natural, an} \cong \mathbb{C}^2 / L_\tau, \quad L_\tau = \{(m + n\tau, n) : m, n \in \mathbb{Z}\}.$$

With coordinates (z, w) on \mathbb{C}^2 :

$$\nu = dw, \quad \omega^{(0)} = dz, \quad \omega^{(1)} = \left(\frac{\theta'_\tau(z)}{\theta_\tau(z)} + 2\pi iw \right) dz, \quad \dots$$

What about the universal KZB?



$$\nabla_{KZB} = d - dz \otimes ad_a F_\tau(z, ad_a) b - \frac{d\tau}{2\pi i} \otimes (ad_a F'_\tau(z, ad_a) b + D_\tau)$$

Universal vector extensions **behave very well in families:**


Universal vector extensions **behave very well in families**:

- ▶ Given a family of elliptic curves $\mathcal{E} \rightarrow S$, get a family of universal vector extensions $f : \mathcal{E}^\natural \rightarrow S$

Universal vector extensions **behave very well in families**:

- ▶ Given a family of elliptic curves $\mathcal{E} \rightarrow S$, get a family of universal vector extensions $f : \mathcal{E}^\natural \rightarrow S$
- ▶ *Canonical* horizontal foliation (isomonodromy of (\mathcal{L}, ∇))


$$0 \longrightarrow T_{\mathcal{E}^\natural/S} \longrightarrow T_{\mathcal{E}^\natural} \xrightarrow{\quad \leftarrow \quad} f^* T_S \longrightarrow 0$$



Universal vector extensions **behave very well in families**:

- ▶ Given a family of elliptic curves $\mathcal{E} \rightarrow S$, get a family of universal vector extensions $f : \mathcal{E}^\natural \rightarrow S$
- ▶ *Canonical* horizontal foliation (isomonodromy of (\mathcal{L}, ∇))

$$0 \longrightarrow T_{\mathcal{E}^\natural/S} \longrightarrow T_{\mathcal{E}^\natural} \xrightarrow{\quad \leftarrow \quad} f^* T_S \longrightarrow 0$$



- ▶ **Laumon '96, Coleman '98**: If $\text{char}(S) = 0$,

$$f_* \mathcal{O}_{\mathcal{E}^\natural} \cong \mathcal{O}_S, \quad R^1 f_* \mathcal{O}_{\mathcal{E}^\natural} = 0.$$

Theorem (F.–Matthes)

The universal KZB connection in level N is algebraic and defined over the number field $\mathbb{Q}(\mu_N)$.

Theorem (F.–Matthes)


The universal KZB connection in level N is algebraic and defined over the number field $\mathbb{Q}(\mu_N)$.

- ▶ **Luo** '19: $N = 1$ by different methods.

Theorem (F.–Matthes)

The universal KZB connection in level N is algebraic and defined over the number field $\mathbb{Q}(\mu_N)$.


- ▶ **Luo** '19: $N = 1$ by different methods.
- ▶ Our construction relies on a theorem which provides **canonical lifts of relative logarithmic differential forms**

$$0 \longrightarrow \Omega_S^1 \longrightarrow f_* \Omega_{\mathcal{E}^{\natural}}^1(\log \pi^{-1} O) \longrightarrow f_* \Omega_{\mathcal{E}^{\natural}/S}^1 \longrightarrow 0$$


Theorem (F.–Matthes)

The universal KZB connection in level N is algebraic and defined over the number field $\mathbb{Q}(\mu_N)$.

- ▶ **Luo '19:** $N = 1$ by different methods.
- ▶ Our construction relies on a theorem which provides **canonical lifts of relative logarithmic differential forms**

$$0 \longrightarrow \Omega_S^1 \longrightarrow f_* \Omega_{\mathcal{E}^\natural}^1(\log \pi^{-1} O) \longrightarrow f_* \Omega_{\mathcal{E}^\natural/S}^1 \longrightarrow 0$$


- ▶ Example:

$$\begin{aligned} \tilde{\omega}^{(1)} = & \left(\frac{\theta'_\tau(z)}{\theta_\tau(z)} + 2\pi i w \right) dz \\ & + \frac{1}{2\pi i} \left(\frac{1}{2} \frac{\theta''_\tau(z)}{\theta_\tau(z)} - \frac{1}{6} \frac{\theta'''_\tau(0)}{\theta'_\tau(0)} - \frac{(2\pi i w)^2}{2} \right) d\tau. \end{aligned}$$

Thank you!