

Epidemics in Networks

SIR: Final Size and Dynamics

Joel C. Miller & Tom Hladish

13–15 July 2015

Introduction

Final Size

Dynamic equations

References

Introduction

We now consider the final size and the temporal dynamics of SIR epidemics in static configuration model networks.

Introduction

Final Size

Dynamic equations

References


Applying percolation results

- ▶ I claimed that for unclustered networks, final size depends on distribution of susceptibility.
- ▶ For both Reed-Frost and continuous time epidemics, susceptibility is uniform.
- ▶ So they should have the same final size.
- ▶ Also, we have proven that for Reed–Frost epidemics $\mathcal{P} = \mathcal{A}$. We'll calculate \mathcal{A} anyways, using a very similar method.
- ▶ To introduce something new, assume that at $t = 0$ some nodes are chosen to be infected so that $S(k, 0)$ is the probability a degree k individual is susceptible [1].

Setting up size calculations

- ▶ Assuming an epidemic occurs, the probability a random node u is infected equals the proportion of the population that is infected.
- ▶ The probability a random node u is infected is unaltered if we prevent u from transmitting to its partners.
- ▶ So consider a random test node u which is prevented from transmitting to its partners, we want to know the probability u is infected if an epidemic occurs.

Reed Frost final size in configuration model networks



$$\Theta = P(v \text{ did not transmit to } u)$$


Probability a random degree- k test individual is susceptible is

$$S = \sum_k P(k) S(k, 0) \Theta^k = \hat{\psi}(\Theta)$$

where

$$\hat{\psi}(x) = \sum_k P(k) S(k, 0) x^k$$

Finding \ominus



Probability a random degree- k partner susceptible is

$$\phi_S = \sum_k \frac{kP(k)}{\langle K \rangle} S(k, 0) \Theta^{k-1} = \frac{\hat{\psi}'(\Theta)}{\langle K \rangle}$$

If p , then probability partner does not transmit to u is

$$\Theta = \phi_S + (1 - p)(1 - \phi_S) = 1 - p + p \frac{\hat{\psi}'(\Theta)}{\langle K \rangle}$$

Final Size

So


$$R = 1 - \hat{\psi}(\Theta)$$

where

$$\Theta = 1 - p + p \frac{\hat{\psi}'(\Theta)}{\langle K \rangle}$$

Contuuous time case

To look at the continuous time case, we use the same figure.



$$\Theta = P(v \text{ did not transmit to } u)$$

Here each Θ is uncorrelated with the other Θ s.

- ▶ Transmission to all partners of an individual v are not independent because they depend on the duration of v 's infection.
- ▶ However, transmissions from different partners to u are independent.
- ▶ So the identical calculation applies.

Introduction

Final Size

Dynamic equations

References

SIR dynamic equations

- ▶ We now derive the equations for the time dependent case, assuming continuous time.
- ▶ The derivation is almost identical to the final size derivation, except that we calculate the probability transmission happened by time t rather than by the end.
- ▶ This is not the only approach [2, 3, 4, 5], but the predictions of the models are identical (subject to some small caveats) [6].

Calculating Dynamics

The network structure alters the infection process (but not the recoveries)

Calculating Dynamics

The network structure alters the infection process (but not the recoveries)



$$I = 1 - S - R, \quad \dot{R} = \gamma I$$

We will switch to a partnership-based perspective to find $S(t)$.

Revisiting the test individual


- ▶ Consider a randomly chosen test individual u in the population.
- ▶ Disallow infection from u to its partners (allows independence assumption for partners).
- ▶ The probability u is Susceptible, Infected, or Recovered at time t is affected by the status of its partners.
- ▶ The fraction of the population that is susceptible $S(t)$ equals the probability u is susceptible.

$$S(t) = P(u \text{ is susceptible})$$

- ▶ Let v be a random partner of u .
- ▶ Define

$$\theta(t) = P(v \text{ not yet transmitted to } u)$$

Finding $S(t)$



$$\theta(t) = P(v \text{ not yet transmitted to } u)$$


Probability a random degree- k test individual still susceptible is

$$S(t) = \sum_k P(k) S(k, 0) \theta(t)^k = \hat{\psi}(\theta(t))$$


where

$$\hat{\psi}(x) = \sum_k S(k, 0) P(k) x^k$$

How does θ evolve?




How does θ evolve?




- $\theta = \phi_S + \phi_I + \phi_R$.
- $\dot{\theta} = -\beta\phi_I$.
- Our goal is to find ϕ_I in terms of θ .

Finding $\phi_R(t)$




Because derivatives are proportional, $\phi_R = \frac{\gamma}{\beta}(1 - \theta)$

Finding $\phi_S(t)$



Probability a random ~~degree k~~ partner still susceptible is

$$\phi_S(t) = \sum_k \frac{kP(k)}{\langle K \rangle} S(k, 0) \theta(t)^{k-1} = \frac{\hat{\psi}'(\theta)}{\langle K \rangle}$$




Since $\phi_I = \theta - \phi_S - \phi_R = \theta - \frac{\hat{\psi}'(\theta)}{\langle K \rangle} - \frac{\gamma}{\beta}(1 - \theta)$, we have

$$\dot{\theta} = -\beta\phi_I = -\beta\theta + \beta\frac{\hat{\psi}'(\theta)}{\langle K \rangle} + \gamma(1 - \theta)$$

Final System

We finally have

$$\begin{aligned}\dot{\theta} &= -\beta\theta + \beta\frac{\hat{\psi}'(\theta)}{\langle K \rangle} + \gamma(1 - \theta) \\ \dot{R} &= \gamma I \quad S = \hat{\psi}(\theta) \quad I = 1 - S - R\end{aligned}$$



Exercise

Derive the Reed–Frost version of this model [7]
 Set the probability of transmitting in a time step to be β .

A major caveat

- ▶ All of this assumes that the strength of partnerships is not degree-dependent. This is a dangerous assumption.
- ▶ I suspect most results from the high-impact papers on powerlaw degree distributions will disappear if we make more realistic assumptions.

Generalizing


Many generalizations are possible:

- ▶ Dynamic networks, other classes of random networks [8].
- ▶ SEIR, demographic groups, different partnership types, asymmetric transmission, ... [9].
- ▶ Competing diseases [10].
- ▶ Spread of ideas or behaviors [11].
- ▶ Including birth and death (ongoing)
- ▶ Degree-dependent transmission probabilities [12] (and ongoing work).

Other formulations


- ▶ Other variable choices exist, which lead to different equations.
- ▶ Subject to mild conditions the equations are all equivalent [6].
- ▶ The number of equations and compartments can vary substantially

Pairwise model flow diagrams




- ▶ The flow diagram underlying the basic pairwise model [2].
- ▶ We track individuals of each status and degree as well as partnerships between individuals of various statuses and degrees.
- ▶ Dashed lines denote transitions that rely on infection coming from a source outside the edge of interest.
- ▶ The triples $[A_{k'} S_k I]$ and $[I S_k A_{k'}]$ can be expressed in terms of the doubles and singles: $[A_{k'} S_k][S_k I]/[S_k]$

Reduced pairwise flow diagram




- ▶ The flow diagram for the reduced system of [4].
- ▶ The $[SS]$, $[SI]$, and $[SR]$ compartments correspond to the sum of the $[S_k S_{k'}]$, $[S_k I_{k'}]$ and $[S_k R_{k'}]$ compartments of the basic pairwise model.
- ▶ $\langle I \rangle = [SI]/([SS] + [SI] + [SR])$

Effective degree model flow diagram



- The flow diagram underlying the effective degree model of [3].
- We include just the fluxes involving the $x_{s,i}$ or $y_{s,i}$ compartments. Fluxes between other compartments exist but are not included.

Compact effective degree model flow diagram



- The flow diagram that underlies the model of [5].
- Only the fluxes into and out of y_j and x_j are included. Fluxes between other compartments exist, but are not included.
- An active edge is eliminated if the partner recovers or if it transmits infection in either direction. The quantity $\langle I \rangle$ represents the probability an active edge joins an individual with an infected partner.

Introduction

Final Size

Dynamic equations

References

References I

- [1] Joel C. Miller.
Epidemics on networks with large initial conditions or changing structure.
[PLoS ONE](#), 9(7):e101421, 2014.
- [2] K.T.D. Eames and M.J. Keeling.
Modeling dynamic and network heterogeneities in the spread of sexually transmitted diseases.
[Proceedings of the National Academy of Sciences](#), 99(20):13330–13335, 2002.
- [3] J. Lindquist, J. Ma, P. van den Driessche, and F.H. Willeboordse.
Effective degree network disease models.
[Journal of Mathematical Biology](#), 62(2):143–164, 2011.
- [4] Laurent Hébert-Dufresne, Oscar Patterson-Lomba, Georg M Goerg, and Benjamin M Althouse.
Pathogen mutation modeled by competition between site and bond percolation.
[Physical Review Letters](#), 110(10):108103, 2013.
- [5] F. Ball and P. Neal.
Network epidemic models with two levels of mixing.
[Mathematical Biosciences](#), 212(1):69–87, 2008.
- [6] Joel C. Miller and Istvan Z. Kiss.
Epidemic spread in networks: Existing methods and current challenges.
[Mathematical Modelling of Natural Phenomena](#), 2014.
- [7] Lucas Daniel Valdez, Pablo Alejandro Macri, and Lidia Adriana Braunstein.
Temporal percolation of the susceptible network in an epidemic spreading.
[PLoS ONE](#), 7(9):e44188, 2012.
- [8] Joel C. Miller, Anja C. Slim, and Erik M. Volz.
Edge-based compartmental modelling for infectious disease spread.
[Journal of the Royal Society Interface](#), 9(70):890–906, 2012.

References II

- [9] Joel C. Miller and Erik M. Volz.
Incorporating disease and population structure into models of SIR disease in contact networks.
[PLoS ONE](#), 8(8):e69162, 2013.
- [10] Joel C. Miller.
Cocirculation of infectious diseases on networks.
[Physical Review E](#), 87(6):060801, 2013.
- [11] Joel C. Miller.
Complex contagions and hybrid phase transitions in unclustered and clustered random networks.
[Under Review; arXiv preprint arXiv:1501.01585](#), 2015.
- [12] P Rattana, JC Miller, and IZ Kiss.
Pairwise and edge-based models of epidemic dynamics on correlated weighted networks.
[Mathematical modelling of natural phenomena](#), 9(02):58–81, 2014.