

# Epidemics in Networks

## SIS: Equilibrium Size and Dynamics

Joel C. Miller & Tom Hladish

13–15 July 2015

Introduction


SIS dynamics

Equilibrium

Other models

References

# Introduction



- For technical reasons it is difficult to handle SIS on configuration model networks.
- The standard solution is to turn to annealed networks.
- There are some subtle but important differences between epidemics in the two cases.

## History

- ▶ The model was introduced in 1980 [1] and used widely in mathematical biology [2, 3]: “Social heterogeneity”
- ▶ In 2001 it was used to study powerlaw degree distributions [4]: “Degree-based mean field”
- ▶ Many people use it as if it were correct for configuration model networks.
- ▶ The major result of [4] is altered if the network is static [5].

# What's the problem?

- ▶ If a high degree node is infected in a static network, it will infect many partners.
- ▶ When it recovers, these partners are likely to reinfect it.
- ▶ A local island of infection forms.
- ▶ These islands occasionally go extinct, and occasionally they spread disease to another high degree node, starting a new island.

Introduction

SIS dynamics

Equilibrium

Other models

References

## SIS dynamics

- ▶ Because partnerships are constantly changing, the probability a partner is infected is independent of the history of a node.
- ▶ Let  $\langle I \rangle$  denote the probability a partner is infected.
- ▶ Then

$$\langle I \rangle = \frac{\sum_k k P(k) I_k}{\sum_k k P(k)} = \frac{\sum_k k P(k) I_k}{\langle K \rangle}$$

Where  $I_k$  is the probability a degree  $k$  individual is infected.

- ▶ We get

$$\begin{aligned}\dot{S}_k &= -\beta k S_k \langle I \rangle + \gamma I_k \\ \dot{I}_k &= \beta k S_k \langle I \rangle - \gamma I_k\end{aligned}$$

Introduction

SIS dynamics

Equilibrium

Other models

References

# SIS equilibrium

At equilibrium infection balances recovery.

$$\begin{aligned}\beta k S_k \langle I \rangle &= \gamma I_k \\ \beta k (1 - I_k) \langle I \rangle &= \gamma I_k \\ \beta k \langle I \rangle &= (\gamma + \beta k \langle I \rangle) I_k \\ \Rightarrow I_k &= \frac{\beta k \langle I \rangle}{\gamma + \beta k \langle I \rangle}\end{aligned}$$

where

$$\langle I \rangle = \frac{\sum_k k P(k) I_k}{\langle K \rangle}$$

Introduction

SIS dynamics

Equilibrium

Other models


References

## Other models

I am aware of two other systems of equations that appear to do a reasonable job on configuration model networks


## Effective degree model

From [6]



$$G = \frac{\sum_{k=1}^M \sum_{j+l=k} j \beta I S_{jl}}{\sum_{k=1}^M \sum_{j+l=k} j S_{jl}}$$

$$H = \frac{\sum_{k=1}^M \sum_{j+l=k} \beta I^2 S_{jl}}{\sum_{k=1}^M \sum_{j+l=k} j I_{jl}}$$




An alternate model by [7] has just 5 equations:

$$\begin{aligned}\frac{d}{dt}[S] &= \gamma[I] - \beta[SI] \\ \frac{d}{dt}[I] &= \beta[SI] - \gamma[I] \\ \frac{d}{dt}[SI] &= \gamma([II] - [SI]) + \beta[SI]([SS] - [SI])Q - \beta[SI] \\ \frac{d}{dt}[SS] &= 2\gamma[SI] - 2\beta[SI][SS]Q \\ \frac{d}{dt}[II] &= -2\gamma[II] + 2\beta[SI]^2Q + 2\beta[SI]\end{aligned}$$

where

$$Q = \frac{1}{[S]n_S} \left( \frac{\langle K^2 \rangle (\langle K^2 \rangle - n_S \langle K \rangle) + \langle K^3 \rangle (n_S - \langle K \rangle)}{n_S (\langle K^2 \rangle - \langle K \rangle^2)} - 1 \right)$$

$$n_S = \frac{[SI] + [SS]}{[S]}$$



Introduction

SIS dynamics

Equilibrium

Other models

References



# References I

- [1] Annett Nold.  
**Heterogeneity in disease-transmission modeling.**  
Mathematical Biosciences, 52(3):227–240, 1980.
- [2] Roy M. Anderson and Robert M. May.  
**Infectious Diseases of Humans.**  
Oxford University Press, Oxford, 1991.
- [3] Herbert W Hethcote, James A Yorke, and Annett Nold.  
**Gonorrhea modeling: a comparison of control methods.**  
Mathematical Biosciences, 58(1):93–109, 1982.
- [4] Romualdo Pastor-Satorras and Alessandro Vespignani.  
**Epidemic spreading in scale-free networks.**  
Physical Review Letters, 86(14):3200–3203, Apr 2001.
- [5] S. Chatterjee and R. Durrett.  
**Contact processes on random graphs with power law degree distributions have critical value 0.**  
The Annals of Probability, 37(6):2332–2356, 2009.
- [6] J. Lindquist, J. Ma, P. van den Driessche, and F.H. Willeboordse.  
**Effective degree network disease models.**  
Journal of Mathematical Biology, 62(2):143–164, 2011.
- [7] Péter L Simon and Istvan Z Kiss.  
**Super compact pairwise model for sis epidemic on heterogeneous networks.**  
arXiv preprint arXiv:1503.01090, 2015.