

National Coverage Determination (NCD)

Counseling to Prevent Tobacco Use

210.4.1

Links in PDF documents are not guaranteed to work. To follow a web link, please use the MCD Website.

Tracking Information

Publication Number

100-3

Manual Section Number

210.4.1

Manual Section Title

Counseling to Prevent Tobacco Use

Version Number

1

Effective Date of this Version

08/25/2010

Ending Effective Date of this Version

09/26/2017

Implementation Date

01/03/2011

Description Information

Benefit Category

Additional Preventive Services

Please Note: This may not be an exhaustive list of all applicable Medicare benefit categories for this item or service.

Item/Service Description

A. General

Tobacco use remains the leading cause of preventable morbidity and mortality in the U.S. and is a major contributor to the nation's increasing medical costs. Despite the growing list of adverse health effects associated with smoking, more than 45 million U.S. adults continue to smoke and approximately 1,200 die prematurely each day from tobacco-related diseases. Annual smoking-attributable expenditures can be measured both in direct medical costs (\$96 billion) and in lost productivity (\$97 billion), but the results of national surveys have raised concerns that recent declines in smoking prevalence among U.S. adults may have come to an end. According to the U.S. Department of Health and Human Services (DHHS) Public Health Service (PHS) Clinical Practice Guideline on Treating Tobacco Use and Dependence (2008), 4.5 million adults over 65 years of age smoke cigarettes. Even smokers over age 65, however, can benefit greatly from abstinence, and older smokers who quit can reduce their risk of death from coronary heart disease, chronic obstructive lung disease and lung cancer, as well as decrease their risk of osteoporosis.

Medicare Part B (section 210.4 of the National Coverage Determination (NCD) Manual) already covers cessation counseling for individuals who use tobacco and have been diagnosed with a recognized tobacco-related disease or who exhibit symptoms consistent with tobacco-related disease. In November 2009, based upon authority to cover "additional preventive services" for Medicare beneficiaries if certain statutory requirements are met, the Centers for Medicare & Medicaid Services (CMS) initiated a new national coverage analysis to evaluate whether the existing evidence on counseling to prevent tobacco use is sufficient to extend national coverage for cessation counseling to those individuals who use tobacco but do not have signs or symptoms of tobacco-related disease. One of these statutory requirements is that the service be categorized as a grade A (strongly recommends) or grade B (recommends) rating by the US Preventive Services Task Force (USPSTF).

Indications and Limitations of Coverage

B. Nationally Covered Indications

Effective for claims with dates of service on or after August 25, 2010, CMS will cover tobacco cessation counseling for outpatient and hospitalized Medicare beneficiaries

1. Who use tobacco, regardless of whether they have signs or symptoms of tobacco-related disease;
2. Who are competent and alert at the time that counseling is provided; and,
3. Whose counseling is furnished by a qualified physician or other Medicare-recognized practitioner.

Intermediate and intensive smoking cessation counseling services will be covered under Medicare Part B when the above conditions of coverage are met, subject to frequency and other limitations. That is, similar to existing tobacco cessation counseling for symptomatic individuals, CMS will allow 2 individual tobacco cessation counseling attempts per 12-

month period. Each attempt may include a maximum of 4 intermediate OR intensive sessions, with a total benefit covering up to 8 sessions per 12-month period per Medicare beneficiary who uses tobacco. The practitioner and patient have the flexibility to choose between intermediate (more than 3 minutes but less than 10 minutes), or intensive (more than 10 minutes) cessation counseling sessions for each attempt.

C. Nationally Non-Covered Indications

Inpatient hospital stays with the principal diagnosis of tobacco use disorder are not reasonable and necessary for the effective delivery of tobacco cessation counseling services. Therefore, we will not cover tobacco cessation services if tobacco cessation is the primary reason for the patient's hospital stay.

D. Other

Section 4104 of the Affordable Care Act provided for a waiver of the Medicare coinsurance and Part B deductible requirements for this service effective on or after January 1, 2011. Until that time, this service will continue to be subject to the standard Medicare coinsurance and Part B deductible requirements.

(This NCD last reviewed August 2010.)

Claims Processing Instructions

[TN 2058 \(Medicare Claims Processing\)](#) 

[TN 3848 \(Medicare Claims Processing\)](#) 

Transmittal Information

Transmittal Number

126

Coverage Transmittal Link

<https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/downloads/R126NCD.pdf> 

Revision History

07/2017 - This Change Request (CR) constitutes a maintenance update of International Code of Diseases, Tenth Revision (ICD-10) conversions and other coding updates specific to National Coverage Determinations (NCDs). These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at:

<https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. (TN 1875) (CR10184)

06/2016 - Transmittal 1665, dated May 13, 2016, is being rescinded and replaced by Transmittal 1672 to: (1) 9631.1: Remove FISS responsibility and 1st sentence; (2) 9631.2: Remove additional procedure codes, including 0075T, 0076T; (3) 9631.4: Revise descriptor of dx L59.8; (4) 9631.6: Add deletion of dx C49.10, C65.9, remove deletion of dx C54.9, remove deletion of invalid dx C47.90; (5) 9631.8: Remove deletion of invalid dx C51.29, replace with deletion of dx C50.029; (6) 9631.9: Add deletion of 0V504ZZ, 0V500ZZ, override capability, and contractor discretion verbiage. All other information remains the same. (TN 1672) (CR9631)

05/2016 - This change request (CR) is the 7th maintenance update of ICD-10 conversions and other coding updates specific to national coverage determinations (NCDs). The majority of the NCDs included are a result of feedback received from previous ICD-10 NCD CR7818, CR8109, CR8197, CR8691, CR9087, CR9252, and CR9540. Some are the result of revisions required to other NCD-related CRs released separately. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent, quarterly releases as needed. No policy-related changes are included with these updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. (TN 1665) (CR9631)

03/2013 - CMS translated the information for this policy from ICD-9-CM/PCS to ICD-10-CM/PCS according to HIPAA standard medical data code set requirements and updated any necessary and related coding infrastructure. These updates do not expand, restrict, or alter existing coverage policy. Implementation date: 10/07/2013 Effective date: 10/1/2015. (TN 1199) (TN 1199) (CR 8197)

09/2010 - These individuals who do not have signs or symptoms of tobacco-related disease will be covered under Medicare Part B when the above conditions of coverage are met, subject to certain frequency and other limitations. Effective date 08/25/2010 Implementation date 01/03/2011 (TN 126) (CR 7133)

09/2010 - Effective for claims with dates of service on and after August 25, 2010, CMS will cover tobacco cessation counseling for outpatient and hospitalized Medicare beneficiaries:

1. Who use tobacco, regardless of whether they have signs or symptoms of tobacco-related disease;
2. Who are competent and alert at the time that counseling is provided; and

3. Whose counseling is furnished by a qualified physician or other Medicare-recognized practitioner.

National Coverage Analyses (NCAs)

This NCD has been or is currently being reviewed under the National Coverage Determination process. The following are existing associations with NCAs, from the National Coverage Analyses database.

- [Original Consideration for Counseling to Prevent Tobacco Use \(CAG-00420N\)](#) 

Additional Information

Other Versions

Title	Version	Effective Between	
Counseling to Prevent Tobacco Use	2	09/26/2017 - N/A	View
Counseling to Prevent Tobacco Use	1	08/25/2010 - 09/26/2017	You are here