

National Coverage Determination (NCD)

Erythropoiesis Stimulating Agents (ESAs) in Cancer and Related Neoplastic Conditions

110.21

Links in PDF documents are not guaranteed to work. To follow a web link, please use the MCD Website.

Tracking Information

Publication Number

100-3

Manual Section Number

110.21

Manual Section Title

Erythropoiesis Stimulating Agents (ESAs) in Cancer and Related Neoplastic Conditions

Version Number**Effective Date of this Version**

07/30/2007

Implementation Date

04/07/2008

Description Information

Benefit Category

Drugs and Biologicals

Please Note: This may not be an exhaustive list of all applicable Medicare benefit categories for this item or service.

Item/Service Description**A. General**

Erythropoiesis stimulating agents (ESAs) stimulate the bone marrow to make more red blood cells and are United States Food and Drug Administration (FDA) approved for use in

reducing the need for blood transfusion in patients with specific clinical indications. The FDA has issued alerts and warnings for ESAs administered for a number of clinical conditions, including cancer. Published studies report a higher risk of serious and life-threatening events associated with oncologic uses of ESAs.

Indications and Limitations of Coverage

B. Nationally Covered Indications

ESA treatment for the anemia secondary to myelosuppressive anticancer chemotherapy in solid tumors, multiple myeloma, lymphoma, and lymphocytic leukemia is only reasonable and necessary under the following specified conditions:

- The hemoglobin level immediately prior to initiation or maintenance of ESA treatment is < 10 g/dL (or the hematocrit is < 30%).
- The starting dose for ESA treatment is the recommended FDA label starting dose, no more than 150 U/kg/3 times weekly for epoetin and 2.25 mcg/kg/1 time weekly for darbepoetin alpha. Equivalent doses may be given over other approved time periods.
- Maintenance of ESA therapy is the starting dose if the hemoglobin level remains below 10g/dL (or hematocrit is < 30%) 4 weeks after initiation of therapy and the rise in hemoglobin is $\geq 1\text{g/dL}$ (hematocrit $\geq 3\%$);
- For patients whose hemoglobin rises < 1g/dl (hematocrit rise < 3%) compared to pretreatment baseline over 4 weeks of treatment and whose hemoglobin level remains < 10g/dL after the 4 weeks of treatment (or the hematocrit is < 30%), the recommended FDA label starting dose may be increased once by 25%. Continued use of the drug is not reasonable and necessary if the hemoglobin rises < 1g/dl (hematocrit rise < 3%) compared to pretreatment baseline by 8 weeks of treatment.
- Continued administration of the drug is not reasonable and necessary if there is a rapid rise in hemoglobin > 1g/dl (hematocrit > 3%) over 2 weeks of treatment unless the hemoglobin remains below or subsequently falls to < 10g/dL (or the hematocrit is < 30%). Continuation and reinstitution of ESA therapy must include a dose reduction of 25% from the previously administered dose.
- ESA treatment duration for each course of chemotherapy includes the 8 weeks following the final dose of myelosuppressive chemotherapy in a chemotherapy regimen.

C. Nationally Non-Covered Indications

ESA treatment is not reasonable and necessary for beneficiaries with certain clinical conditions, either because of a deleterious effect of the ESA on their underlying disease or because the underlying disease increases their risk of adverse effects related to ESA use. These conditions include:

- Any anemia in cancer or cancer treatment patients due to folate deficiency, B-12 deficiency, iron deficiency, hemolysis, bleeding, or bone marrow fibrosis;
- The anemia associated with the treatment of acute and chronic myelogenous leukemias (CML, AML), or erythroid cancers;
- The anemia of cancer not related to cancer treatment;
- Any anemia associated only with radiotherapy;
- Prophylactic use to prevent chemotherapy-induced anemia;
- Prophylactic use to reduce tumor hypoxia;
- Patients with erythropoietin-type resistance due to neutralizing antibodies; and
- Anemia due to cancer treatment if patients have uncontrolled hypertension.

D. Other

Local Medicare Administrative Contractors may continue to make reasonable and necessary determinations on all other uses of ESAs not specified in this National Coverage Determination.

See the [Medicare Benefit Policy Manual](#), chapter 11, section 90 and chapter 15, section 50.5.2 for coverage of ESAs for end-stage renal disease-related anemia.

Claims Processing Instructions

TN 1413 (Medicare Claims Processing) [↗](#)

Transmittal Information

Transmittal Number

80

Coverage Transmittal Link

<https://www.cms.gov/regulations-and-guidance/guidance/transmittals/downloads/dwnlds/r80ncdpdf> [↗](#)

Revision History

10/2024 - The purpose of the Change Request (CR) is to provide a quarterly maintenance update of ICD-10 coding conversions and other coding updates specific to National Coverage Determinations (NCDs). No policy is being changed as a result of these updates. ([TN 12903](#) [↗](#)) (CR13818)

01/2021 - Transmittal 10515, dated December 10, 2020, is being rescinded and replaced by Transmittal 10566, dated, January 14, 2021 to remove FISS Reason Codes (RCs) 59041, 59042, 59209, and 59210 from the spreadsheet for NCD 160.18. All other information remains the same. ([TN 10566](#) [↗](#)) (CR12027)

12/2020 - This Change Request (CR) constitutes a maintenance update of ICD-10 conversions and other coding updates specific to NCDs. These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, longstanding NCD process. (TN 10515) (CR12027)

10/2020 - This Change Request (CR) constitutes a maintenance update of ICD-10 conversions and other coding updates specific to NCDs. These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, longstanding NCD process. (TN 10432) (CR12027)

11/2018 - This Change Request (CR) constitutes a maintenance update of ICD-10 conversions and other coding updates specific to NCDs. These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. (TN 2202) (CR11005)

11/2018 - This Change Request (CR) constitutes a maintenance update of ICD-10 conversions and other coding updates specific to NCDs. These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other

CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. ([TN 2200](#)) (CR10859)

09/2018 - This Change Request (CR) constitutes a maintenance update of ICD-10 conversions and other coding updates specific to NCDs. These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. ([TN 2138](#)) (CR10859)

08/2018 - This Change Request (CR) constitutes a maintenance update of ICD-10 conversions and other coding updates specific to NCDs. These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. ([TN 2122](#)) (CR10859)

02/2018 - Transmittal 2033, dated February 16, 2018, is being rescinded and replaced by Transmittal 2039, dated, February 28, 2018 to correct instructions in business requirement 7, NCD210.3, Colorectal Cancer Screening, and its accompanying spreadsheet. All other information remains the same. ([TN 2039](#)) (CR10473)

02/2018 - This Change Request (CR) constitutes a maintenance update of International Code of Diseases, Tenth Revision (ICD-10) conversions and other coding updates specific to National Coverage Determinations (NCDs). These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other

CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. ([TN 2033](#)) (CR10473)

01/2018 - This Change Request (CR) constitutes a maintenance update of International Code of Diseases, Tenth Revision (ICD-10) conversions and other coding updates specific to National Coverage Determinations (NCDs). These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received.

Previous NCD coding changes appear in ICD-10 quarterly updates that can be found at: <https://www.cms.gov/Medicare/Coverage/CoverageGenInfo/ICD10.html>, along with other CRs implementing new policy NCDs. Edits to ICD-10 and other coding updates specific to NCDs will be included in subsequent quarterly releases and individual CRs as appropriate. No policy-related changes are included with the ICD-10 quarterly updates. Any policy-related changes to NCDs continue to be implemented via the current, long-standing NCD process. ([TN 2005](#)) (CR10318)

11/2017 - This Change Request (CR) constitutes a maintenance update of International Code of Diseases, Tenth Revision (ICD-10) conversions and other coding updates specific to National Coverage Determinations (NCDs). These NCD coding changes are the result of newly available codes, coding revisions to NCDs released separately, or coding feedback received. ([TN 1975](#)) (CR10318)

12/2015 - This change request (CR) is the 3rd maintenance update of ICD-10 conversions/updates specific to national coverage determinations (NCDs). The majority of the NCDs included are a result of feedback received from previous ICD-10 NCD CR7818, CR8109, CR8197, CR8691, & CR9087. Some are the result of revisions required to other NCD-related CRs released separately that included ICD-10 coding. Implementation date: 01/04/2016 Effective date: 10/1/2015. ([TN 1580](#)) (CR9252)

08/2015 - This change request (CR) is the 3rd maintenance update of ICD-10 conversions/updates specific to national coverage determinations (NCDs). The majority of the NCDs included are a result of feedback received from previous ICD-10 NCD CR7818, CR8109, CR8197, CR8691, & CR9087. Some are the result of revisions required to other NCD-related CRs released separately that included ICD-10 coding. These updates do not expand, restrict, or alter existing coverage policy. Implementation date: 01/04/2016 Effective date: 10/1/2015. ([TN 1537](#)) (CR 9252)

09/2012 - CMS translated the information for this policy from ICD-9-CM/PCS to ICD-10-CM/PCS according to HIPAA standard medical data code set requirements and updated any

necessary and related coding infrastructure. These updates do not expand, restrict, or alter existing coverage policy. Implementation date: 01/07/2013 Effective date: 10/1/2015. [\(TN 1122\)](#) [\(TN 1122\)](#) (CR 7818)

01/2008 - CMS has determined that ESA treatment is reasonable and necessary for anemia secondary to myelosuppressive anticancer chemotherapy in solid tumors, multiple myeloma, lymphoma and lymphocytic leukemia under specified conditions. CMS has also determined that ESA treatment is not reasonable and necessary for beneficiaries with certain clinical conditions, either because of a deleterious effect of the ESA on their underlying disease or because the underlying disease increases their risk of adverse effects related to ESA use. Effective date: 07/30/2007 [\(TN 80\)](#) [Implementation date: 04/07/2008.](#) (CR5818)

National Coverage Analyses (NCAs)

This NCD has been or is currently being reviewed under the National Coverage Determination process. The following are existing associations with NCAs, from the National Coverage Analyses database.

- [Original Consideration for Erythropoiesis Stimulating Agents \(ESAs\) for non-renal disease indications \(CAG-00383N\)](#)

Additional Information

Other Versions

Title	Version	Effective Between	
Erythropoiesis Stimulating Agents (ESAs) in Cancer and Related Neoplastic Conditions	1	07/30/2007 - N/A	You are here