
A
Overall survival probability
1.00
0.75
0.50
0.25
0.00
CR (n=273)
PR (n=51)
SD (n=18)
PD (n=30)
Log-rank P<0.001
CV C-index = 0.630
(CV n=343)
0
24
48
72
96
120
Months since end of first-line therapy
Number at risk
CR 273 192 108 56 27 8
PR 51 28 8 0 0 0
SD 18 6 1 0 0 0
PD 30 15 2 0 0 0
B
Overall survival probability
CR (n=273)
NCR (n=99)
Log-rank P<0.001
CV C-index = 0.632
(CV n=343)
Months since end of first-line therapy
Number at risk
CR 273 192 108 56 27 8
NCR 99 49 11 0 0 0
C
Adjusted OS HR (95% CI)
10
5
2
1
PR
SD
PD
Response (vs CR)
Analysis setting
Diagnosis: complete cases (n=408)
6-month landmark (n=382)
Therapy-end landmark (n=343)
D
Unadjusted response-only Cox models
1000 bootstrap samples; ORR selected: 0%
Therapy end (n=372)
Diagnosis Same cohort (n=372)
Diagnosis All labelled (n=453)
11%
62%
27%
9%
68%
23%
15%
24%
62%
0%
25%
50%
75%
100%
Lowest-AIC selection frequency
Response coding
Four-category
Objective-response
CR versus NCR
Three-category (post hoc)
E
Clinical-only median C-index = 0.615; n=343
20×5-fold CV (empirical 5th–95th percentiles)
Objective-response
CR versus NCR
Three-category (post hoc)
Four-category
0.028
0.067
0.066
0.065
0.00
0.03
0.06
0.09
Paired ΔC-index vs clinical model