

A
TCGA-OV cohort and landmark selection
TCGA-OV clinical records, n = 630
Sources: TCGA-CDR, Xena, GDC
Non-primary-tumour records, n = 35
Primary tumours n = 595
Not linked to U133A expression, n = 28
Linked to U133A expression n = 567
No canonical response label, n = 113
Canonical response label, n = 454
CR 319 | PR 64 | SD 30 | PD 41
OS unavailable, n = 1
Overall survival available n = 453
No first-line treatment record, n = 35
First-line treatment record n = 418
No recorded end date n = 21
Recorded end of first-line therapy n = 397
End outside 60–365 days, n = 14
L = end of first-line therapy, 60–365 days after diagnosis n = 383
Died or censored on or before L, n = 11
Landmark cohort
n = 372 (220 deaths)
Alive and under observation beyond L
CR 273 | NCR 99
(PR 51, SD 18, PD 30)
Response codings compared
M1 four-category
M2 objective response
M3 CR versus NCR; M_PD post hoc
Selection and treatment sensitivity
B
External serous cohorts
Response-labelled cohorts
curatedOvarianData and GEO
Fixed 6-month (183-day) landmark from diagnosis
Cohort All Serous Landmark n (deaths)
GSE63885 75 70 69 (61)
GSE23554 28 28 25 (13)
GSE30161 55 45 44 (30)
Cohort-level REML meta-analysis (k = 3)
Total n = 138 (104 deaths)
Pooled CR versus NCR OS (k = 3)
Four-category OS: GSE63885 only
C
Exploratory cross-cancer analyses
TCGA-CDR, 23 tumour types
n = 5,377 labelled
Response-label audit
Treatment definition, resection and power
Four-category topology
OV n = 396; LGG n = 262
Treated BLCA n = 102
(exploratory)
CR versus NCR
from therapy end
6 tumour types, n = 1,142
(OV n = 396)
Parallel analyses; eligibility from TCGA-CDR, not panel A