

A S2 Protein-DNA Span vs Trapping (two-state) B AAI vs Trapping (sensitivity, two-state)

neutral: -0.67 (p 0.267)

niraparib-protonated: -0.05 (p 1.000)


rucaparib-protonated: -0.67 (p 0.267)

neutral: S1 0.31 | AAI +0.21 (n.s.)


niraparib-protonated: S1 0.46 | AAI +0.21 (n.s.)

rucaparib-protonated: S1 0.31 | AAI +0.31 (n.s.)

S2 CV1 Protein-DNA Span (kcal/mol)



Trapping rank (1 = weakest, 5 = most potent)



C Descriptor-state-estimator matrix

no cell attains nominal significance (all exact $p \geq 0.27$); red = inverse, blue = positive

	Neutral				Niraparib-prot.			
S2 CV1	-0.67 (p 0.27)	-0.36 (p 0.63)	-0.36 (p 0.63)	n.d.	-0.05 (p 1.00)	-0.20 (p 0.77)	-0.05 (p 1.00)	n.d.
S2 CV2	-0.05 (p 1.00)	-0.36 (p 0.63)	-0.36 (p 0.63)	n.d.	-0.20 (p 0.77)	-0.36 (p 0.63)	-0.20 (p 0.77)	n.d.
S1	+0.31 (p 0.67)	-0.20 (p 0.77)	+0.05 (p 1.00)	n.d.	+0.46 (p 0.43)	-0.20 (p 0.77)	+0.05 (p 1.00)	n.d.
AAI	+0.20 (p 0.77)	+0.36 (p 0.63)	+0.36 (p 0.63)	n.d.	+0.20 (p 0.77)	+0.36 (p 0.63)	+0.36 (p 0.63)	n.d.
	Rucaparib-prot.				Audited-dominant			
S2 CV1	-0.67 (p 0.27)	-0.36 (p 0.63)	-0.36 (p 0.63)	n.d.	-0.20 (p 0.77)	-0.36 (p 0.63)	-0.20 (p 0.77)	n.d.
S2 CV2	-0.31 (p 0.67)	-0.36 (p 0.63)	-0.62 (p 0.30)	n.d.	-0.31 (p 0.67)	-0.36 (p 0.63)	-0.46 (p 0.43)	n.d.
S1	+0.31 (p 0.67)	+0.05 (p 1.00)	+0.30 (p 1.00)	n.d.	+0.46 (p 0.43)	+0.05 (p 1.00)	+0.05 (p 1.00)	n.d.
AAI	+0.31 (p 0.67)	+0.36 (p 0.63)	+0.46 (p 0.43)	n.d.	+0.31 (p 0.67)	+0.36 (p 0.63)	+0.46 (p 0.43)	n.d.
	min-max	5-95%	95% trunc	eff. support	min-max	5-95%	95% trunc	eff. support
Span estimator								

