
A S2 CV1 Span vs Trapping (state panels)
neutral: -0.67 (p 0.267)
niraparib-protonated: -0.05 (p 1.000)
rucaparib-protonated: -0.67 (p 0.267)
audit-corrected (all three): -0.21 (p 0.767)
S2 CV1 Protein–DNA Span (kcal/mol)
Trapping rank (1 = weakest, 5 = most potent)
Type II
Type III
Rucaparib-protonated
Olaparib
Niraparib-protonated
Veliparib
Rucaparib
Niraparib
Veliparib-protonated
Talazoparib
B Span ratio vs Trapping (sensitivity, state panels)
neutral: S1 0.31 | span ratio +0.21 (n.s.)
niraparib-protonated: S1 0.46 | span ratio +0.21 (n.s.)
rucaparib-protonated: S1 0.31 | span ratio +0.31 (n.s.)
audit-corrected: S1 0.62 | span ratio -0.10 (n.s.)
Trapping rank (1 = weakest, 5 = most potent)
S1/S2 span ratio
Talazoparib
Niraparib
Niraparib-protonated
Rucaparib
Rucaparib-protonated
Olaparib
Veliparib-protonated
Veliparib
C Descriptor–state–estimator matrix
no cell attains nominal significance (all exact p ≥ 0.267); red = inverse, blue = positive
Neutral
Niraparib-prot.
Rucaparib-prot.
Audit-corrected
S2 CV1
S2 CV2
S1
span ratio
-0.67 (p 0.267)
-0.36 (p 0.633)
-0.36 (p 0.633)
n.d.
-0.05 (p 1.000)
-0.21 (p 0.767)
-0.05 (p 1.000)
-0.21 (p 0.767)
-0.36 (p 0.633)
+0.31 (p 0.667)
+0.05 (p 1.000)
+0.46 (p 0.433)
+0.21 (p 0.767)
+0.36 (p 0.633)
-0.31 (p 0.667)
-0.62 (p 0.300)
+0.41 (p 0.500)
+0.67 (p 0.267)
+0.56 (p 0.367)
+0.62 (p 0.300)
-0.10 (p 0.900)
Spearman ρ
1.0
0.5
0.0
-0.5
-1.0
min–max
5–95%
95% trunc
eff. support
Span estimator