Data purity, not model capability, dominates apparent AlphaFold3 performance on phosphorylation benchmarks

Qingqing Mo[1,2], Pingbo Chen[1,2], Cheng Xu[1,2], Ya Wang[1,2], Ting Hu[1,2], Qian Sun[1,2], Tao Zhu[1,2,*]

[1]Department of Obstetrics and Gynecology, National Clinical Research Center for Obstetrics and Gynecology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

[2]Key Laboratory of Cancer Invasion and Metastasis (Ministry of Education), Hubei Key Laboratory of Tumor Invasion and Metastasis, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

[*]Corresponding author. E-mail: zhutao@tjh.tjmu.edu.cn. ORCID: 0009-0001-0779-2245

November 9, 2026

Abstract

Motivation: AlphaFold3 (AF3) accepts explicit post-translational modifications, which is widely taken to imply that it can model the conformational consequences of phosphorylation. Existing benchmarks, however, rarely audit whether the "phosphorylated" and "unphosphorylated" experimental structures that they pair are genuinely comparable, so complex-composition effects can be mistaken for phosphorylation effects.

Results: We assembled a phosphorylation structure benchmark and subjected every pair to a five-criterion biological purity gate (P1–P5). Of 18 curated entries, 16 are two-state pairs and 15 have AF3 models for both states; only 9 pass the full gate. On this cohort the correlation between AF3-predicted and experimentally observed per-residue Cα displacements is not significant ($r = +0.11$, 95% CI $[-0.34, +0.57]$, $n = 15$), while the predicted displacement is nearly constant (0.52 ± 0.36 Å) against an experimental span of 0.25–2.51 Å. Both AF3 internal confidence signals fare no better: the association between site pLDDT and predictive agreement ($r = +0.48$) and that between ΔPAE and predictive agreement ($r = +0.60$) are each carried by a single leverage point and collapse when it is removed ($r = +0.21$ and $r = +0.14$). Restricting to purely paired structures changes the headline statistic, demonstrating that pairing purity, not model capability, governs the apparent result. A second, sequence-only ESMFold line, run with a phosphomimetic substitution, is equally uninformative ($r = -0.36$).

Availability and Implementation: Benchmark data, analysis code and frozen intermediate results are available at https://github.com/tjogzt/ptm-phospho-af3-benchmark (archived at Zenodo: https://doi.org/10.5281/zenodo.24087962); implemented in Python 3.12.

Contact: zhutao@tjh.tjmu.edu.cn

Supplementary information: Supplementary figures and tables are available at *Bioinformatics* online.

Keywords: AlphaFold3, phosphorylation, conformational change, benchmark, data purity, prediction confidence

1 Introduction

Phosphorylation is the most prevalent regulatory post-translational modification (PTM). By altering local electrostatics, hydrogen bonding and backbone flexibility, a single phosphate group can propagate

structural rearrangements far from the modified residue, switching a protein between functional states [1]. Reconstructing the structural consequences of phosphorylation is therefore central to mechanistic interpretation of signalling, and it is precisely the kind of "instance-specific" question that structure-prediction models are now expected to answer.

AlphaFold3 (AF3) marked a qualitative change in this expectation: unlike earlier protein-only predictors, it accepts explicit chemical components and modified residues as input, including phosphoserine (SEP), phosphothreonine (TPO) and phosphotyrosine (PTR) [2]. Because the modification is part of the input, it seems natural to read a difference between an unmodified and a modified AF3 prediction as the model's view of the phosphorylation-induced structural change. Several studies have begun to use AF3 in exactly this "mechanistic" mode [3], and AF3's per-residue confidence (pLDDT) and predicted aligned error (PAE) are frequently quoted as reassurance that the predicted change is trustworthy [4, 5].

Three gaps complicate this practice. First, benchmarking structural responses to phosphorylation requires a pair of experimental structures that differ *only* in the modification. In the Protein Data Bank (PDB), however, a phospho form and a non-phospho form of the same protein are usually crystallised as different constructs: different oligomeric states, different bound ligands, different domain boundaries, sometimes different partners. Any of these covariates induces large conformational differences that have nothing to do with the phosphate, yet such pairs are routinely treated as "phosphorylation pairs". Second, the metrics used to summarise agreement—typically a mean displacement magnitude or an amplitude ratio—are dominated by a handful of disordered, highly mobile residues, so that a large apparent "under-prediction" can be a measurement artefact. Third, pLDDT and PAE measure *self-consistency* of a single prediction, not agreement with a reference; using them as proxies for correctness of a PTM-induced change is a category error unless it is tested.

This work reports a deliberately conservative benchmark that addresses these gaps and, in doing so, arrives at a cautionary negative result. Our contributions are threefold.

1. **A purity audit with biological criteria.** We formalise a five-criterion gate (P1–P5) covering identity, presence of a genuine modified residue, two-state coverage, overlap about the target site, and chain-composition consistency, and we show that a substantial fraction of apparently usable phosphorylation pairs fail at least one criterion.

2. **A negative result on a cleaned cohort.** On the pairs that survive the gate we find no evidence that AF3's predicted phosphorylation-induced displacements track the experimentally observed ones, we show that AF3's predicted displacement is close to a constant, and we find that a second, sequence-only predictor (ESMFold) behaves the same way.

3. **Two falsified confidence hypotheses.** We test whether site pLDDT or ΔPAE predicts agreement, and show that both apparent associations are produced by a single leverage point and disappear when it is removed.

We do not claim that AF3 is incapable of modelling phosphorylation effects; we claim that the widely used benchmark protocol cannot currently demonstrate that it does, and we give concrete, low-cost practices that avoid the two failure modes we identify.

2 Background and related work

AF3 and explicit modified residues. AF3 takes an explicit list of residues and ligands, so a phospho form can be specified directly [2]. This capability is orthogonal to accuracy on modification-induced change, yet the two are often conflated. Prior evaluations of AF3 on modified or non-

protein components [3] focus on local geometry around the modified residue rather than on global conformational responses.

Structural-change benchmarks. Comparing two states of a protein is a mature problem: one superposes a common core (e.g. Kabsch [6]) and measures per-residue displacement, or decomposes the motion into rigid-block (hinge) components [7]. The choice of core and the treatment of disordered regions strongly affect the reported magnitude, which is why amplitude summaries must be interpreted against a noise model.

Confidence as a proxy for correctness. pLDDT and PAE are estimates of local and pairwise geometric confidence within a single prediction [2]. Their use as proxies for agreement with an independent reference is common but rarely validated on a paired structural-change benchmark; when it has been examined, the relationship is weak or confounded [4, 5].

3 Materials and methods

3.1 Benchmark construction

Candidate pairs were collected from the PDB as pairs of structures of the same UniProt entry in which exactly one state carries a phosphorylated residue, and were labelled with an effect type (*local rearrangement*, *allosteric*, or *disorder-to-order*). After curation the benchmark contains 18 entries, of which 16 are two-state pairs; AF3 models for both states were obtained for 15 of them, and these 15 form the *primary cohort* (Table 1). All downstream analysis uses a single frozen manifest (`manifest_frozen.json`) that binds each protein to its UniProt accession, phosphorylated site(s) in UniProt coordinates, and its phospho/unphospho PDB identifiers and chains, so that every number in this paper is traceable to one immutable table.

3.2 Pairing-purity gate (P1–P5)

Each candidate pair is adjudicated by five biological criteria, evaluated independently for the phospho and non-phospho structures.

P1 (identity). Both structures map to the same UniProt entry at high sequence identity.

P2 (genuine modification). The phospho structure contains at least one modified residue (SEP, TPO or PTR) recorded in the PDB MODRES records; a structure that merely lacks the phosphate but is otherwise wild type qualifies as the non-phospho state.

P3 (two-state coverage). Each state covers at least 50% of the UniProt sequence.

P4 (overlap about the site). The two states share at least 20 common residues and the shared region includes the target site.

P5 (composition consistency). The two states have the same chain composition, so that differences in oligomeric state or in bound partners cannot be mistaken for a phosphorylation effect. A pair that fails any of P1–P4 is excluded; a pair that fails only P5 is retained but flagged (*impure*), because P5 violations are the most common and most insidious source of confounding. In the primary cohort, 9 pairs pass all five criteria (the *pure* cohort) and 6 do not (4 flagged on P5, 2 excluded on P3/P5).

3.3 Prediction protocols

AlphaFold3. For each protein we submitted the full-length UniProt sequence with the approved site(s) specified as modified residues using the corresponding CCD codes, and collected 15 models per state (five diffusion samples across three random seeds). Analysis aggregates all models unless otherwise stated; the seed axis is treated as sampling repetition, not biological replication.

ESMFold (comparison line). As an independent, sequence-only predictor we folded the same full-length sequences with ESMFold [8]. ESMFold does not parse CCD modified residues, so the phospho state was represented by a phosphomimetic substitution (pSer/pThr→Asp, pTyr→Glu); this limitation is stated explicitly and the comparison is therefore qualitative.

3.4 Displacement metrics

All comparisons share one coordinate protocol. The non-phospho experimental structure defines the reference frame; the phospho experimental structure is superposed onto it over common Cα atoms by the Kabsch algorithm [6], giving the experimental displacement field Δ_{exp}. Predicted fields are obtained by superposing each predictor's non-phospho model onto the same reference and applying that transform to its phospho model, giving Δ_{pred} without ever superposing predicted onto experimental coordinates. From these fields we compute: (i) the per-residue displacement magnitude; (ii) the amplitude ratio, $|\Delta_{\text{pred}}|/|\Delta_{\text{exp}}|$; (iii) the direction cosine between Δ_{pred} and Δ_{exp}; (iv) the Pearson correlation r of the per-residue magnitude profiles; (v) a hinge index from a rigid-block superposition; and (vi) a site window of ± 10 residues around each modified site. Modified residues (SEP/TPO/PTR) are retained throughout; omitting them, as a naive HETATM filter does, silently removes the residue of interest from both sides of the comparison.

3.5 Statistics

The independent unit is the *protein*, not the model: the 15 AF3 models of one protein share a single experimental reference and are not independent observations. Cohort correlations are therefore computed over proteins, and bootstrap confidence intervals resample proteins, not models. Sensitivity to single points is assessed by leave-one-out and leverage diagnostics. Correlations are reported as Pearson r with Spearman ρ where a monotone alternative is relevant; no correction for multiplicity is applied to the exploratory correlations, which are labelled as such.

3.6 Same-state noise floor

To decide whether experimentally observed displacements are distinguishable from noise, we computed two *same-state* controls: (i) an experimental floor, the Cα dispersion after superposing crystallographically independent copies (NCS) of the same chain in the same state; and (ii) a predictor floor, the dispersion among the 15 AF3 models of the same state. Because AF3 predicts full-length sequences whereas experimental structures cover only the ordered core, the predictor floor is evaluated on the residues actually resolved in the experiment, so that unmodelled termini do not inflate it.

4 Results

4.1 The benchmark shrinks under a purity gate

Of 18 curated entries, 16 are two-state pairs and 15 have AF3 models for both states (Fig. 1a). Applying P1–P5 leaves 9 pure pairs; the remaining 6 are either compositionally inconsistent (P5; IKBKB, IRF3, CDK7, CDK1) or fail coverage/overlap (P3/P5; AMPKα2, TBK1). The dominant failure mode is P5: the phospho and non-phospho entries differ in chain count or partners, exactly the covariate that a naive amplitude comparison would attribute to phosphorylation.

This is not a cosmetic filter. Stratifying the same frozen per-protein results by purity changes the apparent behaviour of the model (Fig. 1b): the median amplitude ratio is 0.46 for pure pairs but 0.30 for impure pairs, and the magnitude correlation is $r = +0.22$ ($n = 9$) for pure pairs but $r = -0.24$

Protein	UniProt	Effect	PTM	Site(s)	Unphos PDB	Phos PDB	P1	P2	P3	P4	P5	Verdict	Pure	In cohort
CDK2	P24941	local	pT	Thr160 (pT)	2C5N	1H1R	Y	Y	Y	Y	Y	PASS	Y	Y
MAPK1	P28482	local	pT/pY	Thr185/Tyr187 (pT/pY)	1ERK	2ERK	Y	Y	Y	Y	Y	PASS	Y	Y
AURKA	O14965	local	pT	Thr288 (pT)	4J8M	5DNR	Y	Y	Y	Y	Y	PASS	Y	Y
IRF3	Q14653	allost.	pS	Ser386/Ser402 (pS)	1QWT	3A77	Y	Y	Y	Y	N	WARN	N	Y
MAPK14	Q16539	local	pT/pY	Thr180/Tyr182 (pT/pY)	1WFC	3PY3	Y	Y	Y	Y	Y	PASS	Y	Y
CDK7	P50613	local	pT	Thr170 (pT)	8P79	8PYR	Y	Y	Y	Y	N	WARN	N	Y
SRC	P12931	allost.	pY	Tyr419 (pY)	7NG7	9NS0	Y	Y	Y	Y	Y	PASS	Y	Y
PRKAA2	P54646	allost.	pT	Thr172 (pT)	2H6D	6B1U	Y	Y	N	Y	N	FAIL	N	Y
PRKACA	P17612	allost.	pT	Thr198 (pT)	–	6QJ7	Y	Y	Y	N	N	PASS	N	N
POLR2A	P24928	disorder	pS	Ser1614/Ser1621 (pS)	–	4JXT	Y	Y	N	N	N	PASS	N	N
PYGM	P00489	disorder	pS	Ser15 (pS)	2GJ4	1LWO	Y	Y	Y	Y	Y	PASS	Y	Y
CDK1	P06493	local	pT	Thr161 (pT)	4YC3	9SKQ	Y	Y	Y	Y	N	WARN	N	Y
MAPK3	P27361	local	pT/pY	Thr202/Tyr204 (pT/pY)	4QTB	2ZOQ	Y	Y	Y	Y	Y	PASS	N	N
GSK3B	P49841	local	pY	Tyr216 (pY)	1J1B	6Y9S	Y	Y	Y	Y	Y	PASS	Y	Y
PAK1	Q13153	local	pT	Thr423 (pT)	1YHW	3FXZ	Y	Y	Y	Y	Y	PASS	Y	Y
AKT1	P31749	local	pT	Thr308 (pT)	3O96	4GV1	Y	Y	Y	Y	Y	PASS	Y	Y
TBK1	Q9UHD2	local	pS	Ser172 (pS)	4IM0	4EUU	Y	Y	N	Y	N	FAIL	N	Y
IKBKB	O14920	local	pS	Ser177/Ser181 (pS)	4E3C	4KIK	Y	Y	Y	Y	N	WARN	N	Y

Table 1: Benchmark cohort and pairing-purity gate (P1–P5). P1 identity, P2 real phospho-residue (MODRES), P3 coverage ≥0.5 in both states, P4 ≥20-residue overlap covering the target site, P5 identical chain composition.

($n = 6$) for impure ones. Any headline computed on the uncleaned cohort is therefore a mixture of model behaviour and pairing confounding. We report all subsequent statistics on the full 15-pair cohort *and* flag the pure subset, and we avoid any claim whose sign depends on the mixture.

4.2 No evidence that AF3 tracks the phosphorylation-induced displacement

Protein	UniProt	n_{common}	exp disp (Å)	AF3 disp (Å)	AF3 ratio	AF3 cos	AF3 r	ESM disp (Å)	ESM ratio	ESM cos	ESM r	Pure
AKT1	P31749	262	2.05	0.81	0.39	-0.40	+0.73	0.08	0.04	+0.55	+0.68	Y
PRKAA2	P54646	255	2.22	0.20	0.09	+0.02	+0.10	0.04	0.02	+0.18	+0.77	N
AURKA	O14965	263	2.44	0.29	0.12	+0.34	+0.85	0.04	0.02	+0.20	+0.68	Y
CDK1	P06493	283	1.44	0.71	0.49	+0.43	+0.62	0.06	0.04	+0.41	+0.39	N
CDK2	P24941	296	0.78	0.44	0.56	+0.26	+0.37	0.10	0.13	+0.17	+0.63	Y
CDK7	P50613	259	0.53	0.18	0.34	+0.22	+0.43	0.07	0.12	-0.72	+0.10	N
MAPK1	P28482	352	1.64	1.06	0.65	+0.66	+0.61	0.07	0.04	-0.01	+0.55	Y
GSK3B	P49841	344	0.63	0.15	0.24	+0.17	+0.45	1.53	2.43	+0.14	+0.13	Y
PYGM	P00489	803	0.94	0.43	0.46	-0.08	+0.34	0.61	0.65	+0.73	+0.99	Y
IKBKB	O14920	617	1.96	0.53	0.27	+0.17	+0.38	0.05	0.02	+0.15	+0.40	N
IRF3	Q14653	232	0.33	0.73	2.22	+0.06	+0.38	0.61	1.85	+0.31	+0.33	N
MAPK14	Q16539	338	1.57	1.39	0.89	+0.70	+0.96	0.06	0.04	+0.38	+0.52	Y
PAK1	Q13153	291	0.25	0.24	0.95	–	+0.26	0.03	0.12	–	+0.00	Y
TBK1	Q9UHD2	279	1.50	0.40	0.27	+0.56	+0.40	0.05	0.03	+0.07	+0.67	N
SRC	P12931	274	2.51	0.25	0.10	-0.07	+0.03	0.40	0.16	+0.09	+0.72	Y

Table 2: Per-protein comparison of experimental versus predicted phospho-induced Cα displacement. Displacements are mean per-residue magnitudes (Å). Ratio = predicted/experimental. cos = mean direction cosine over residues moving >1 Å. r = Pearson r of the per-residue magnitude profiles.

Over the 15-pair cohort the per-protein mean predicted displacement is essentially uncorrelated with the experimental displacement: Pearson $r = +0.11$ (95% CI $[-0.34, +0.57]$), slope $+0.053$ Å/Å, Spearman $\rho = +0.15$ ($p = 0.59$) (Fig. 2a). The confidence interval is wide—$n = 15$ is small—but it excludes the strong positive association that "AF3 models phosphorylation" would require. The predicted displacement is also close to constant: 0.52 ± 0.36 Å across all proteins, against an experimental span of 0.25–2.51 Å (median 1.50 Å). Put differently, the model's output moves by roughly the same small amount regardless of whether the true response is negligible (PAK1, 0.25 Å) or large (Src, 2.51 Å).

Leave-one-out analysis shows the coefficient is not driven by a single outlier in the sense of an

extreme r; the largest change is the removal of Src, which raises r to +0.22, still far from significant. The per-protein picture is heterogeneous rather than uniformly null: a few proteins show high magnitude correlation (p38, $r = +0.96$; AURKA, $r = +0.85$), but several show none (Src, $r = +0.03$) even though their experimental displacements are among the largest. Aggregated over the cohort, this heterogeneity averages out to noise.

4.3 A second, sequence-only predictor is equally uninformative

The non-response could in principle be specific to AF3's structure module, which consumes an MSA. We therefore repeated the identical coordinate protocol with ESMFold, a single-sequence language-model predictor, using a phosphomimetic substitution to represent the phospho state (Sec. 3.3). The outcome is the same and, quantitatively, even flatter: the cohort-mean predicted displacement is 0.25 ± 0.41 Å (against 0.52 ± 0.36 Å for AF3) and shows no association with the experimental displacement ($r = -0.36$, 95% CI $[-0.72, +0.31]$; Spearman $\rho = -0.24$, $p = 0.40$) (Table 2, Fig. 2a). For most proteins the phosphomimetic substitution moves the predicted structure by less than 0.1 Å, i.e. essentially not at all, so the near-constant prediction is not unique to AF3. Leave-one-out again exposes the small cohort: removing the lowest- displacement protein (PAK1) moves the correlation to $r = -0.47$, while no single removal produces a significant positive association. ESMFold's per-protein direction cosines are also centred near zero and its per-residue magnitude correlations are uninformative (Table 2, Fig. 2b). The failure to track the phosphorylation response is thus shared across two predictors, two input modalities (MSA-based and sequence-only) and two ways of representing the phosphate, which argues that it is a property of the benchmark question rather than of one model.

4.4 The predicted signal sits at the same-state noise floor

The negative result is only meaningful if the experimental signal exceeds noise. It does. The same-state experimental floor has a median of 0.335 Å and a maximum of 1.309 Å across the seven proteins with crystallographically independent copies, whereas the phosphorylation-induced experimental displacement has a median of 1.50 Å and a span of 0.25–2.51 Å (Fig. 3). The experimentally observed change therefore exceeds the upper bound of same-state noise, so the absence of predicted tracking is not an artefact of noisy references. The predictor's own same-state floor (median 0.389 Å) is of the same order as its putative phospho signal (0.52 Å), reinforcing that AF3's between-state displacement is close to its own sampling variability.

4.5 Two confidence-based hypotheses are refuted

If the model's internal confidence flagged where it was right, a user could at least triage predictions. We tested two natural hypotheses on the 15-pair cohort.

Site pLDDT. The correlation between the unphosphorylated site pLDDT and the amplitude ratio is $r = +0.48$ (Fig. 4). The relation is carried almost entirely by IRF3, whose amplitude ratio (2.22) and site pLDDT (89.9) are both extreme: removing this single protein drops the correlation to $r = +0.21$ ($n = 14$). A companion analysis using the site-minus-global pLDDT contrast ($r = +0.43$) behaves the same way.

ΔPAE. PAE is the more natural signal for a *pairwise* question such as "does the domain carrying the site move relative to the rest?". The correlation between ΔPAE (site→rest) and the amplitude ratio is $r = +0.60$ (Fig. 5), the strongest association we observe. It is again an IRF3 effect: removing IRF3 drops it to $r = +0.14$ ($n = 14$), and the analogous statistic for global ΔPAE falls from $r = +0.49$ to $r = +0.06$.

Both confidence signals are thus indistinguishable from single-point artefacts on this benchmark. Their apparent strength is an artefact of a small cohort containing one protein whose prediction and reference happen to disagree in both magnitude and confidence in the same direction.

4.6 Exploratory: what does the model respond to?

Seeking a physical explanation, we asked whether AF3 under-responds most where the true motion is largest. Across the 15 pairs the amplitude ratio is inversely related to the overall deformation magnitude ($r = -0.60$, $p = 0.017$ against the full-structure RMSD; $r = -0.62$, $p = 0.014$ against mean displacement) and to the block-wise deformation ($r = -0.60$, $p = 0.018$); the relation with the hinge index is weaker and not significant ($r = -0.40$, $p = 0.14$). A residue-level view agrees: binning all 3,355 common residues by experimental displacement, the median predicted/experimental ratio falls monotonically from 0.54 in the smallest-displacement bin to 0.05 in the largest, i.e. AF3's response saturates while the true response grows (Fig. 6). Consistent with a hinge explanation, large-motion residues cluster about the modified site; residues far from the site (> 80 residues away) show the weakest and least correlated response. Because these stratifications are exploratory and uncorrected for multiplicity, we present them as hypotheses, not conclusions; in particular, the hinge comparison is not significant and we do not claim that hinge motion is harder to predict.

5 Discussion

What the benchmark does and does not show. Our central empirical claim is narrow: on a 15-pair phosphorylation structure benchmark, AF3's predicted conformational response is not related to the observed one, and the two most commonly cited internal confidence signals do not identify where it succeeds. We deliberately do *not* claim that AF3 cannot model phosphorylation. The cohort is small; the metric is a single, rigid-body-based displacement field; and "no detectable correlation at $n = 15$" is weaker than "no correlation". The honest statement is that the current, commonly used protocol cannot demonstrate the capability it is assumed to have.

Why the result is easy to miss. Two artefacts work in the same direction. First, pairing impurity inflates apparent agreement: phospho and non-phospho entries that differ in oligomeric state or partner differ in structure for reasons unrelated to the phosphate, and pooling such pairs changes the sign of the cohort correlation (Sec. 4.1, Fig. 1b). Second, amplitude summaries are dominated by a few highly mobile residues, so a large apparent "under-prediction" can reflect disorder rather than a mechanical failure. Both artefacts are avoidable at essentially no cost: apply a composition check before pairing, and report displacement statistics against a same-state noise floor.

Confidence is not correctness. Our two refutations share a mechanism: a single protein (IRF3) is extreme in both the predictor's confidence and in its disagreement, and on a 15-point cohort one such point can manufacture a correlation of 0.5–0.6. This is a warning about study design as much as about AF3: on small structural benchmarks, leverage diagnostics are not optional.

Practical guidance. (i) Audit pair purity before computing any agreement statistic; a composition check (P5) removes the largest confounder. (ii) Never drop modified residues from the comparison; they are the object of study. (iii) Calibrate measured displacements against same-state replicates; an effect smaller than the same-state floor is not interpretable. (iv) Do not use pLDDT or PAE as evidence that a PTM-induced change is correct; test that association explicitly and check its leverage.

Limitations. The cohort is modest ($n = 15$, 9 pure) and kinase-dominated, so the effect-type strata are under-powered and we report them only descriptively. ESMFold is included as a second, sequence-only line; because it cannot represent CCD modified residues its phospho state is phosphomimetic, so that line is qualitative, yet it agrees with AF3 in showing no tracking ($r = -0.36$). No molecular-

dynamics reference is used. These limitations bound the generality of the negative result; they do not affect the two methodological findings, which are properties of the protocol.

6 Conclusion

On a phosphorylation structure benchmark cleaned by biological purity criteria, AF3's predicted conformational response does not track the experimental one, its predicted displacement is near-constant, and its internal confidence signals fail to identify where it is right once single-point leverage is removed. The more transferable lesson is methodological: on paired structural-change benchmarks, pairing purity and same-state noise calibration—not model capability—determine the apparent result, and both are cheap to enforce.

Availability and implementation

The frozen benchmark manifest, analysis scripts (Python 3.12), and all intermediate results are archived at https://github.com/tjogzt/ptm-phospho-af3-benchmark (Zenodo DOI: https://doi.org/10.5281/zenodo.24087962). The analysis is deterministic given the frozen manifest; random seeds are fixed. AF3 predictions were generated with the AlphaFold Server (alphafoldserver.com); ESMFold inference was run locally with `transformers` on macOS (Apple Silicon, MPS backend). Codes are released under the MIT licence.

Funding

This work was supported by the National Natural Science Foundation of China (Grant Nos. 82403759 to C.X. and 82403616 to Y.W.).

Conflict of interest

None declared.

References

[1] Louise N. Johnson and Richard J. Lewis. Structural basis for control by phosphorylation. *Chemical Reviews*, 101(8):2209–2242, 2001. doi: 10.1021/cr000225s.

[2] Josh Abramson, Jonas Adler, Jack Dunger, Richard Evans, Tim Green, Alexander Pritzel, Olaf Ronneberger, Lindsay Willmore, Andrew J. Ballard, Joshua Bambrick, et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*, 630(8016):493–500, 2024. doi: 10.1038/s41586-024-07487-w.

[3] Ki Hyun Nam. Evaluation of AlphaFold3 prediction for post-translational modification, oligomeric assembly, and quenchable metal binding of fluorescent proteins. *Journal of Molecular Graphics and Modelling*, 142:109169, 2026. doi: 10.1016/j.jmgm.2025.109169.

[4] Oliviero Carugo. pLDDT values in AlphaFold2 protein models are unrelated to globular protein local flexibility. *Crystals*, 13(11):1560, 2023. doi: 10.3390/cryst13111560.




Figure 1: **Pairing purity shapes the benchmark and the conclusion.** (a) Funnel from the curated candidate pool through the two-state and AF3-availability filters to the pairs passing the P1–P5 purity gate. (b) The same frozen per-protein results stratified by pairing purity, showing the median amplitude ratio, median direction cosine and magnitude correlation for pure ($n = 9$), impure ($n = 6$) and all ($n = 15$) pairs.

[5] Mehmet Akdel, Douglas E. V. Pires, Eduard Porta Pardo, Jürgen Jänes, Arthur O. Zalevsky, Bálint Mészáros, et al. A structural biology community assessment of AlphaFold2 applications. *Nature Structural & Molecular Biology*, 29(11):1056–1067, 2022. doi: 10.1038/s41594-022-00849-w.

[6] Wolfgang Kabsch. A solution for the best rotation to relate two sets of vectors. *Acta Crystallographica Section A*, 32(5):922–923, 1976. doi: 10.1107/S0567739476001873.

[7] Steven Hayward and Herman J. C. Berendsen. Systematic analysis of domain motions in proteins from conformational change: New results on citrate synthase and t4 lysozyme. *Proteins: Structure, Function, and Genetics*, 30(2):144–154, 1998. doi: 10.1002/(SICI)1097-0134(19980201)30:2⟨144::AID-PROT4⟩3.0.CO;2-N.

[8] Zeming Lin, Halil Akin, Roshan Rao, Brian Hie, Zhongkai Zhu, Wenting Lu, Nikita Smetanin, Robert Verkuil, Ori Kabeli, Yaniv Shmueli, et al. Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science*, 379(6637):1123–1130, 2023. doi: 10.1126/science.ade2574.




Figure 2: **Predicted versus experimental phosphorylation-induced displacement.** (a) Per-protein mean experimental displacement against the predicted displacement for AF3 (circles) and ESMFold with a phosphomimetic phospho state (triangles); the dashed line is 1:1. Open markers denote purely paired structures. (b) Per-protein direction cosine and magnitude correlation for AF3 (bars) and ESMFold (diamonds). Neither predictor's displacement tracks the experimental one.




Figure 3: **Same-state noise floor.** Experimental same-state dispersion (crystallographically independent copies) and predictor same-state dispersion (across AF3 models) compared with the phosphorylation-induced experimental displacement.




Figure 4: **Site pLDDT does not predict agreement.** Site pLDDT against amplitude ratio; the association is carried by IRF3 and weakens sharply when it is removed.




Figure 5: **ΔPAE association is a single-point effect.** ΔPAE (site→rest) against amplitude ratio, all pairs (left) and after removing IRF3 (right).




Figure 6: **Residue-level response saturates.** Median predicted/experimental displacement ratio as a function of experimental displacement magnitude, showing monotone decline as the true response grows.