


Enumerating Weak Order Unit Fubini Rankings

Eban Barrett, Maritza Sanchez, and Tung (TJ) Vo

July 29, 2026

Plan for Talk

- Definitions
  - Tuples and Weak Order
  - Fubini Rankings
  - Unit Interval Parking Functions
  - Unit Fubini Rankings
- Motivation
- Main Result (Enumerating Sets)




Definitions: Tuples and Weak Order

Tuples and Weak Order




Definition (Tuple)

A *tuple* of length n is a finite set of ordered elements, denoted by $\alpha = (a_1, a_2, \ldots, a_n) \in [n]^n$.

Tuples and Weak Order




Definition (Tuple)

A *tuple* of length n is a finite set of ordered elements, denoted by $\alpha = (a_1, a_2, \ldots, a_n) \in [n]^n$.

Definition (Weakly increasing rearrangement)

The *weakly increasing rearrangement* of α is $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$ such that $a'_i \leq a'_{i+1}$ for all $i \in [n-1]$.

Tuples and Weak Order

Definition (Tuple)

A *tuple* of length n is a finite set of ordered elements, denoted by $\alpha = (a_1, a_2, \ldots, a_n) \in [n]^n$.

Definition (Weakly increasing rearrangement)

The *weakly increasing rearrangement* of α is $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$ such that $a'_i \leq a'_{i+1}$ for all $i \in [n-1]$.

Definition (Weakly decreasing rearrangement)

The *weakly decreasing rearrangement* of α is $\alpha^{\downarrow} = (a'_1, a'_2, \ldots, a'_n)$ such that $a'_i \geq a'_{i+1}$ for all $i \in [n-1]$.




Definitions: Fubini Rankings

Fubini Rankings

Definition (Fubini ranking)

Let $\alpha = (a_1, a_2, ..., a_n) \in [n]^n$. Then, α is a *Fubini ranking* of length n if for all $r \in [n]$, if k entries of α are equal to r, then the next larger value in α is $r + k$. The set of Fubini rankings of length n is denoted FR_n.

Fubini Rankings





Example

Take $\alpha = (\underline{3}, \underline{1}, \underline{3}, \underline{6}, \underline{3}, \underline{1}) \in \mathrm{FR}_6$, and read as "competitor i placed in a_i-th rank."

Fubini Rankings

Example


Take $\alpha = (\underline{3}, \underline{1}, \underline{3}, \underline{6}, \underline{3}, \underline{1}) \in \mathrm{FR}_6$, and read as "competitor i placed in a_i-th rank."



Fubini Rankings

Example

Take $\alpha = (\underline{3}, \underline{1}, \underline{3}, \underline{6}, \underline{3}, \underline{1}) \in \mathrm{FR}_6$, and read as "competitor i placed in a_i-th rank."



Note: FR_n is permutation invariant!

Lemma [1, Ager-Hart et al., 2026]

Given $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n$, let $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$ be its weakly increasing rearrangement. Then, for all $i \in [n]$, either $a'_i = i$ or $a'_i = a'_{i-1}$.

Fubini Rankings

Lemma [1, Ager-Hart et al., 2026]

Given $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n$, let $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$ be its weakly increasing rearrangement. Then, for all $i \in [n]$, either $a'_i = i$ or $a'_i = a'_{i-1}$.

Example

- Take $\alpha = (3, 1, 1) \in \mathrm{FR}_3$. We have $\alpha^{\uparrow} = (a'_1, a'_2, a'_3) = (1, 1, 3)$.

Fubini Rankings

Lemma [1, Ager-Hart et al., 2026]

Given $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n$, let $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$ be its weakly increasing rearrangement. Then, for all $i \in [n]$, either $a'_i = i$ or $a'_i = a'_{i-1}$.

Example

- Take $\alpha = (3, 1, 1) \in \mathrm{FR}_3$. We have $\alpha^{\uparrow} = (a'_1, a'_2, a'_3) = (1, 1, 3)$.
- Take $\alpha = (2, 1, 1) \notin \mathrm{FR}_3$. We have $\alpha^{\uparrow} = (a'_1, a'_2, a'_3) = (1, 1, 2)$.




Definitions: Unit Interval Parking Functions

Unit Interval Parking Functions




Definition (Unit interval parking function)

Let $\alpha = (a_1, a_2, ..., a_n) \in [n]^n$, and read as "car i has parking preference a_i." Then, α is a *unit interval parking function* of length n if for all $i \in [n]$, car i can park in its preferred spot a_i or the spot right after, which is $a_i + 1$, i.e., all cars park and no parking spots are left empty. The set of unit interval parking functions of length n is denoted UPF_n.

Unit Interval Parking Functions






Example

Take $\alpha = (1, 1, 2) \in \mathrm{UPF}_3$.

Unit Interval Parking Functions

Example

Take $\alpha = (1, 1, 2) \in \mathrm{UPF}_3$. Let's park!



Unit Interval Parking Functions





Example


Take $\alpha = (1, 1, 2) \in \mathrm{UPF}_3$. Let's park!



Unit Interval Parking Functions

Example

Take $\alpha = (1, 1, 2) \in \mathrm{UPF}_3$. Let's park!



Unit Interval Parking Functions



Example

Take $\alpha = (1, 1, 2) \in \mathrm{UPF}_3$. Let's park!

Unit Interval Parking Functions

Example

Take $\alpha = (1, 1, 2) \in \mathrm{UPF}_3$. Let's park!



Caution: UPF_n is not permutation invariant!

Unit Interval Parking Functions




Example


We have $(1,1,2) \in \mathrm{UPF}_3$, but $(2,1,1) \notin \mathrm{UPF}_3$.

Unit Interval Parking Functions




Example

We have $(1,1,2) \in \mathrm{UPF}_3$, but $(2,1,1) \notin \mathrm{UPF}_3$. Let's park!



Unit Interval Parking Functions

Example


We have $(1,1,2) \in \mathrm{UPF}_3$, but $(2,1,1) \notin \mathrm{UPF}_3$. Let's park!



Unit Interval Parking Functions

Example


We have $(1,1,2) \in \mathrm{UPF}_3$, but $(2,1,1) \notin \mathrm{UPF}_3$. Let's park!



Unit Interval Parking Functions

Example

We have $(1, 1, 2) \in \mathrm{UPF}_3$, but $(2, 1, 1) \notin \mathrm{UPF}_3$. Let's park!






Lemma

Let $\alpha \in \mathrm{UPF}_n$. If entry $i \in \alpha$ repeats, it repeats once.




Lemma

Let $\alpha \in \mathrm{UPF}_n$. If entry $i \in \alpha$ repeats, it repeats once.

Proof by Example


Consider $\alpha = (1, 1, 1) \notin \mathrm{UPF}_3$.

Lemma

Let $\alpha \in \mathrm{UPF}_n$. If entry $i \in \alpha$ repeats, it repeats once.

Proof by Example

Consider $\alpha = (1, 1, 1) \notin \mathrm{UPF}_3$. Let's park!



Lemma

Let $\alpha \in \mathrm{UPF}_n$. If entry $i \in \alpha$ repeats, it repeats once.

Proof by Example

Consider $\alpha = (1, 1, 1) \notin \mathrm{UPF}_3$. Let's park!




Lemma

Let $\alpha \in \mathrm{UPF}_n$. If entry $i \in \alpha$ repeats, it repeats once.

Proof by Example

Consider $\alpha = (1, 1, 1) \notin \mathrm{UPF}_3$. Let's park!




Lemma

Let $\alpha \in \mathrm{UPF}_n$. If entry $i \in \alpha$ repeats, it repeats once.

Proof by Example

Consider $\alpha = (1, 1, 1) \notin \mathrm{UPF}_3$. Let's park!






Definitions: Unit Fubini Rankings

Unit Fubini Rankings





Lemma (Characterization of unit Fubini rankings)

For all $n \geq 1$, $\alpha = (a_1, a_2, \ldots, a_n) \in [n]^n$ is in $\mathrm{FR}_n \cap \mathrm{UPF}_n$ if and only if the weakly increasing rearrangement of α, denoted by $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$, satisfies the following:

(1) either $a'_i = i$ or $a'_i = a'_{i-1}$ for all $i \in [n]$

(2) if entry $i \in \alpha$ repeats, it repeats once.

Unit Fubini Rankings





Lemma (Characterization of unit Fubini rankings)

For all $n \geq 1$, $\alpha = (a_1, a_2, \ldots, a_n) \in [n]^n$ is in $\mathrm{FR}_n \cap \mathrm{UPF}_n$ if and only if the weakly increasing rearrangement of α, denoted by $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$, satisfies the following:

(1) either $a'_i = i$ or $a'_i = a'_{i-1}$ for all $i \in [n]$

(2) if entry $i \in \alpha$ repeats, it repeats once.

Proof Sketch

(1) Required by Fubini rankings [1, Ager-Hart et al., 2026].

Unit Fubini Rankings

Lemma (Characterization of unit Fubini rankings)

For all $n \geq 1$, $\alpha = (a_1, a_2, \ldots, a_n) \in [n]^n$ is in $\mathrm{FR}_n \cap \mathrm{UPF}_n$ if and only if the weakly increasing rearrangement of α, denoted by $\alpha^{\uparrow} = (a'_1, a'_2, \ldots, a'_n)$, satisfies the following:

(1) either $a'_i = i$ or $a'_i = a'_{i-1}$ for all $i \in [n]$

(2) if entry $i \in \alpha$ repeats, it repeats once.

Proof Sketch

(1) Required by Fubini rankings [1, Ager-Hart et al., 2026].

(2) Required by unit interval parking functions (Lemma).

Motivation

Motivation

Theorem [3, Elder et al., 2025]

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Motivation




Theorem [3, Elder et al., 2025]

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Proof Sketch

1. For $n = 1$, $|\mathrm{FR}_1^{\uparrow} \cap \mathrm{UPF}_1^{\uparrow}| = 1$.

Motivation

Theorem [3, Elder et al., 2025]

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Proof Sketch

1. For $n = 1$, $|\mathrm{FR}_1^{\uparrow} \cap \mathrm{UPF}_1^{\uparrow}| = 1$.
2. For $n = 2$, $|\mathrm{FR}_2^{\uparrow} \cap \mathrm{UPF}_2^{\uparrow}| = 2$.

Motivation

Theorem [3, Elder et al., 2025]

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Proof Sketch

1. For $n = 1$, $|\mathrm{FR}_1^{\uparrow} \cap \mathrm{UPF}_1^{\uparrow}| = 1$.
2. For $n = 2$, $|\mathrm{FR}_2^{\uparrow} \cap \mathrm{UPF}_2^{\uparrow}| = 2$.
3. The cases are disjoint and exhaustive, satisfying a recursion relation.

Motivation





Question: What is the cardinality of

- $\mathrm{UPF}_n^{\downarrow}$
- $\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}$?




Main Result (Enumerating Sets)

Main Result

Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Main Result

Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

First, we want $|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}|$.

Proof of Main Result





Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

Proof of Main Result





Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait!

Proof of Main Result




Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof of Main Result

Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof by Example

Take $\alpha = (4, 5, 2, 1, 2, 5) \in \mathrm{FR}_6 \cap \mathrm{UPF}_6$.

Proof of Main Result

Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof by Example

Take $\alpha = (4, 5, 2, 1, 2, 5) \in \mathrm{FR}_6 \cap \mathrm{UPF}_6$. Then, $\alpha^{\uparrow} = (1, \boxed{2, 2}, 4, \boxed{5, 5})$.

Proof of Main Result

Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof by Example

Take $\alpha = (4, 5, 2, 1, 2, 5) \in \mathrm{FR}_6 \cap \mathrm{UPF}_6$. Then, $\alpha^{\uparrow} = (1, \boxed{2, 2}, 4, \boxed{5, 5})$.

- Cars not in a box park in their preferred spot.

Proof of Main Result

Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof by Example

Take $\alpha = (4, 5, 2, 1, 2, 5) \in \mathrm{FR}_6 \cap \mathrm{UPF}_6$. Then, $\alpha^{\uparrow} = (1, \boxed{2, 2}, 4, \boxed{5, 5})$.

- Cars not in a box park in their preferred spot.
- First car in each box parks in its preferred spot.

Proof of Main Result

Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof by Example

Take $\alpha = (4, 5, 2, 1, 2, 5) \in \mathrm{FR}_6 \cap \mathrm{UPF}_6$. Then, $\alpha^{\uparrow} = (1, \boxed{2, 2}, 4, \boxed{5, 5})$.

- Cars not in a box park in their preferred spot.
- First car in each box parks in its preferred spot.
- Second car in the same box parks in spot right after its preferred spot.

Proof of Main Result

Lemma ($\mathrm{FR}_n \cap \mathrm{UPF}_n$ Permutation Invariance)

For all $n \geq 1$, the set $\mathrm{FR}_n \cap \mathrm{UPF}_n$ is permutation invariant.

But wait! The set UPF_n is not permutation invariant.

Proof by Example

Take $\alpha = (4, 5, 2, 1, 2, 5) \in \mathrm{FR}_6 \cap \mathrm{UPF}_6$. Then, $\alpha^{\uparrow} = (1, \boxed{2, 2}, 4, \boxed{5, 5})$.

- Cars not in a box park in their preferred spot.
- First car in each box parks in its preferred spot.
- Second car in the same box parks in spot right after its preferred spot.
- Entries in a box are indistinguishable and can be permuted.

Proof of Main Result





Definition (rev(α))

Let $\alpha = (a_1, a_2, \ldots, a_n)$. Define rev($\alpha$) to reverse the order of the entries in α such that rev(α) $= (a_n, \ldots, a_2, a_1)$.

Proof of Main Result




Definition (rev(α))

Let $\alpha = (a_1, a_2, \ldots, a_n)$. Define rev($\alpha$) to reverse the order of the entries in α such that rev(α) $= (a_n, \ldots, a_2, a_1)$.

Corollary

For all $n \geq 1$, rev($\alpha^{\uparrow}$) $= \alpha^{\downarrow}$ and rev($\alpha^{\downarrow}$) $= \alpha^{\uparrow}$.

Proof of Main Result

Definition (rev(α))

Let $\alpha = (a_1, a_2, \ldots, a_n)$. Define $\mathrm{rev}(\alpha)$ to reverse the order of the entries in α such that $\mathrm{rev}(\alpha) = (a_n, \ldots, a_2, a_1)$.

Corollary

For all $n \geq 1$, $\mathrm{rev}(\alpha^{\uparrow}) = \alpha^{\downarrow}$ and $\mathrm{rev}(\alpha^{\downarrow}) = \alpha^{\uparrow}$.

Lemma

For all $n \geq 1$, $\alpha \in \mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}$ if and only if $\mathrm{rev}(\alpha) \in \mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}$.

Proof of Main Result

Definition (rev(α))

Let $\alpha = (a_1, a_2, \ldots, a_n)$. Define $\text{rev}(\alpha)$ to reverse the order of the entries in α such that $\text{rev}(\alpha) = (a_n, \ldots, a_2, a_1)$.

Corollary

For all $n \geq 1$, $\text{rev}(\alpha^{\uparrow}) = \alpha^{\downarrow}$ and $\text{rev}(\alpha^{\downarrow}) = \alpha^{\uparrow}$.

Lemma

For all $n \geq 1$, $\alpha \in \text{FR}_n^{\downarrow} \cap \text{UPF}_n^{\downarrow}$ if and only if $\text{rev}(\alpha) \in \text{FR}_n^{\uparrow} \cap \text{UPF}_n^{\uparrow}$.

Thus, $|\text{FR}_n^{\uparrow} \cap \text{UPF}_n^{\uparrow}| = |\text{FR}_n^{\downarrow} \cap \text{UPF}_n^{\downarrow}|$.

Proof of Main Result





Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Proof of Main Result

Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Now, we want $|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}|$.

Proof of Main Result




Lemma

For all $n \geq 1$, $\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow} = \mathrm{UPF}_n^{\downarrow}$.

Proof of Main Result





Lemma

For all $n \geq 1$, $\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow} = \mathrm{UPF}_n^{\downarrow}$.

Proof by Example (Contrapositive)

- We already saw $(2, 1, 1) \notin \mathrm{FR}_3^{\downarrow} \cup \mathrm{UPF}_3^{\downarrow}$.

Proof of Main Result

Lemma

For all $n \geq 1$, $\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow} = \mathrm{UPF}_n^{\downarrow}$.

Proof by Example (Contrapositive)

- We already saw $(2, 1, 1) \notin \mathrm{FR}_3^{\downarrow} \cup \mathrm{UPF}_3^{\downarrow}$.
- Consider $(3, 3, 1) \notin \mathrm{FR}_3^{\downarrow}$. Car 2 fails to park, and $\alpha \notin \mathrm{UPF}_3^{\downarrow}$.

Proof of Main Result

Lemma

For all $n \geq 1$, $\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow} = \mathrm{UPF}_n^{\downarrow}$.

Proof by Example (Contrapositive)

- We already saw $(2,1,1) \notin \mathrm{FR}_3^{\downarrow} \cup \mathrm{UPF}_3^{\downarrow}$.
- Consider $(3,3,1) \notin \mathrm{FR}_3^{\downarrow}$. Car 2 fails to park, and $\alpha \notin \mathrm{UPF}_3^{\downarrow}$.
- Consider $(3,1,1) \in \mathrm{FR}_3^{\downarrow}$. All cars park, so $\alpha \in \mathrm{UPF}_3^{\downarrow}$.



Proof of Main Result

Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

Proof of Main Result

Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

We already know the cardinality is F_{n+1} [3, Elder et al., 2025].

Proof of Main Result





Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

We already know the cardinality is F_{n+1} [3, Elder et al., 2025].

Question: While recursion is nice, can we find a closed formula?

Proof of Main Result





Theorem

For all $n \geq 1$,

$$|\mathrm{FR}_n^{\uparrow} \cap \mathrm{UPF}_n^{\uparrow}| = |\mathrm{FR}_n^{\downarrow} \cap \mathrm{UPF}_n^{\downarrow}| = |\mathrm{UPF}_n^{\downarrow}| = F_{n+1}$$

where F_{n+1} is the $(n+1)$th Fibonacci number. The terms for $1 \leq n \leq 7$ are:

$$1, 2, 3, 5, 8, 13, 21.$$

We already know the cardinality is F_{n+1} [3, Elder et al., 2025].

Question: While recursion is nice, can we find a closed formula?

Yes!

Proof of F_{n+1} Enumeration





Lemma

For all $n \geq 1$, if $\alpha = (a_1, a_2, \ldots, a_n)$, then $k \leq \lfloor \frac{n}{2} \rfloor$ entries $i \in \alpha$ can repeat once.

Proof of F_{n+1} Enumeration





Lemma

For all $n \geq 1$, if $\alpha = (a_1, a_2, \ldots, a_n)$, then $k \leq \lfloor \frac{n}{2} \rfloor$ entries $i \in \alpha$ can repeat once.

Lemma

For $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n \cup \mathrm{UPF}_n$, if entries $i, j \in \alpha$ are repeats, they are not consecutive numbers.

Proof of F_{n+1} Enumeration

Lemma

For all $n \geq 1$, if $\alpha = (a_1, a_2, \ldots, a_n)$, then $k \leq \lfloor \frac{n}{2} \rfloor$ entries $i \in \alpha$ can repeat once.

Lemma

For $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n \cup \mathrm{UPF}_n$, if entries $i, j \in \alpha$ are repeats, they are not consecutive numbers.

Example

Consider $(1, 1, 2, 2) \notin \mathrm{FR}_4 \cup \mathrm{UPF}_4$ due to invalid entries and parking conflicts.

Proof of F_{n+1} Enumeration




Proof

Given $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n \cap \mathrm{UPF}_n$, choose $k \leq \lfloor \frac{n}{2} \rfloor$ distinct entries $i \in \alpha$ to repeat from the $n - k$ options (subtract the numbers greater than and consecutive to each i).

Proof of F_{n+1} Enumeration





Proof

Given $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n \cap \mathrm{UPF}_n$, choose $k \leq \lfloor \frac{n}{2} \rfloor$ distinct entries $i \in \alpha$ to repeat from the $n - k$ options (subtract the numbers greater than and consecutive to each i). Counting the variations, we have

$$\sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k}.$$

Proof of F_{n+1} Enumeration





Proof

Given $\alpha = (a_1, a_2, \ldots, a_n) \in \mathrm{FR}_n \cap \mathrm{UPF}_n$, choose $k \leq \lfloor \frac{n}{2} \rfloor$ distinct entries $i \in \alpha$ to repeat from the $n - k$ options (subtract the numbers greater than and consecutive to each i). Counting the variations, we have

$$\sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k}.$$

For $1 \leq n \leq 7$, the summation yields

$$1, 2, 3, 5, 8, 13, 21,$$

which are the $(n+1)$th Fibonacci numbers, F_{n+1}.

Example of F_{n+1} Enumeration





Example

Determine $|\mathrm{FR}_4^{\downarrow} \cap \mathrm{UPF}_4^{\downarrow}| = |\mathrm{UPF}_4^{\downarrow}|$.

Example of F_{n+1} Enumeration



Example

Determine $|\mathrm{FR}_4^{\downarrow} \cap \mathrm{UPF}_4^{\downarrow}| = |\mathrm{UPF}_4^{\downarrow}|$. We have

$$\sum_{k=0}^{\lfloor \frac{4}{2} \rfloor} \binom{4-k}{k} = \underbrace{\binom{4-0}{0}}_{1} + \underbrace{\binom{4-1}{1}}_{3} + \underbrace{\binom{4-2}{2}}_{1} = 5.$$

Example of F_{n+1} Enumeration

Example

Determine $|\mathrm{FR}_4^{\downarrow} \cap \mathrm{UPF}_4^{\downarrow}| = |\mathrm{UPF}_4^{\downarrow}|$. We have

$$\sum_{k=0}^{\lfloor \frac{4}{2} \rfloor} \binom{4-k}{k} = \underbrace{\binom{4-0}{0}}_{1} + \underbrace{\binom{4-1}{1}}_{3} + \underbrace{\binom{4-2}{2}}_{1} = 5.$$

$$\mathrm{FR}_4^{\downarrow} \cap \mathrm{UPF}_4^{\downarrow} = \mathrm{UPF}_4^{\downarrow} \begin{cases} (4,3,2,1) & (k = 0 \text{ repeats}) \\ (4,3,1,1) & (k = 1 \text{ repeat}) \\ (4,2,2,1) & (k = 1 \text{ repeat}) \\ (3,3,2,1) & (k = 1 \text{ repeat}) \\ (3,3,1,1) & (k = 2 \text{ repeats}). \end{cases}$$

Summary




- Definitions
  - Tuples and Weak Order
  - Fubini Rankings
  - Unit interval parking functions
  - Unit Fubini Rankings
- Motivation
- Main Result (Sets Enumerated by F_{n+1})

Future Work

- Given $\alpha \in \mathrm{FR}_n$, park the cars. If a car's preferred spot is occupied, the car will continue down the street until it finds an empty spot to park.
- Given $\alpha \in \mathrm{UPF}_n$, follow the parking rules stated in the definition.

Future Work

- Given $\alpha \in \mathrm{FR}_n$, park the cars. If a car's preferred spot is occupied, the car will continue down the street until it finds an empty spot to park.
- Given $\alpha \in \mathrm{UPF}_n$, follow the parking rules stated in the definition.

Definition (Total Displacement)

Let $\alpha \in \mathrm{FR}_n \cup \mathrm{UPF}_n$. The total number of spots each car parked away from its preferred spot is called the *total displacement* of α, denoted $d(\alpha)$.

Future Work





Prove the following:

Future Work




Prove the following:

Proposition (Total displacement of UPF_n)

Let $\alpha \in \mathrm{UPF}_n$. Then $0 \leq d(\alpha) \leq n-1$.

Future Work

Prove the following:

Proposition (Total displacement of UPF_n)

Let $\alpha \in \mathrm{UPF}_n$. Then $0 \leq d(\alpha) \leq n-1$.

Proposition (Total displacement of $\mathrm{FR}_n \cap \mathrm{UPF}_n$)

Let $\alpha \in \mathrm{FR}_n \cap \mathrm{UPF}_n$. Then $0 \leq d(\alpha) \leq \lfloor \frac{n}{2} \rfloor$.

References

[1] Bjorn Andreas Ager-Hart et al.
Counting ℓ-interval Fubini rankings through their parking outcome. 2026. arXiv: 2603.00693 [math.CO]. URL: https://arxiv.org/abs/2603.00693.

[2] S. Alex Bradt et al. "Unit interval parking functions and the r-Fubini numbers". In: Matematica 3.1 (2024), pp. 370–384. ISSN: 2730-9657. DOI: 10.1007/s44007-024-00089-y. URL: https://doi.org/10.1007/s44007-024-00089-y.

[3] Jennifer Elder et al. "Parking functions, Fubini rankings, and Boolean intervals in the weak order of $\mathfrak{S}_n$". In: Journal of Combinatorics 16.1 (2025), pp. 65–89. ISSN: 2150-959X. DOI: 10.4310/joc.241216213152. URL: http://dx.doi.org/10.4310/JOC.241216213152.

Acknowledgments

- National Science Foundation (NSF) for funding the Combinatorics and Coding Theory in the Tropics REU
- University of Puerto Rico at Ponce for hosting the Combinatorics and Coding Theory in the Tropics REU




Dr. Pamela Harris

Principal Investigator



Dr. Fernando Piñero

Professional Mentor



Cyrus Young

Graduate Mentor



Melissa Beerbower

Graduate Mentor

Questions and Contact







Eban Barrett

emb1295@uah.edu

Maritza Sanchez

sanchezm36@southernct.edu

Tung (TJ) Vo

vo.tung@stthom.edu