

A Brief History of Generative Models for Power Law and Lognormal Distributions

Michael Mizenmacher

Presenter: Bo Jiang

Sept 17, 2010

Basic Definitions

Power law distribution

$$\bar{F}_X(x) = \mathbb{P}\{X > x\} \sim cx^{-\alpha} \quad \text{for } c, \alpha > 0$$

Basic Definitions

Power law distribution

$$\bar{F}_X(x) = \mathbb{P}\{X > x\} \sim cx^{-\alpha} \quad \text{for } c, \alpha > 0$$

Pareto distribution

$$\bar{F}_X(x) = \mathbb{P}\{X > x\} = \left(\frac{x}{k}\right)^{-\alpha} \quad \text{for } k, \alpha > 0$$

Basic Definitions

Power law distribution

$$\bar{F}_X(x) = \mathbb{P}\{X > x\} \sim cx^{-\alpha} \quad \text{for } c, \alpha > 0$$

Pareto distribution


$$\bar{F}_X(x) = \mathbb{P}\{X > x\} = \left(\frac{x}{k}\right)^{-\alpha} \quad \text{for } k, \alpha > 0$$

Lognormal distribution


$$\log X \sim \mathcal{N}(\mu, \sigma^2), \quad f(x) = \frac{1}{\sqrt{2\pi}\sigma x} e^{-(\log x - \mu)^2 / 2\sigma^2}$$

If $\sigma \gg 1$, density approx. linear for large range in log-log scale.

Power Law vs Lognormal



Power Law vs Lognormal



Preferential Attachment

Prototype model

- ▶ Start with single node with one self-loop.
- ▶ At each time step, add one new node of outdegree 1
 - ▶ $\alpha < 1$.
 - ▶ With probability α , link to node uniformly at random.
 - ▶ With probability $1 - \alpha$, link to node \propto *indegree*.

Preferential Attachment

Prototype model

- ▶ Start with single node with one self-loop.
- ▶ At each time step, add one new node of outdegree 1
 - ▶ $\alpha < 1$.
 - ▶ With probability α , link to node uniformly at random.
 - ▶ With probability $1 - \alpha$, link to node $\propto \text{indegree}$.

Indegree Distribution

- ▶ CCDF: $\bar{F}(k) \sim ck^{-1/(1-\alpha)}$, power law
- ▶ Simple mean field analysis.
- ▶ Can be justified using martingales.

Preferential Attachment (cont'd)

Mean field analysis

- ▶ $X_j(t)$: # of nodes of indegree j at time t
- ▶ At time t , total # of nodes = sum of indegrees = t , i.e.

$$\sum_j X_j(t) = \sum_j jX_j(t) = t$$

Preferential Attachment (cont'd)

Mean field analysis

- ▶ $X_j(t)$: # of nodes of indegree j at time t
- ▶ At time t , total # of nodes = sum of indegrees = t , i.e.

$$\sum_j X_j(t) = \sum_j jX_j(t) = t$$

- ▶ For $j \geq 1$,

$$\frac{d}{dt}X_j = \alpha \frac{X_{j-1}}{t} + (1 - \alpha) \frac{(j-1)X_{j-1}}{t} + \alpha \frac{X_j}{t} + (1 - \alpha) \frac{jX_j}{t}$$

- ▶

$$\frac{d}{dt}X_0 = 1 - \alpha \frac{X_0}{t}$$

- ▶ Solve for $c_j = X_j/t$ in steady state, i.e. $t \rightarrow \infty$.

Optimization

Mandelbrot's model for word rank-frequency distribution

- ▶ Average amount of information per transmission:

$$\text{entropy: } H = - \sum_{j=1}^n p_j \log_2 p_j$$

n = total # of words.

Optimization

Mandelbrot's model for word rank-frequency distribution

- ▶ Average amount of information per transmission:

$$\text{entropy: } H = - \sum_{j=1}^n p_j \log_2 p_j$$

n = total # of words.

- ▶ Optimize average amount of information per unit cost

$$\min A = \frac{C}{H} = \frac{\sum_{j=1}^n p_j C_j}{H} \implies p_j = e^{-1} 2^{-HC_j/C}$$

C_j cost for j th word.

Optimization

Mandelbrot's model for word rank-frequency distribution

- ▶ Average amount of information per transmission:

$$\text{entropy: } H = - \sum_{j=1}^n p_j \log_2 p_j$$

n = total # of words.

- ▶ Optimize average amount of information per unit cost

$$\min A = \frac{C}{H} = \frac{\sum_{j=1}^n p_j C_j}{H} \implies p_j = e^{-1} 2^{-HC_j/C}$$

C_j cost for j th word.



$$C_j \sim \log_d j \implies \text{power law for } p_j \sim e^{-1} j^{-H \log_d 2 / C}$$

d alphabet size.

Multiplicative Processes

Basic Operation

$$X_j = F_j X_{j-1} \quad \text{for r.v. } F_j$$

X_0, F_j 's independent.

Multiplicative Processes

Basic Operation

$$X_j = F_j X_{j-1} \quad \text{for r.v. } F_j$$

X_0, F_j 's independent.

Limit Distribution

$$\log X_j = \log X_0 + \sum_{k=1}^j \log F_k$$

- ▶ F_j 's independent lognormal $\implies X_j$ lognormal $\forall j$
- ▶ In general, CLT $\implies X_j$ converges to lognormal

Multiplicative Processes

Basic Operation

$$X_j = F_j X_{j-1} \quad \text{for r.v. } F_j$$

X_0, F_j 's independent.

Limit Distribution


$$\log X_j = \log X_0 + \sum_{k=1}^j \log F_k$$

- ▶ F_j 's independent lognormal $\implies X_j$ lognormal $\forall j$
- ▶ In general, CLT $\implies X_j$ converges to lognormal

However, seemingly trivial modifications can lead to power law!


Multiplicative Processes (cont'd)

- ▶ $X_0 = m$.
- ▶ $\mathbb{P}\{F_j = \gamma\} = 1/(1 + \gamma), \mathbb{P}\{F_j = \gamma^{-1}\} = \gamma/(1 + \gamma), \gamma > 1$
- ▶ X_j lognormal




Multiplicative Processes (cont'd)

- ▶ $X_0 = m$.
- ▶ $\mathbb{P}\{F_j = \gamma\} = 1/(1 + \gamma), \mathbb{P}\{F_j = \gamma^{-1}\} = \gamma/(1 + \gamma), \gamma > 1$
- ▶ X_j lognormal



- ▶ But, if $X_j = \max\{F_j X_{j-1}, m\} \implies X_j$ power law!



Monkey Typing Randomly

The model

- ▶ Alphabet $\mathcal{X} = \{x_1, \dots, x_n\}$ plus space
- ▶ Hit space with probability q
- ▶ Hit x_j with probability $q_j = (1 - q)p_j$, where $\sum_j p_j = 1$
- ▶ Space separates words

Monkey Typing Randomly

The model

- ▶ Alphabet $\mathcal{X} = \{x_1, \dots, x_n\}$ plus space
- ▶ Hit space with probability q
- ▶ Hit x_j with probability $q_j = (1 - q)p_j$, where $\sum_j p_j = 1$
- ▶ Space separates words

Frequency-rank Distributions

uniform $p_j = \frac{1}{n} \implies$ power law

Monkey Typing Randomly

The model

- ▶ Alphabet $\mathcal{X} = \{x_1, \dots, x_n\}$ plus space
- ▶ Hit space with probability q
- ▶ Hit x_j with probability $q_j = (1 - q)p_j$, where $\sum_j p_j = 1$
- ▶ Space separates words

Frequency-rank Distributions

uniform $p_j = \frac{1}{n} \implies$ **power law**

- ▶ length: $L(w)$
- ▶ frequency: $f(w) \approx \mathbb{P}(\{w\}) = \left(\frac{1-q}{n}\right)^{L(w)} q$

Monkey Typing Randomly

The model

- ▶ Alphabet $\mathcal{X} = \{x_1, \dots, x_n\}$ plus space
- ▶ Hit space with probability q
- ▶ Hit x_j with probability $q_j = (1 - q)p_j$, where $\sum_j p_j = 1$
- ▶ Space separates words

Frequency-rank Distributions

uniform $p_j = \frac{1}{n} \implies$ **power law**

- ▶ length: $L(w)$
- ▶ frequency: $f(w) \approx \mathbb{P}(\{w\}) = \left(\frac{1-q}{n}\right)^{L(w)} q$
- ▶ rank: $r(w) \approx n^{L(w)-1} \sim n^{L(w)} \implies L(w) \approx \log_n r(w)$

Monkey Typing Randomly

The model

- ▶ Alphabet $\mathcal{X} = \{x_1, \dots, x_n\}$ plus space
- ▶ Hit space with probability q
- ▶ Hit x_j with probability $q_j = (1 - q)p_j$, where $\sum_j p_j = 1$
- ▶ Space separates words

Frequency-rank Distributions

uniform $p_j = \frac{1}{n} \implies$ **power law**

- ▶ length: $L(w)$
- ▶ frequency: $f(w) \approx \mathbb{P}(\{w\}) = \left(\frac{1-q}{n}\right)^{L(w)} q$
- ▶ rank: $r(w) \approx n^{L(w)-1} \sim n^{L(w)} \implies L(w) \approx \log_n r(w)$

$$f \approx q r^{\log_n(1-q)-1}$$

Monkey Typing Randomly (cont'd)

Distributions

non-uniform p_j : $\exists i \neq j$ s.t. $p_i \neq p_j$

- ▶ Ω_m = set of all distinct words of length m , $|\Omega_m| = n^m$
- ▶ $\Omega = \bigcup_{m=0}^{\infty} \Omega_m$ = set of all distinct words

Monkey Typing Randomly (cont'd)

Distributions

non-uniform p_j : $\exists i \neq j$ s.t. $p_i \neq p_j$

- ▶ Ω_m = set of all distinct words of length m , $|\Omega_m| = n^m$
- ▶ $\Omega = \bigcup_{m=0}^{\infty} \Omega_m$ = set of all distinct words
- ▶ $C_i(w) = i$ -th letter of w , $i = 1, 2, \dots, L(w)$

Monkey Typing Randomly (cont'd)

Distributions

non-uniform p_j : $\exists i \neq j$ s.t. $p_i \neq p_j$

- ▶ Ω_m = set of all distinct words of length m , $|\Omega_m| = n^m$
- ▶ $\Omega = \bigcup_{m=0}^{\infty} \Omega_m$ = set of all distinct words
- ▶ $C_i(w) = i$ -th letter of w , $i = 1, 2, \dots, L(w)$
- ▶ frequency of w ,

$$f(w) = \mathbb{P}(\{w\}) = q \prod_{i=1}^{L(w)} q_{C_i(w)}$$

Monkey Typing Randomly (cont'd)

- ▶ Let \mathbb{Q}_m be uniform distribution on Ω_m

Monkey Typing Randomly (cont'd)

- ▶ Let \mathbb{Q}_m be uniform distribution on Ω_m
- ▶ As r.v. on Ω_m , $f : \Omega_m \rightarrow [0, 1]$ has CCDF \bar{G}_m w.r.t. \mathbb{Q}_m ,

Monkey Typing Randomly (cont'd)

- ▶ Let \mathbb{Q}_m be uniform distribution on Ω_m
- ▶ As r.v. on Ω_m , $f : \Omega_m \rightarrow [0, 1]$ has CCDF \bar{G}_m w.r.t. \mathbb{Q}_m ,
- ▶ Rank of w within Ω_m

$$\begin{aligned}r_m(w) &\approx \#\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \mathbb{Q}_m\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \bar{G}_m(f(w))\end{aligned}$$

Monkey Typing Randomly (cont'd)

- ▶ Let \mathbb{Q}_m be uniform distribution on Ω_m
- ▶ As r.v. on Ω_m , $f : \Omega_m \rightarrow [0, 1]$ has CCDF \bar{G}_m w.r.t. \mathbb{Q}_m ,
- ▶ Rank of w within Ω_m

$$\begin{aligned}r_m(w) &\approx \#\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \mathbb{Q}_m\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \bar{G}_m(f(w))\end{aligned}$$

- ▶ CLT $\implies \bar{G}_m$ is roughly lognormal
For fixed length, frequency-rank distribution is lognormal

Monkey Typing Randomly (cont'd)

- ▶ Let \mathbb{Q}_m be uniform distribution on Ω_m
- ▶ As r.v. on Ω_m , $f : \Omega_m \rightarrow [0, 1]$ has CCDF \bar{G}_m w.r.t. \mathbb{Q}_m ,
- ▶ Rank of w within Ω_m

$$\begin{aligned}r_m(w) &\approx \#\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \mathbb{Q}_m\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \bar{G}_m(f(w))\end{aligned}$$

- ▶ CLT $\implies \bar{G}_m$ is roughly lognormal
For fixed length, frequency-rank distribution is lognormal
- ▶ Upper bound word length, Anscombe's theorem \implies
lognormal frequency-rank distribution

Monkey Typing Randomly (cont'd)

- ▶ Let \mathbb{Q}_m be uniform distribution on Ω_m
- ▶ As r.v. on Ω_m , $f : \Omega_m \rightarrow [0, 1]$ has CCDF \bar{G}_m w.r.t. \mathbb{Q}_m ,
- ▶ Rank of w within Ω_m

$$\begin{aligned}r_m(w) &\approx \#\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \mathbb{Q}_m\{w' \in \Omega_m : f(w') > f(w)\} \\&= |\Omega_m| \bar{G}_m(f(w))\end{aligned}$$

- ▶ CLT $\implies \bar{G}_m$ is roughly lognormal
For fixed length, frequency-rank distribution is lognormal
- ▶ Upper bound word length, Anscombe's theorem \implies
lognormal frequency-rank distribution
- ▶ No bound on word length \implies power law (very complicated proof involving analytical number theory)

Double Pareto Distribution

- ▶ For $t \geq 0$, let $X(t)$ be lognormal, i.e.

$$\log X(t) \sim \mathcal{N}(\log x_0 + \mu t, \sigma^2 t)$$

where $x_0 > 0$, $\mu \in \mathbb{R}$, $\sigma \in \mathbb{R}^+$.

Double Pareto Distribution

- ▶ For $t \geq 0$, let $X(t)$ be lognormal, i.e.

$$\log X(t) \sim \mathcal{N}(\log x_0 + \mu t, \sigma^2 t)$$

where $x_0 > 0$, $\mu \in \mathbb{R}$, $\sigma \in \mathbb{R}^+$.

- ▶ Let $T \sim \exp(\lambda)$

Double Pareto Distribution

- ▶ For $t \geq 0$, let $X(t)$ be lognormal, i.e.

$$\log X(t) \sim \mathcal{N}(\log x_0 + \mu t, \sigma^2 t)$$

where $x_0 > 0$, $\mu \in \mathbb{R}$, $\sigma \in \mathbb{R}^+$.

- ▶ Let $T \sim \exp(\lambda)$
- ▶ Distribution of $Y = X(T)$?


Double Pareto Distribution

- ▶ For $t \geq 0$, let $X(t)$ be lognormal, i.e.

$$\log X(t) \sim \mathcal{N}(\log x_0 + \mu t, \sigma^2 t)$$

where $x_0 > 0$, $\mu \in \mathbb{R}$, $\sigma \in \mathbb{R}^+$.

- ▶ Let $T \sim \exp(\lambda)$
- ▶ Distribution of $Y = X(T)$? In log-log scale,



Double Pareto Distribution

- ▶ If population grows exponentially, age distribution will be truncated exponential.

Double Pareto Distribution

- ▶ If population grows exponentially, age distribution will be truncated exponential.
- ▶ $X(t)$ is geometric Brownian motion

$$\dot{X} = (\mu + \xi)X \quad \text{or} \quad dX(t) = X(\mu dt + \sigma dW)$$

ξ white noise, W standard Brownian motion.

- ▶ “noisy” exponential growth
- ▶ “continuous” multiplication

Double Pareto Distribution

- ▶ If population grows exponentially, age distribution will be truncated exponential.
- ▶ $X(t)$ is geometric Brownian motion

$$\dot{X} = (\mu + \xi)X \quad \text{or} \quad dX(t) = X(\mu dt + \sigma dW)$$

ξ white noise, W standard Brownian motion.

- ▶ “noisy” exponential growth
 - ▶ “continuous” multiplication
- ▶ If T is geometric: lognormal body, power law tail

$$X(T) = x_0 \prod_{i=1}^T X_i, \quad \text{where } X_i \text{ i.i.d } \sim \text{log-normal}(\mu, \sigma)$$

Double Pareto Distribution

- ▶ If population grows exponentially, age distribution will be truncated exponential.
- ▶ $X(t)$ is geometric Brownian motion

$$\dot{X} = (\mu + \xi)X \quad \text{or} \quad dX(t) = X(\mu dt + \sigma dW)$$


ξ white noise, W standard Brownian motion.

- ▶ “noisy” exponential growth
 - ▶ “continuous” multiplication
- ▶ If T is geometric: lognormal body, power law tail

$$X(T) = x_0 \prod_{i=1}^T X_i, \quad \text{where } X_i \text{ i.i.d } \sim \text{log-normal}(\mu, \sigma)$$

- ▶ Exponential growth + exponential (geometric) stopping
 \implies power law

Double Pareto Degree Distribution?



Degree distribution of Facebook [Gjoka et al 2010]