

NOTES FOR MACAULAY2 REAL PROJECT

JORDY LOPEZ, KELLY MALUCCIO, THOMAS J. YAHL, AND FRANK SOTTILE

ABSTRACT. These are notes around the project on implementing routines to study real roots of polynomial systems in Macaulay2.

1. INTRODUCTION

For a sequence $c = (c_0, \dots, c_n)$ of real numbers, let $\text{var}(c)$ be the variation (number of changes in sign) in the sequence c . This is the number of times consecutive elements of the sequence c have opposite signs, after removing any occurrences of 0 from c . Given a sequence $F = (f_0(t), \dots, f_m(t))$ of real univariate polynomials and $a \in \mathbb{R}$, let $\text{var}(F, a)$ be the variation in the sequence $(f_0(a), \dots, f_m(a))$. We define $\text{var}(F, \infty)$ to be the variation in the leading coefficients of the polynomials in F , which is $\text{var}(F, a)$ for $a \gg 0$ sufficiently positive, and set $\text{var}(F, -\infty)$ to be $\text{var}(F, a)$ for $a \ll 0$ sufficiently negative.

For a univariate polynomial $f \in \mathbb{R}[t]$ of degree m , its derivative sequence δf is the sequence of its derivatives, $\delta f := (f(t), f'(t), f''(t), f^{(3)}(t), \dots, f^{(m)}(t))$. For numbers $a < b$ in $\mathbb{R} \cup \{\pm\infty\}$, let $r(f, a, b)$ be the number of roots of f in the interval $(a, b]$.

Theorem 1.1 (Budan-Fourier Theorem). *Let $f \in \mathbb{R}[t]$ be a univariate polynomial and $a < b$ be numbers in $\mathbb{R} \cup \{\pm\infty\}$. Then $\text{var}(\delta f, a) - \text{var}(\delta f, b) \geq r(f, a, b)$, and the difference is even.*

(This implies Descartes' rule of signs for roots in $(0, \infty)$.)

Given univariate polynomials f, g , their *Sturm sequence* $\text{St}(f, g)$ is the sequence

$$f_0 := f, f_1 := f'g, f_2, f_3, \dots, f_m,$$

where f_m is a greatest common divisor of f and $f'g$, and for each $i \geq 1$, $-f_{i+1}$ is the negative of the remainder from the Euclidean algorithm. That is, f_{i+1} is the unique polynomial of degree less than the degree of f_i such that there is a polynomial q_i with $f_{i-1} = q_i f_i - f_{i+1}$.

Theorem 1.2 (Sylvester's Theorem). *Let $f, g \in \mathbb{R}[x]$ be univariate polynomials and suppose that $a < b$ are numbers in $\mathbb{R} \cup \{\pm\infty\}$ such that neither is a root of f . Then $\text{var}(\text{St}(f, g), a) - \text{var}(\text{St}(f, g), b)$ is the difference of the number of roots $x \in (a, b)$ of f with $g(x) > 0$ and those with $g(x) < 0$.*

Theorem 1.3 (Sturm's Theorem). *Let $f \in \mathbb{R}[x]$ be a univariate polynomial suppose that $a < b$ are two numbers in $\mathbb{R} \cup \{\pm\infty\}$ such that neither is a root of f . Then $\text{var}(\text{St}(f, 1), a) - \text{var}(\text{St}(f, 1), b)$ is the number of roots of f in the open interval (a, b) .*

1991 *Mathematics Subject Classification.* 14P99.

Lemma 1.4. *Let $f = c_0 + c_1t + \cdots + c_mt^m$ be a univariate polynomial and $M := |c_0/c_m| + \cdots + |c_{m-1}/c_m| + 1$. Then all roots of f lie in the interval $(-M, M)$.*

Proof. Let $|x| > M$. If $b_i := c_i/c_m$, then $f(x) = c_mx^m(b_0x^{-m} + \cdots + b_{m-1}x^{-1} + 1)$. As

$$|b_0x^{-m} + \cdots + b_{m-1}x^{-1}| < (|b_0| + \cdots + |b_{m-1}|)M^{-1} < 1,$$

we see that $f(x) \neq 0$. □

This gives the start of a bisection algorithm for locating the roots of f .

Email address: `jordy.lopez@math.tamu.edu`

Email address: `kmaluccio@math.tamu.edu`

Email address: `thomasjyahl@math.tamu.edu`

Email address: `sottile@math.tamu.edu`

URL: `www.math.tamu.edu/~sottile`