



E Message length and field occupancy are different quantities

Representation-level constants one value per map, set by the fixed-width ±d.dddddd serialiser and the tokenizer. Point values, not distributions.



Field-dependent quantity how sparse the physical field is, counted in bins, not characters. It does not change message length: both bin encodings print all 24 rows.



Across the 48 stored replay fields (16 peers each) 1–14 of the 24 bins print non-zero mass (median 11). The example field of panel A is denser: 16 of its 24 bins print non-zero mass, and the remaining 8 print 0.000000 only because their mass falls below 5×10^{-7} and rounds to zero at six decimals; every one of the 24 bins holds mass, none is empty. Occupancy is what the field controls; the message length above is not.

F Retained-feature inventory: the maps differ in content, not only in format

	moments	centers	intervals	what the difference does
First harmonic $m=1$ (polar order r_1, ψ_1)	✓	○	○	printed outright by moments; a 24-term sum for the other two
Second and third harmonics $m=2,3$	✓	○	○	the same three complex numbers, arrived at differently
Harmonics $m \geq 4$	✗	○	○	discarded by moments: 24 masses carry 12 harmonics, moments keep 3
Per-bin mass, all 24 bins	✗	✓	✓	local detail; identical numbers in centers and intervals
Explicit bin-centre angle	✗	✓	✗	intervals leaves the centre implicit as the interval midpoint
Explicit half-open bin edges	✗	✗	✓	only intervals states the boundary convention in the text
Six-decimal fixed-width presentation	✓	✓	✓	identical numeric precision, so precision cannot explain the gap
Peer count / absolute phase / agent identity	✗	✗	✗	never serialised by any map
✓ printed explicitly	○ recoverable from what is printed			✗ not present in the message