

TITAN research-software demonstration

24 August 2026

Contents

Research-software output: internally derived TCGA estimates	1
COAD - TCGA-AA-3972	1
Prominent tissue-source-site sensitivity warning	1
Case context (not used as model input)	2
Continuous internally derived estimates	4
Binary research-model calls	4
Binary development reliability	5
Quality and provenance	5
Endpoint interpretation and target sources	6

Research-software output: internally derived TCGA estimates

COAD - TCGA-AA-3972

Models applied	Continuous	Binary	Slides pooled
19	10	9	1

This research-software demonstration applies models developed and evaluated only in TCGA. Every displayed value is an **internally derived TCGA estimate**, not external performance evidence or a diagnostic output. Binary LDA scores are uncalibrated. Their displayed TCGA out-of-fold score ranks describe relative position only and must not be read as probabilities.

By default, TITANPred excludes binary models with fewer than 50 participants in either development class. Such models appear only after the caller explicitly sets `include_limited_evidence = TRUE`.

Prominent tissue-source-site sensitivity warning

12 model(s) in this output fell below their original effect threshold under TCGA tissue-source-site-grouped internal validation. These calls must not be highlighted without this warning. Tissue-source-site grouping is not institutional or scanner-level external validation.

Endpoint (type)	TSS-grouped metric	Grouped minus random	Warning category
APC (binary)	BA 0.543	-0.250	Near chance or worse
PI3K (binary)	BA 0.593	-0.020	Below effect threshold
Aneuploidy score (continuous)	Q2 0.173	-0.086	Below effect threshold
Deleted arm count (continuous)	Q2 0.147	-0.066	Below effect threshold
MANTIS score (continuous)	Q2 0.113	-0.104	Below effect threshold
MSIsensor score (continuous)	Q2 0.026	-0.229	Below effect threshold
Aneuploidy Score (continuous)	Q2 0.174	-0.096	Below effect threshold
Lymphocyte Infiltration Signature Score (continuous)	Q2 0.048	-0.238	Below effect threshold
Nonsilent Mutation Rate (continuous)	Q2 0.088	-0.127	Below effect threshold
SNV Neoantigens (continuous)	Q2 -0.013	-0.252	Near chance or worse
Silent Mutation Rate (continuous)	Q2 -0.003	-0.236	Near chance or worse
TIL Regional Fraction (continuous)	Q2 0.176	-0.278	Below effect threshold

Case context (not used as model input)

Shared selection features. Both cases were men with sigmoid-colon, moderately differentiated (G2), pT3 adenocarcinoma and tumour-free resection margins.

Pathology summary. The sigmoid colectomy contained an ulcerated grade-2 colorectal adenocarcinoma extending through the bowel wall into adjacent mesocolic fat (pT3). The proximal and distal resection margins were uninvolved; the supplied slide summary did not state a nodal category.


Treatment and follow-up. NCI GDC records list capecitabine and oxaliplatin beginning on day 60, with the recorded outcome ‘Progressive Disease’; recurrence was recorded on day 1216. Later records list fluorouracil, leucovorin, oxaliplatin and bevacizumab through day 1369, also with progressive disease. Follow-up on day 1551 was ‘with tumor’; the recorded vital status was alive.

Clinical-context source. Pathology was paraphrased from TCGA-Slide-Reports.csv distributed with the TITAN study (Ding et al., Nature Medicine 2025; doi:10.1038/s41591-025-03982-3). Treatment and follow-up fields were retrieved from the public NCI GDC Cases API (<https://api.gdc.cancer.gov/cases>) on 16 August 2026.

The pathology and treatment fields are descriptive public metadata; they were not predictors, training targets, or validation outcomes for TITANPred. No treatment-effect inference should be drawn from these examples.


Continuous internally derived TCGA estimates

[SITE-SENSITIVE] marks models below the original effect threshold under TCGA TSS grouping



Binary research-model calls

TCGA out-of-fold LDA-score rank; [SITE-SENSITIVE] models require prominent c



Continuous internally derived estimates

Family	Endpoint	Prediction	Reference percentile	Category
thorsson	Aneuploidy Score	15.76	91.5%	Moderate effect
microsatellite_instability	MANTIS score	0.2664	9.8%	Moderate effect
thorsson	Nonsilent Mutation Rate	1.152	15.5%	Moderate effect
thorsson	Silent Mutation Rate	0.5383	15.8%	Moderate effect
aneuploidy	Aneuploidy score	16.71	82.0%	Moderate effect
thorsson	Lymphocyte Infiltration Signature Score	-0.4594	22.4%	Moderate effect
thorsson	SNV Neoantigens	3.968	26.7%	Moderate effect
microsatellite_instability	MSIsensor score	1.183	28.8%	Moderate effect
aneuploidy	Deleted arm count	8.558	69.7%	Moderate effect
thorsson	TIL Regional Fraction	5.048	30.3%	Higher effect

Binary research-model calls

Calls and development denominators.

Endpoint	Class	LDA score	TCGA OOF score rank (not probability)	Training n (+/-)
Genome doubling	1	1.557	84.7%	408 (+165/-243)
APC	0	-3.918	4.3%	382 (+196/-186)
KRAS	0	-4.625	9.6%	382 (+126/-256)
TP53 [driver_mutation]	0	-3.37	15.1%	382 (+161/-221)
MSI-H (MANTIS >0.4)	0	-4.847	42.3%	391 (+74/-317)
MSI-H strict (MANTIS >0.6)	0	-9.961	23.7%	374 (+57/-317)
MYC	0	-2.331	36.1%	323 (+79/-244)
PI3K	0	-1.268	22.8%	323 (+148/-175)
TP53 [oncogenic_pathway]	1	1.99	77.2%	323 (+208/-115)

Held-out repeated-CV evidence. PPV and NPV use the observed TCGA prevalence.

Endpoint	BA	AUROC	PR-AUC	PPV	NPV	Evidence
Genome doubling	0.692	0.760	0.678	0.636	0.750	>=50 per class
APC	0.794	0.848	0.814	0.760	0.851	>=50 per class
KRAS	0.692	0.781	0.581	0.600	0.794	>=50 per class
TP53 [driver_mutation]	0.778	0.854	0.793	0.718	0.828	>=50 per class
MSI-H (MANTIS >0.4)	0.732	0.834	0.634	0.677	0.894	>=50 per class
MSI-H strict (MANTIS >0.6)	0.782	0.915	0.741	0.750	0.931	>=50 per class
MYC	0.605	0.676	0.417	0.515	0.801	>=50 per class
PI3K	0.607	0.640	0.583	0.582	0.635	>=50 per class
TP53 [oncogenic_pathway]	0.668	0.734	0.815	0.752	0.624	>=50 per class

Binary development reliability

Fold class counts and component selection.

Endpoint	Minimum outer test (+/-)	Minimum inner training (+/-)	Components: median (range)
Genome doubling	+33 / -48	+105 / -155	6.0 (4-10)
APC	+39 / -37	+124 / -118	2.0 (1-4)
KRAS	+25 / -51	+80 / -163	4.0 (2-10)
TP53 [driver_mutation]	+32 / -44	+102 / -140	2.0 (2-7)
MSI-H (MANTIS >0.4)	+14 / -63	+47 / -202	6.0 (4-9)
MSI-H strict (MANTIS >0.6)	+11 / -63	+36 / -202	7.0 (3-10)
MYC	+15 / -48	+50 / -156	4.0 (2-5)
PI3K	+29 / -35	+94 / -112	4.0 (2-10)
TP53 [oncogenic_pathway]	+41 / -23	+132 / -73	3.0 (1-6)

Repeat-to-repeat held-out prediction stability.

Endpoint	Score Spearman rho	Class agreement
Genome doubling	0.850	0.857
APC	0.945	0.944
KRAS	0.825	0.858
TP53 [driver_mutation]	0.948	0.931
MSI-H (MANTIS >0.4)	0.856	0.940
MSI-H strict (MANTIS >0.6)	0.804	0.948
MYC	0.861	0.905
PI3K	0.721	0.787
TP53 [oncogenic_pathway]	0.853	0.895

Quality and provenance

- Patient-level aggregation: arithmetic mean across all supplied slide vectors.
- Maximum fraction outside endpoint-specific TCGA training ranges: 0.4%.
- TITAN input schema: 768 named features (`titan_000`–`titan_767`).
- Decomposition: seeded rSVD in fastPLS; binary classification: PLS-LDA.

- Tissue-source-site metadata: every row includes the grouped internal metric, random-minus-grouped change, analysed-site count, threshold-retention status, fold-size range and inner-site-separation audit.
- Binary reliability metadata: held-out average precision, cohort-specific PPV/NPV, outer and inner class-count minima, selected-component distribution and repeat-to-repeat prediction stability.
- Default model selection: binary models require at least 50 TCGA participants in each class; 17 smaller-class models are available only by explicit opt-in.
- Scope: TCGA tissue-source-site-grouped internal validation is not institutional or scanner-level external validation.
- Intended use: research-software demonstration only; TCGA benchmark without independent external validation.

Endpoint interpretation and target sources

For continuous endpoints, the percentile locates the prediction within repeated TCGA out-of-fold predictions and is not a clinical reference interval. For binary endpoints, the displayed value is the rank of the uncalibrated LDA score among repeated TCGA out-of-fold scores. It is not a probability, confidence, or clinical-risk estimate. PR-AUC is non-interpolated average precision for positive class 1. PPV and NPV are held-out estimates at the observed endpoint prevalence in this TCGA analysis and will change when prevalence changes. Rows marked exploratory/limited evidence are excluded from default inference and require explicit opt-in.

- **Genome doubling.** Binary whole-genome-doubling call derived from the tumour copy-number profile. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* Taylor et al., Cancer Cell 2018, Table S2; doi:10.1016/j.ccell.2018.03.007.
- **APC.** Binary prediction of a qualifying protein-altering PASS mutation in APC; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* MC3 consensus calls: Ellrott et al., Cell Systems 2018, doi:10.1016/j.cels.2018.03.002; cancer-gene eligibility: Bailey et al., Cell 2018, doi:10.1016/j.cell.2018.02.060.
- **KRAS.** Binary prediction of a qualifying protein-altering PASS mutation in KRAS; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* MC3 consensus calls: Ellrott et al., Cell Systems 2018, doi:10.1016/j.cels.2018.03.002; cancer-gene eligibility: Bailey et al., Cell 2018, doi:10.1016/j.cell.2018.02.060.
- **TP53 [driver_mutation].** Binary prediction of a qualifying protein-altering PASS mutation in TP53; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* MC3 consensus calls: Ellrott et al., Cell Systems 2018, doi:10.1016/j.cels.2018.03.002; cancer-gene eligibility: Bailey et al., Cell 2018, doi:10.1016/j.cell.2018.02.060.
- **MSI-H (MANTIS >0.4).** Binary call indicating whether the source MANTIS score exceeds 0.4; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* Bonneville et al., JCO Precision Oncology 2017; doi:10.1200/PO.17.00073.
- **MSI-H strict (MANTIS >0.6).** Sensitivity-analysis call: MANTIS >0.6 is positive, <0.4 is negative, and intermediate source values were excluded from training. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* Threshold documentation distributed with the cBioPortal TCGA PanCancer Atlas clinical sample files.
- **MYC.** Binary prediction that the MYC oncogenic pathway carries at least one qualifying genomic alteration; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* Sanchez-Vega et al., Cell 2018, Table S4; doi:10.1016/j.cell.2018.03.035.
- **PI3K.** Binary prediction that the PI3K oncogenic pathway carries at least one qualifying genomic alteration; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* Sanchez-Vega et al., Cell 2018, Table S4; doi:10.1016/j.cell.2018.03.035.
- **TP53 [oncogenic_pathway].** Binary prediction that the TP53 oncogenic pathway carries at least one qualifying genomic alteration; the displayed LDA score is not a probability. *Analysed scale:* PLS-LDA class and uncalibrated discriminant score *Model-target source:* Sanchez-Vega et al., Cell 2018, Table S4; doi:10.1016/j.cell.2018.03.035.
- **Aneuploidy score.** Number of chromosome arms with arm-level copy-number gain or loss; larger values indicate greater chromosomal imbalance. *Analysed scale:* source endpoint units *Model-target source:* Taylor et al., Cancer Cell 2018, Table S2; doi:10.1016/j.ccell.2018.03.007.

- **Deleted arm count.** Number of chromosome arms classified as deleted; larger values indicate a greater arm-level deletion burden. *Analysed scale:* source endpoint units *Model-target source:* Taylor et al., Cancer Cell 2018, Table S2; doi:10.1016/j.ccell.2018.03.007.
- **MANTIS score.** Genome-wide microsatellite-instability score from the MANTIS algorithm; larger values indicate stronger MSI evidence. *Analysed scale:* source endpoint units *Model-target source:* Kautto et al., Oncotarget 2017; doi:10.18632/oncotarget.20432. TCGA values were obtained from cBioPortal PanCancer Atlas records.
- **MSIsensor score.** Microsatellite-instability score from MSIsensor; larger values indicate a larger fraction of unstable microsatellite loci. *Analysed scale:* source endpoint units *Model-target source:* Niu et al., Bioinformatics 2014; doi:10.1093/bioinformatics/btt755. TCGA values were obtained from cBioPortal PanCancer Atlas records.
- **Aneuploidy Score.** Aneuploidy burden distributed with the TCGA immune landscape resource; larger values indicate more arm-level copy-number alterations. *Analysed scale:* source endpoint units *Model-target source:* Thorsson et al., Immunity 2018, PanImmune_MS; doi:10.1016/j.immuni.2018.03.023.
- **Lymphocyte Infiltration Signature Score.** Gene-expression signature summarising lymphocyte-associated transcriptional infiltration; it is not a direct cell count. *Analysed scale:* source endpoint units *Model-target source:* Thorsson et al., Immunity 2018, PanImmune_MS; doi:10.1016/j.immuni.2018.03.023.
- **Nonsilent Mutation Rate.** Rate of coding mutations that alter the encoded protein. TITANPred reports the model output on the analysis $\log(1+x)$ scale. *Analysed scale:* $\log_{1p}$ -transformed source endpoint units *Model-target source:* Thorsson et al., Immunity 2018, PanImmune_MS; doi:10.1016/j.immuni.2018.03.023.
- **SNV Neoantigens.** Predicted neoantigen burden arising from single-nucleotide variants. TITANPred reports the model output on the analysis $\log(1+x)$ scale. *Analysed scale:* $\log_{1p}$ -transformed source endpoint units *Model-target source:* Thorsson et al., Immunity 2018, PanImmune_MS; doi:10.1016/j.immuni.2018.03.023.
- **Silent Mutation Rate.** Rate of coding mutations that do not change the encoded amino acid. TITANPred reports the model output on the analysis $\log(1+x)$ scale. *Analysed scale:* $\log_{1p}$ -transformed source endpoint units *Model-target source:* Thorsson et al., Immunity 2018, PanImmune_MS; doi:10.1016/j.immuni.2018.03.023.
- **TIL Regional Fraction.** Estimated fraction of tumour image regions containing tumour-infiltrating lymphocytes in the TCGA immune landscape resource. *Analysed scale:* source endpoint units *Model-target source:* Thorsson et al., Immunity 2018, PanImmune_MS; doi:10.1016/j.immuni.2018.03.023.