
Models of both families and the reference models, by AIC
Standard (resistance)
SAMC
Reference model
dor_pc_pva (inverse monomolecular, shape 1); resistance 1-145 per km
dor_pc_pva (inverse monomolecular, shape 1); resistance 1-194 per km
tmp_dc_cyr (reverse Ricker, shape 1) + dor_pc_pva (inverse Ricker, shape 7); resistance 1-93.5 per km
sgr_dk_rav (reverse-inverse monomolecular, shape 1, symmetric)
tmp_dc_cyr (reverse-inverse Ricker, shape 4); resistance 1-19.7 per km
tmp_dc_cyr (reverse-inverse Ricker, shape 3, symmetric) + flow -0.002/km (gene flow downstream)
SAMC flow alone (-0.100/km)
flow -0.082/km (gene flow downstream)
run_mm_cyr (Ricker, shape 8, symmetric) + flow -0.100/km (gene flow downstream)
SAMC distance alone
distance only (km; standard model)
population effects only (null)
w = 0.51
w = 0.49
ΔAIC = 2
0
2
5
10
20
50
100
AIC difference from the best model (symmetric log scale)
Summed Akaike weight
Standard (resistance)
1.00
SAMC
<0.01
Reference model
<0.01
0.0
0.5
1.0