Once-Daily Prokinetic Agent

GASTIIN® CR Tab.

Mosapride citrate **15 mg**

Drug information

Composition	Mosapride citrate 15mg
Mechanism of Action	Selective 5-HT_4 receptor agonist
Indication	Treatment of gastrointestinal symptoms associated with functional dyspepsia (heartburn, nausea, vomiting).
Dosage and Administration	Adults : 15 mg of mosapride citrate is orally administered once a day. GASTIIN CR should be taken on an empty stomach. Tablet must be swallowed without chewing or crushing.

For the Best Global Pharmaceutical Company
KOREA UNITED PHARM. INC.

GST1609PA1-HI

For the Best Global Pharmaceutical Company
KOREA UNITED PHARM. INC.

Once-Daily Prokinetic Agent

GASTIIN® CR Tab.

Mosapride citrate **15 mg**

Once Daily One Tablet, Prokinetics

GASTIIN® CR Tab.

With the **BILDAS** (Bilayered Drug Absorption System)*...

*Patent No. 1612931 (2016.04.08)

Once-daily prokinetic agent,
GASTIIN CR Tab. improves the drug compliance
of the patient with 24 hours of long-acting effect.

Reduced the risk of side effects as a selective 5-HT_4 receptor agonist.
Effective in all gastrointestinal tract from esophagus to stomach and intestines.
Little drug interactions and convenient once-daily dosing make it easy
to be co-administered with other drugs.

GASTIIN® CR Tab.

improves the drug compliance by controlled release for 24 hours with BILDAS (Bilayered Drug Absorption System) technology.

Once-Daily Prokinetic Agent
GASTIIN® CR Tab.
Mosapride citrate 15mg



Improvement of the drug compliance
(3 times a day -> once a day)

GASTIIN® CR Tab. has been proven for the efficacy and safety through phase 3 clinical trials where 19 medical centers across the country have participated.

GASTIIN® CR Tab. 15mg Phase 3 Clinical Trials

MARS (*MosARpride Study*)

Multi-center, Randomized, Active-controlled, Double-blind, Non-inferiority, Phase 3 Clinical Trial To Evaluate The Efficacy And Safety of GASTIIN CR Tab. in Functional Dyspepsia

Study Title	Multi-center, Randomized, Active-controlled, Double-blind, Non-inferiority, Phase 3 Clinical Trial To Evaluate The Efficacy And Safety of GASTIIN CR Tab. in Functional Dyspepsia
Period	November 5, 2014 ~ September 5, 2015 (10 months)
Participating Medical Center	19 medical centers
Treatment Profile	Functional dyspepsia
Outcome Measure	Primary outcome measures : Total Gastrointestinal system score (GIS)
Number of Subject	143
Study Design	Screening → 2 weeks Wash-out → Randomization → GASTIIN CR Tab. 15 mg Once daily PO / Mosapride citrate 5 mg 3 times a day PO (Total 4 weeks); V1 Informed consent, V2 (W0) Baseline, V3 (W2), V4 (W4) End of Study Primary outcome

Primary Outcome Measures : Total GIS(Gastrointestinal system score)

GIS Evaluation Method

The severity of 10 symptoms for nausea, vomiting, bloating, abdominal cramps, early satiety, acid eructation/heartburn, sickness, loss of appetite, retrosternal discomfort, and epigastric or upper abdominal pain experienced by the subject over the past 2 weeks is evaluated on the 5-point Likert scale of 0 (no symptom), 1 (mild), 2 (moderate), 3 (severe), 4 (very severe), and each symptom score is added up.

Total GIS per visit



Visit	GASTIIN CR	Reference Drug
Baseline	13.3	12.9
2 weeks	6.0	5.6
4 weeks	3.5	2.9

Changes of total GIS from baseline per visit



Visit	GASTIIN CR	Reference Drug	*P-value*
2 weeks	-7.2	-7.3	0.656
4 weeks	-9.8	-10.0	0.824

* Reference Drug : GASMOTIN Tab. 5 mg t.i.d

As the result of clinical trial where GASTIIN CR Tab. has been administered for 4 weeks on patients with functional dyspepsia, the efficacy and safety have been proved compared to the reference drug of mosapride immediate-release tablet.

Once-Daily Prokinetic Agent
GASTIIN® CR Tab.
Mosapride citrate 15mg

Secondary Outcome Measures : NDI-K (Nepean Dyspepsia Index-Korean Version) score

NDI-K Evaluation Method

The questions of quality of life consist of 25 questions, 5 areas, which include 9 questions on tension/sleep, 6 questions on interference with daily activities, 3 questions on eating/drinking, 4 questions on knowledge/control, and 3 questions of work/study. Each question is calculated from 1 to 5 points, and the higher the point, the better quality of life it indicates.

Total NDI-K score



Visit	GASTIIN CR	Reference Drug
Baseline	55.8	55.4
4 weeks	70.1	72.3

Change of total NDI-K score from Baseline



Visit	GASTIIN CR	Reference Drug	*P-value*
4 weeks	14.3	16.9	0.263

Secondary Outcome Measures : GIS per symptoms

2 weeks Baseline	GASTIIN CR N=59	Reference Drug N=58	*P-value*
Nausea	-0.92±1.29	-0.86±1.05	0.924
Vomiting	-0.58±1.02	-0.41±0.82	0.485
Bloating	-1.03±1.08	-1.09±1.01	0.892
Abdominal Cramps	-0.49±0.92	-0.41±0.75	0.870
Early Satiety	-0.85±1.17	-0.93±1.14	0.620
Acid Eructation /Heartburn	-1.20±1.06	-0.97±0.90	0.261
Sickness	-0.49±1.06	-0.84±0.99	0.096
Loss of Appetite	-0.46±0.88	-0.62±0.81	0.150
Retrosternal Discomfort	-0.46±1.01	-0.50±0.94	0.885
Epigastric or Upper Abdominal Pain	-0.76±0.93	-0.67±1.00	0.707

4 weeks Baseline	GASTIIN CR N=59	Reference Drug N=58	*P-value*
Nausea	-1.19±1.24	-1.19±1.02	0.765
Vomiting	-0.63±1.05	-0.48±0.94	0.432
Bloating	-1.44±0.91	-1.53±0.92	0.542
Abdominal Cramps	-0.63±1.02	-0.55±0.88	0.849
Early Satiety	-1.25±1.18	-1.45±0.92	0.308
Acid Eructation /Heartburn	-1.69±1.00	-1.50±1.01	0.501
Sickness	-0.80±0.92	-0.97±0.99	0.332
Loss of Appetite	-0.47±1.02	-0.81±0.93	0.090
Retrosternal Discomfort	-0.73±0.94	-0.69±0.99	0.832
Epigastric or Upper Abdominal Pain	-0.97±1.08	-0.86±1.05	0.502

Safety Outcome Measures : Adverse reactions

System Organ Class	GASTIIN CR N=68		Reference Drug N=58	
	n (%)	case	n (%)	case
Gastrointestinal disorders	3 (4.41)	3	1 (1.43)	1
Abdominal discomfort	0 (0.00)	0	1 (1.43)	1
Diarrhoea	**1 (1.47)**	**1**	**0 (0.00)**	**0**
Frequent bowel movements	1 (1.47)	1	0 (0.00)	0
Nausea	**1 (1.47)**	**1**	**0 (0.00)**	**0**

The adverse reactions were observed in 1 case of diarrhoea and nausea for a total of 2 cases for GASTIIN CR group, showing no statistically significant differences between the two groups.

* Reference Drug : GASMOTIN Tab. 5 mg t.i.d