HYDRINE Capsules

Hydroxyurea 500mg

$CH_4N_2O_2$ (MW 76.06)

HYDRINE is an antineoplastic agent used in chronic myelocytic leukemia and various other neoplasms.

PHARMACOLOGY

▶ Mechanism of Action

Various studies support the hypothesis that hydroxyurea causes an immediate inhibition of DNA synthesis by acting as a ribonucleotide reductase inhibitor, without interfering with the synthesis of ribonucleic acid or of protein.

▶ Antineoplastic Effect

Hydroxyurea blocks replication in accurately infected lymphocytes and macrophages as well as in blood cells infected in vivo obtained from individuals with neoplasms.

▶ Other Effects

Hydroxyurea is a urease inhibitor.
It also inhibits the production of ammonia and thereby enhance the efficacy of anti-infective agents.
Hydroxyurea may have effects on HIV. It does not attack HIV directly, but instead boosts the potency of some nucleoside analogue reverse transcriptase inhibitors, once they get inside human CD4 cells .

PHARMACOKINETICS

▶ Absorption

Hydroxyurea is readily absorbed from the GI tract. Peak serum concentrations are attained within 2 hours following a single oral dose.

▶ Distribution

Hydroxyurea crosses the blood-brain barrier; peak hydroxyurea CSF concentrations are attained within 3 hours following oral administration.

▶ Elimination

Studies using 14C-labeled hydroxyurea indicate that about one-half an orally administered dose is degraded in the liver and is excreted as respiratory carbon dioxide and in urine as urea. The remaining portion of the drug is excreted intact in urine.

Antineoplastic

HYDRINE Capsules

Hydroxyurea 500mg

Drug Information

Composition

Each capsule contains 500 mg of Hydroxyurea USP

Description

Each capsule, with pink-colored cap and green-colored body, is filled with white to off-white powder.

Indications

Melanoma, resistant chronic myelocytic leukemia, and recurrent, metastatic, or inoperable carcinoma of the ovary.
Hydroxyurea used concomitantly with irradiation therapy is intended for use in the local control of primary squamous cell(epidermoid) carcinomas of the head and neck, excluding the lip.

Dosage & Administration

1) SOLID TUMOR
- Intermittent therapy-80mg/kg administered orally as a single dose every third day.
- Continuous therapy-20 to 30 mg/kg administered orally as a single dose daily.
- Concomitant therapy-80 mg/kg administered orally as a single dose every third day.

2) RESISTANT CHRONIC MYELOCYTIC LEUKEMIA
Until the intermittent therapy regimen has been evaluated, continuous therapy (20-30 mg/kg administered orally as a single dose daily) is recommended.

Storage

Store at room temperature (Below 30℃)

Package

100 Capsules

Manufactured by
KOREA UNITED PHARM. INC.
404-10, Nojang-Ri, Jeondong-Myeon, Yeongi-Kun, Chungnam, Korea

HOME PAGE ON INTERNET ▶ http : // www. kup. co. kr
E-mail ▶ trade @ kup. co. kr

Antineoplastic