

RESEARCH ARTICLE

Strategic Planning of Agroforestry Supply Chains

Thomas Kirschstein^{1,2}¹Fraunhofer IKTS Center for Economics and Management of Materials, Halle, Germany²Hochschule RheinMain – University of Applied Sciences and Arts, Wiesbaden Business School, Wiesbaden, Germany

Correspondence

Thomas Kirschstein

Email: thomas.kirschstein@ikts.fraunhofer.de

Abstract

Agroforestry systems integrate fast-growing woody perennials into agricultural land forming a biological-industrial symbiosis. They simultaneously deliver multiple biomass fractions of differing quality to food, chemical, pulp-and-paper, as well as energy industries through a shared regional supply network. The woody biomass feedstock produced in agroforestry systems can be used to substitute fossil feedstocks particularly in chemical and energy industry. Additionally, biogenic carbon can be stored in long-lived products and agroforestry increases climate resiliency and sustainability of agricultural production. In contrast to the traditional forestry-based supply of woody biomass, agroforestry-based woody biomass shows substantial feedstock potentials as most land is used agriculture in central Europe. Despite these environmental and economic potentials, agroforestry is only applied in marginal scale in central European agriculture. A core challenge in fostering agroforestry is to establish economically viable regional bioeconomy networks which connect partners from agriculture as well as downstream industries. Such a agroforestry-based supply chain needs to be structured and organized in a way that allows all partners to benefit economically in the long run. This paper presents a supply chain design model that helps to support design decisions for agroforestry-based supply chains. Therefore, it models agroforestry establishment and harvest scheduling on farm-level in a large-scaled linear multi-echelon network design model. It relies on biomass growth models which describe the production of multiple biomass types in agroforestry systems. These biomass fractions are pretreated and transported to consumers from chemical, paper-and-pulp, and energy industry depending on quality and demand restrictions. The model is applied in a case study focussing on central Germany where a heterogeneous industrial cluster is located within a fertile and large agricultural production area. The optimization results indicate that agroforestry-based supply chains can be configured in an economically viable way. Agroforestry systems can serve as feedstock suppliers for industrial consumers in the same region at a considerable scale. Profitability of agroforestry-based agricultural practice seem to be competitive with traditional agricultural production and at the same time allow producing woody biomass at a competitive cost, provided transports are not too long.

KEYWORDS

Industrial symbiosis, Agroforestry, Biomass cascading, Bioeconomy network design, Supply chain optimisation, Biobased production

1 | INTRODUCTION

The industrial bio-economy depends on connecting primary biological production systems to multiple industrial sectors through networks that efficiently allocate biomass fractions of varying quality to their

highest-value application (Hildebrandt, O’Keeffe, Bezama, & Thrän, 2019). Agroforestry systems (AFS) that deliberately integrate fast-growing woody perennials into agricultural land represent realisation of this principle. By co-producing heterogeneous biomass fractions like stems, branches, twigs, leaves etc. which are more or less suitable for chemical, paper-and-pulp industry, as well as energy generation, AFS create diverse feedstocks that can serve multiple industrial consumer sectors through a shared regional supply network (Toensmeier, 2017). This biological-industrial symbiosis substitutes fossil feedstocks with renewable bio-based alternatives and stores biogenic carbon in long-lived material products.

Despite their ecological and industrial potentials, temperate AFS occupy only a marginal share of agricultural land in practice. In Germany, approximately 1,200 hectares of silvoarable AFS were established in 2025, accounting for barely 0.02% of total agricultural land (Deutscher Fachverband für Agroforstwirtschaft (DeFAF) e.V., 2026). This gap between supposed potentials and observed practices indicates implementation challenges which often root in economic, organizational, and regulatory obstacles. Establishing AFS at regional scale requires coordinated decisions on stand establishment, harvest scheduling, and biomass allocation across spatially distributed industrial consumers [REF]. E.g. farmers will only invest in AFS when they are convinced to generate revenues from selling wood to industrial customers at the time of harvest which is, however, 5-15 years after establishing an AFS. This long time span imposes a severe investment risk as investment cost are clear and immediate but revenue is uncertain and far away. On the side, industrial consumers particularly in process industries like chemical and paper-and-pulp industry prefer reliable and high-volume biomass supply contracts which imposes a challenge for farmers as AFS growth depends on weather as do harvest decisions. Next to those, a lot of business decisions need to be made. Early harvests promise earlier revenues, but harvested biomass is dominated by small-scaled branches, twigs and leaves with limited usability in downstream industries and lower value [REFS]. At longer rotations, biomass quantity is higher and mostly composed of stems whose diameters surpass quality thresholds such that the share of chemical- and paper-grade material rises markedly (Huber, Amann, Hülsbergen, & Schmid, 2018; Niemczyk & Thomas, 2021). The biomass feedstock quantity and quality in an AFS-based supply chain therefore depend on the rotation strategies and the industrial demand structures of multiple farms and industrial consumers. Current biomass supply chain models, reviewed in subsection 2.3, often do not consider this aspect as biomass availability is usually given at exogenous time series or optimise a single downstream supply chain rather (Zahraee, Shiwakoti, & Stasinopoulos, 2020; De Meyer, Cattrysse, & Van Orshoven, 2016).

This paper closes this gap by formulating an AFS supply chain design (AFS-SCD) problem. The linear program integrates design and operation decisions of biomass production in AFS with multiple downstream industries via logistical processes. This way, interdependencies between biomass production decisions in AFS, logistical handling decisions, and biomass allocation decisions in downstream processes are modeled explicitly. Biomass production decisions are based on growth models which estimate the quantity of biomass types in AFS. Thereby, biomass types are defined based on its suitability for downstream processing. Three different grades are distinguished: chemical/biorefinery grade, pulp and paper grade, and energy grade. Thereby, a quality cascade is assumed. I.e., biorefinery-grade biomass can also be used in pulp-and-paper as well as energy applications, but the reverse doesn't hold. Depending on stand age, quantities of these grades are available in AFS which guide harvest decisions. Once harvests are scheduled, biomass flows to pre-treatment and storage hubs are planned where biomass grades are allocated and transshipments to consumers are prepared depending on demand quantity and the consumers' willingness-to-pay. This way, the proposed AFS-SCD configures a supply chain for a regional AFS-based industrial ecosystem. It helps to study effects of ecological, regulatory, and economic parameters on the profitability of AFS-based supply chains. Thereby, it integrates relevant decisions of different actors from various industries all relevant in AFS-based supply chains including farmers and land-owners, wood logistics, as well as industrial downstream sectors (like chemicals, paper or energy).

The remainder of the paper is structured as follows. Section 2 reviews the relevant literature on AFS, biomass growth modelling, and biomass supply chain design. Section 3 presents the mathematical formulation of the AFS-SCD. Section 4 describes a case study illustrating the application of the AFS-SCD to a real-world problem. Section 5 summarizes the findings of the paper and discusses directions for future research.

2 | LITERATURE REVIEW

Agroforestry biomass systems, stand-level growth models, and biomass supply chain optimisation have each been studied extensively, but they are rarely combined in an integrated framework that links age-dependent stand dynamics to regional product-cascading value chains. This section reviews literature on (i) agroforestry systems and economics, (ii) biomass growth models for AFS considering multiple biomass grades, and (iii) biomass supply chain design problems.

2.1 | Agroforestry systems

Agroforestry deliberately integrates woody perennials with crops and/or livestock on the same management unit, with the aim of enhancing productivity, profitability, and environmental performance (Nair, 1993; Toensmeier, 2017). In temperate climate, three distinct system types can be distinguished based on the spatial arrangement of trees and the rotation length at which woody biomass is harvested.

Short-rotation coppice (SRC) is defined as a monoculture plantation of fast-growing woody species covering the entire management unit, with planting densities of 8,000–20,000 trees per hectare. After each harvest, the root system regenerates without replanting. Rotation cycles are typically 3–7 years and total stand lifetime is 20–25 years (Trnka et al., 2008; Faasch & Patenaude, 2012). SRCs are not AFS in the narrow sense, but it is helpful to benchmark biomass production.

Short-rotation agroforestry systems (SRAFS) integrate SRC-type tree strips (typically 1–4 rows of the same fast-growing species like poplar or willow) into an otherwise arable field in an alley-cropping arrangement. Crop production continues in the inter-row strips (Huber et al., 2018). Thus, tree strips typically cover 10–30% of the total parcel area (Graves et al., 2007). Rotation lengths and total stand lifetime in SRAFS are comparable to SRCs. In SRAFS most trees are effectively edge trees, exposed to higher light and water availability than interior SRC trees. This boundary effect raises individual-tree biomass accumulation but also intensifies competition with adjacent crop rows (Huber et al., 2018).

Long-rotation agroforestry systems (LRAFS) use the same alley-cropping spatial structure as SRAFS but with rotation cycles of 12–15 years or longer (Graves et al., 2007; Nair, 1993). At this rotation length, individual trees develop substantial stem diameters and the proportion of high-value timber increases markedly, enabling premium material uses (veneer, sawlogs, engineered wood products). Therefore, also timber-quality tree species (such as walnut, cherry, oak, or high-grade poplar clones) are planted [REFS].

SRAFS and LRAFS are not clearly differentiated in practice. First, a clear cut-off age is hard to define and, more importantly, the intended use of AFS biomass is not always clear when establishing the AFS and can also change over time. Nonetheless, when establishing AFS, the structure of AFS and its intended use after harvest affect economic viability substantially. Thus, in the present paper SRAFS and LRAFS are not considered as separate AFS systems, but the interplay of intended use, establishment, and harvesting decisions is analysed.

The decision of a landowner to establish an AFS involves weighing substantial up-front costs and long-term opportunity costs against a diversified revenue stream. On the cost side, establishment requires site

preparation, planting material, and planting labour, typically amounting to 1,500–3,500 /ha for SRC or SRAFS (Faasch & Patenaude, 2012; Testa, Di Trapani, Foderà, Sgroi, & Tudisca, 2014). Recurrent harvest and refitting costs (machine hire, chipping, transport to roadside) add approximately 200–400 /ha per rotation depending on stand density and equipment availability (Testa et al., 2014; Spinelli, Nati, & Magagnotti, 2009). During the rotation, maintenance costs of AFS arise on a yearly basis. Next to expenses for establishment, maintenance and harvesting, opportunity costs arise as no revenues from crops are generated on tree strips. In regions with fertile soils and high cereal prices, yearly crop revenues may sum up to 400–600 /ha (Faasch & Patenaude, 2012; Graves et al., 2007). Thus, opportunity costs are often the dominant cost driver in SRAFS profitability analyses.

On the benefit side, the primary revenue of AFS is the sale of harvested woody biomass, whose magnitude and frequency depends on rotation length, species, and site productivity. For poplar-based SRAFS in Central Europe, typical yields range from 10 to 25 t of dry matter biomass per ha for typical rotation lengths (Huber et al., 2018; Niemczyk & Thomas, 2021). Additionally, positive crop-yield effects in the inter-row strips of SRAFS and LRAFS may result. Tree rows act as windbreaks, reducing soil erosion and evapotranspiration. They provide partial shading that moderates heat stress during drought years and their root systems may improve water retention in top-level soil (Graves et al., 2007; Grünewald, Böcker, Schneider, Bens, & Hüttl, 2007). Empirical measurements in Central European alley-cropping trials indicate that these microclimate effects can increase cereal yields in adjacent strips by 5–15% relative to open-field controls, partially or fully compensating the loss of arable area occupied by the tree strips (Graves et al., 2007). Beyond direct yield effects, AFS deliver broader ecosystem services (like soil organic matter accumulation, carbon sequestration, biodiversity enhancement) that may lead to agri-environment subsidies and carbon credits which have to be considered as additional financial benefits (Toensmeier, 2017; Faasch & Patenaude, 2012).

Nonetheless, economic attractiveness of AFS depends primarily on revenues of AFS biomass. Thereby, revenues are different for the types of biomass and in which value chain it is used. Often AFS biomasses are used energetically, i.e. for heat and power generation, or anaerobic digestion, which typically creates low prices per tonne of biomass [REF]. Alternatively, woody parts of the biomass (i.e., the stem and large branches) can also be used as pulp and paper grade material as well as feedstock for chemical industry or as timber achieving higher prices (Lamers, Junginger, Hamelinck, & Faaij, 2011) [EXTEND REFS + Explain Forestry Product Classification]. Thus, fostering AFS establishment by maximizing economic returns from AFS biomass requires a strategy for allocating biomass fractions to the highest-value applications,

while ensuring that the rotation length is aligned with the growth dynamics of the stand and the market demand for different biomass types (Lamers et al., 2011).

Thus, for making educated decisions on AFS establishment and harvesting, the expected biomass quantities at harvest for stem, branches, and residues need to be estimated. Thereby, the shares of the biomass types change with stand age. E.g.~the share of stem weight rises from roughly 55% at age 4 to over 75% at age 10 (Huber et al., 2018; Testa et al., 2014). Therefore, in subsection 2.2 growth modelling approaches are reviewed that capture the age-dependent growth dynamics of different biomass types in AFS.

2.2 | Biomass growth models

Reliable estimation of stand-level biomass quantities and their age-dependent composition is a necessary prerequisite for strategic analysis of AFS-based supply chains. Individual-tree biomass is commonly approximated by species-specific allometric power laws relating stem base diameter to tree biomass. Such relationships have been calibrated e.g. for Central European poplar clones under SRC and SRAFS conditions (Huber, Hülsbergen, & Schmid, 2016; Huber et al., 2018). For supply chain planning, however, the relevant quantity is stand-level biomass per hectare over a multi-decade rotation horizon, which requires a temporal growth model. In the following, temporal biomass growth models for different biomass fractions are derived. Therefore, a set of variables is used which are summarized in Table 1 in the appendix.

Stand-level biomass per hectare is denoted by $M(t)$. Thereby, stand-level biomass growth can be described by different growth functions like the logistic, Gompertz, Bertalanffy, and Richards growth model [REFS]. All these functions show a right-skewed sigmoidal pattern. Thereby, the shape of these functions can be controlled by various parameters. The growth functions need more or less parameters whereby more parameters usually imply a more flexible growth model [REF]. On the other side, more parameters require more calibration efforts which have to be counterweighted with modelling accuracy (Tjørve & Tjørve, 2017). Empirical studies like (Barrio-Anta, Sixto-Blanco, Cañellas-Rey de Viñas, & Castedo-Dorado, 2008) and (Paula et al., 2013) indicate that most growth models can be calibrated to empirical observations quite accurately. Therefore, in the following, a comparatively simple Gompertz model is chosen for modelling biomass quantity, which is defined as

$$M(t) = A(N) \cdot \exp(-\exp(-k \cdot (t - t_0))), \quad (1)$$

where t_0 is the age of maximum instantaneous biomass increment and k determines how quickly the stand approaches its site-specific asymptote $A(N)$. For Central European poplar under medium-rotation AFS management (10–20 yr), analysing empirical data from (Huber et al., 2018), (Niemczyk & Thomas, 2021), and (Jha, 2018) yields parameter intervals of $k \in [0.17, 0.2]$ and $t_0 \in [9, 10]$ years, depending on local specifications of the AFS under study. Parameter t_0 indicates an upper bound on the optimal rotation age as growth slows down or diminishes entirely close to this age.

The asymptote $A(N)$ depends largely on planting density and can be modelled by a competition-density power law (Kira, Shinozaki, & Hozumi, 1961):

$$A(N) = C_{\text{site}} \cdot N^{1-\beta}, \quad (2)$$

where C_{site} encodes site quality, N indicates tree density (e.g. trees per hectare), and β is the competition exponent. Equivalently, the per-tree asymptote $A_{\text{tree}}(N) = A(N)/N = C_{\text{site}} \cdot N^{-\beta}$ decreases with density as individual trees have access to less growing space. Parameters C_{site} and β are highly site- and species-specific. For Poplar clones under European conditions, $\beta \in [0.3, 0.5]$ and $C_{\text{site}} \in [4, 6.5]$ (Niemczyk & Thomas, 2021). Figure 2 in the appendix shows asymptote functions according to (2) for advantageous and disadvantageous site conditions.

When assessing potential downstream application of AFS biomass, biomass fractions are to be defined meeting the requirements of downstream consumers. E.g. for timber and biochemical applications, stem diameters of AFS trees need to exceed a minimum threshold in order to qualify as a suitable feedstock. In the following three biomass fractions are distinguished: stems, branch wood, and residue. The stem fraction consists of stems with a diameter of at least 15 cm, while branch wood requires a diameter of at least 7 cm. The residue fraction comprises the smaller parts of the tree. To subdivide the total AFS biomass modelled by (1) into the three fractions depending on tree age, a logistic growth model is used:

$$q_p(t) = \frac{f_p}{1 + e^{-r_p(t-t_{50,p})}}, \quad (3)$$

where $t_{50,p}$ is the age at which 50% of compartment p biomass surpasses the diameter threshold and f_p denotes the asymptotic share of component p for mature trees. Using empirical data from (Civitarese, Acampora, Sperandio, Assirelli, & Picchio, 2019), (Jha, 2018) and (Niemczyk & Thomas, 2021), $r_{\text{stem}} = 0.467, \text{yr}^{-1}$, $t_{50,\text{stem}} = 8.19$ years, and $f_{\text{stem}} = 0.75$ is estimated for the stem fraction (threshold

$d > 15$ cm) and $r_{bran} = 0.698 \text{ yr}^{-1}$, $t_{50,bran} = 7.55 \text{ yr}$, and $f_{bran} = 0.15$ for the branch fraction (diameter at least 7 cm at insertion). Figure 1 illustrates the development of the fraction shares over stand age.

Combining equations (1)–(3), the dry biomass yield of fraction p per hectare at harvest age t is:

$$\eta_p(t) = q_p(t) \cdot M(t), \quad (4)$$

with all non-merchantable fractions aggregated into the residue fraction. Fresh-weight yields at felling are obtained from dry matter yields via $\eta_p^{fw}(t) = \eta_p(t)/(1 - \omega)$, where the moisture content at felling $\omega = 0.55$ is taken from (Civitarese et al., 2019). Figure 1 shows the fractions' stacked biomass quantities $\eta_p(t)$ depending on stand age compared with empirical observations from (Jha, 2018) and (Huber et al., 2018).




FIGURE 1 Stand-level aboveground biomass components for poplar under AFS management with $N = 250$ trees, ha^{-1} . Grey band: typical rotation times (5-15 years).

2.3 | Biomass supply chain design

The design and planning of biomass supply chains has attracted substantial research attention over the past two decades, driven by the need to ensure cost-efficient and sustainable feedstock provision for bioenergy and biorefinery applications. Several systematic reviews provide comprehensive overviews of this

literature. (Cambero & Sowlati, 2014) survey economic, social, and environmental assessment and optimisation studies for forest biomass supply chains, identifying MILP as the dominant modelling approach and noting a growing trend towards multi-objective formulations integrating life cycle assessment.

(Yue, You, & Snyder, 2014) review biomass-to-bioenergy and biofuel supply chain models with a focus on multi-scale modelling, uncertainty treatment, and sustainability metrics, and conclude that the coupling of upstream feedstock dynamics with downstream network design remains an underexplored area. (Malladi & Sowlati, 2018) survey logistics-specific features of biomass supply chain models—seasonal availability, moisture content degradation, preprocessing steps—and highlight that most models treat biomass as a homogeneous commodity, neglecting quality differentiation. (Zahraee et al., 2020) provide a recent meta-review of 140+ studies, confirming that cost minimisation is still the dominant objective, and that multi-product cascading formulations are rare.

The structure of biomass supply chain models typically encompasses three decision layers: (i) strategic network design (facility location, capacity sizing, technology selection), (ii) tactical planning (harvest scheduling, seasonal flows, inventory), and (iii) operational routing. The following paragraphs discuss key individual contributions and their specific methodological advances.

(Ekiolu, Acharya, Leightley, & Arora, 2009) develop one of the earliest large-scale MILP models for biomass-to-biorefinery supply chain design, integrating facility location, capacity sizing, and multimodal transport for lignocellulosic feedstocks. Their model minimises total system cost and introduces binary facility-opening variables that have since become standard in the field. A key limitation, however, is that biomass supply is treated as an exogenous input, so feedstock availability is not endogenised as a function of land-use or growth decisions. (Mansoornejad, Pistikopoulos, & Stuart, 2013) extend the strategic design perspective to forest biorefineries by integrating product portfolio design with supply chain configuration in a single MILP. Their model explicitly accounts for product conversion efficiencies along different processing pathways, enabling joint optimisation of process technology selection and logistics.

(De Meyer, Cattrysse, Rasinmäki, & Van Orshoven, 2014) introduce the OPTIMASS framework, a generic MILP for biomass-based supply chain optimisation that explicitly models product quality cascades: higher-quality biomass classes can substitute for lower-quality demands downstream, whereas the reverse is infeasible. This cascade hierarchy, formalised through ordered product sets, is particularly relevant for woody biomass where stem, branch, and residue fractions command different market prices. The companion t-OPTIMASS extension (De Meyer et al., 2016) incorporates temporal dynamics by explicitly representing biomass growth and regeneration over multiple periods, enabling simultaneous optimisation

of harvest timing and network flows. However, in both OPTIMASS formulations the growth dynamics are represented as exogenous time-series rather than as age-structured stand models, so rotation-length decisions and age-dependent biomass quality are not endogenised. (Arabi, Yaghoubi, & Tajik, 2019) demonstrate multi-period MILP extensions for bio-based supply chains with production dynamics, but focus on microalgae rather than woody biomass and do not address product quality cascades.

(Sharma, Ingalls, Jones, & Khanchi, 2013) identify supply-side uncertainty (yield variability, seasonal availability, moisture content) as the most critical source of risk in biomass supply chains. In response, a strand of literature has developed stochastic and robust optimisation extensions. (Zahraee et al., 2020) document that two-stage stochastic MILP and robust optimisation approaches now account for roughly 30% of the modelling literature, with biomass yield uncertainty and price volatility as the dominant stochastic parameters.

Biomass supply chain research focusing specifically on agroforestry or SRC contexts is comparatively sparse. (Espinoza, Bautista, Narvaez, Alfaro, & Camargo, 2017) note that most supply chain models assume continuous feedstock availability and do not represent the periodic harvest cycles characteristic of coppice systems. (Lamers et al., 2011) analyse feedstock design principles for lignocellulosic supply chains and argue that the allocation of different biomass fractions to valorisation pathways—product cascading—is a key lever for economic viability, yet is rarely optimised jointly with network design decisions. The model developed in Section 3 of this paper directly addresses this gap by endogenising stand age dynamics through binary age-lag variables and coupling component-specific yield coefficients with a product quality cascade.

Table 1 summarises the key modelling features of the reviewed contributions.

TABLE 1 Key modelling features of selected biomass supply chain design studies.

Study	Objective	Multi-product	Quality cascade	Age dynamics	Uncertainty	Level
Eksioglu et al. (2009)	Cost min.	—	—	—	—	Strategic
Sharma & Taaffe (2013)	Cost min.	Partial	—	—	—	Strategic/Tactical
Mansoornejad et al. (2013)	Profit max.	✓	—	—	—	Strategic
Cambero & Sowlati (2014)	Cost/GHG/jobs	—	—	—	Stochastic	Strategic
Yue et al. (2014)	Cost/GHG	✓	—	—	—	Multi-scale
De Meyer et al. (2015)	Cost min.	✓	✓	Exogenous	—	Strategic/Tactical
De Meyer et al. (2016)	Cost min.	✓	✓	Exogenous TS	—	Multi-period
Lamers et al. (2015)	Cost/quality	✓	Partial	—	Scenario	Strategic
Espinoza et al. (2017)	Sustainability	✓	—	—	Scenario	Strategic
Malladi & Sowlati (2018)	Cost min.	Partial	—	—	—	Tactical/Oper.
Arabi et al. (2019)	Profit max.	✓	—	Exogenous	—	Multi-period
Zahraee et al. (2020)	Multi-obj.	✓	—	—	Stochastic	Strategic/Tactical
This paper	NPV max.	✓	✓	Endogenous	Scenario	Strategic/Multi-period

3 | MILP MODEL FOR AGROFORESTRY SUPPLY CHAIN DESIGN

Building on the biomass growth models introduced in subsection 2.2 and the supply chain design literature reviewed in subsection 2.3, this section describes the AFS supply chain design problem (AFS-SCD), a mixed-integer linear programming (MILP) model for the integrated design and operation of an agroforestry biomass supply chain with multiple AFS biomass types (i.e., stem, branch, and residue fractions) used as feedstocks for potential consumers in chemical, pulp and paper as well as energy industry.

3.1 | Sets, Indices, and Time Structure

The model considers a set of agroforestry sites $\mathcal{I}$, storage/pre-treatment facilities $\mathcal{J}$, consumer sites $\mathcal{K}$, and biomass types $\mathcal{P} = \{stem, branches, residues\}$, whose growth models are outlined above. Time is discretised into annual periods $t \in \mathcal{T} = \{1, \dots, T_{\max}\}$. To reduce complexity, reflect legal definitions of SRAFS and LRAFS as well as discard implausible options (e.g. extremely long rotation cycles), $A^{\min}$ and $A^{\max}$ define minimum and maximum stand ages. Thus, harvest are possible in periods $\mathcal{T}^{\text{harv}} = \{A^{\min} + 1, \dots, T_{\max}\}$. To consistently model establishment and consecutive harvest decisions, the following arc sets are defined: $\mathcal{S}^{\text{est}}$ defines potential establishments of AFS. If an AFS is established in period $t \in \mathcal{T}$, arc $(0, t)$ is selected. Harvest set $\mathcal{S}^{\text{harv}}$ summarizes all potential consecutive harvest periods within the age interval $[A^{\min}, A^{\max}]$. I.e., if an AFS is established or harvested in period s and next harvest is in period t , arc (s, t) is selected. Finally, if there is no subsequent harvest after period t , the AFS is terminated by selecting arc $(t, T^{\max} + 1)$. Termination arcs are summarized in set $\mathcal{S}^{\text{term}}$. All arc sets are formally defined as follows:

$$\begin{aligned}\mathcal{S}^{\text{est}} &= \{(0, t) \mid t \in \{1, \dots, T^{\max} - A^{\min}\}\}, \\ \mathcal{S}^{\text{harv}} &= \{(s, t) \mid s, t \in \mathcal{T} \wedge t - s \in \{A^{\min}, \dots, A^{\max}\}\}, \\ \mathcal{S}^{\text{term}} &= \{(s, T^{\max} + 1) \mid s \in \{A^{\min} + 1, \dots, T_{\max}\}\}, \\ \mathcal{S} &= \mathcal{S}^{\text{est}} \cup \mathcal{S}^{\text{harv}} \cup \mathcal{S}^{\text{term}}.\end{aligned}$$

Figure 2a illustrates this network structure of all arc sets for $T = 8$, $A^{\min} = 3$, $A^{\max} = 5$.




FIGURE 2 Network structure of set $\mathcal{S}$ for $T = 8$, $A_{\min} = 3$, $A_{\max} = 5$. Dashed blue arcs: establishment arcs $\mathcal{S}^{est}$, solid green arcs: harvest arcs $\mathcal{S}^{harv}$, solid red arcs: termination arcs $\mathcal{S}^{term}$.

3.2 | Decision Variables

Temporal flow Variables z_{ist} indicate on which quantity of the area of site i an AFS is operated between $(s, t) \in \mathcal{S}$. Depending on the arc type, this implies establishment, harvest, or termination. Selected arcs build a network flow over the complete planning horizon from $t = 0$ to $T^{max} + 1$. Figure 2b illustrates a simple network flow with establishment of an AFS in period 1 and harvests in periods 5 and 8.

Spatial flow variables X_{ijpt} and $X_{jkpp't}$ represent transport of products from AFS sites to storage/pre-treatment facilities and from there to consumers, respectively, while S_{jpt} captures end-of-period inventory at facility j . Table 2 in the Appendix summarises all sets, decision variables, and parameters used in the MILP formulation, while η_{pa} refers to (4).

3.3 | Objective Function

As discussed in Section subsection 2.1, the economic attractiveness of AFS supply chains is modelled from a global perspective taking into account all processes, activities, and participants involved in finally serving customer markets with AFS-based feedstocks. Therefore, revenues for the different biomass fractions and are weighed against costs occurring at all value-adding stages of the supply chains. Revenues created by selling AFS-based feedstocks to different industrial consumers are counterweighted by establishment, opportunity, harvest, transport, and storage costs of landowners, agricultural firms as well as biomass

logistics companies. The model therefore maximizes total profit of all participants in the AFS supply chain over the planning horizon as follows:

$$\begin{aligned}
 \max Z = & \sum_{k \in \mathcal{K}} \sum_{p \in \mathcal{P}} R_{kp} \cdot \sum_{j \in \mathcal{J}} \sum_{(p', p) \in \mathcal{Q}} \sum_{t \in \mathcal{T}} X_{jkp't} \\
 & - \sum_{i \in \mathcal{I}} c_i^{est} \cdot \sum_{(0, t) \in \mathcal{S}^{est}} z_{i0t} \\
 & - \sum_{i \in \mathcal{I}} (c_i^{main} + c_i^{opp}) \cdot (t - s) \cdot \sum_{(s, t) \in \mathcal{S}^{harv}} z_{ist} \\
 & - \sum_{i \in \mathcal{I}} c_i^{harv} \cdot \sum_{(s, t) \in \mathcal{S}^{harv}} z_{ist} \\
 & - \sum_{i \in \mathcal{I}} \sum_{j \in \mathcal{J}} \sum_{p \in \mathcal{P}} \sum_{t \in \mathcal{T}} c_p^{tr-raw} \cdot d_{ij} \cdot X_{ijpt} \\
 & - \sum_{j \in \mathcal{J}} \sum_{k \in \mathcal{K}} \sum_{(p, p') \in \mathcal{Q}} \sum_{t \in \mathcal{T}} c_{p'}^{tr-pre} \cdot d_{jk} \cdot X_{jkpp't} \\
 & - \sum_{j \in \mathcal{J}} \sum_{p \in \mathcal{P}} \sum_{t \in \mathcal{T}} c_j^{stor} \cdot S_{jpt}
 \end{aligned} \tag{5}$$

Here, R_{kp} are product- and consumer-specific revenues for each ton of biomass fraction ($\{ \}$ /t). The opportunity cost term penalises the land use when AFS are established. Thereby, c_i^{opp} can be negative or positive depending on the site-specific conditions. It comprises effect of lost field crop revenues due to AFS as well as higher yields of remaining field crop areas e.g. due to wind and water erosion protection by AFS. While the assessment of lost field crop revenues is usually well predictable, the economic evaluation of positive ecological effects of AFS (e.g. on neighboring crop yields) is more difficult. Thus, c_i^{opp} can also contain regulatory subsidies to support establishing and maintaining AFS. Transport cost terms mirror the two-stage flow structure of the AFS supply chain with differentiated rates for raw and pre-treated biomass depending on the biomass type. Cost rates differ because as e.g. the density and water content of the shipped biomass types differ or because specialized equipment (e.g. for log transports) is used [REFERENCES]. Moreover, site-specific establishment cost for planting the AFS trees is considered, next to yearly maintenance and harvesting costs. All these costs scale with the size of a site ($AREA_i$) measured in hectare. Maintenance cost comprise e.g. costs for trimming, replanting, and pruning unnecessary shoots. Storage costs are considered if biomass types are stored for more than a year. Storage cost rate c_j^{stor} measures the costs for storing a ton of biomass for one year and comprises direct storage cost (e.g. for storage site maintenance) as well as opportunity cost due to potential material losses and capital commitment.

Biomass types harvested and consumed in the same year incur no storage cost, as usually AFS are harvested at the beginning of a year. Harvested biomass has to dry during spring and summer such that the biomass can be used in autumn or winter the same year. As under-year drying is necessary for some industrial applications and beneficial for transport economics, drying storage in the year of harvest is not associated with additional cost in this study.

3.4 | Constraints

Constraints (6) limits the area dedicated to AFS operations to the maximum size of site i . Constraints (7) enforce flow conservation in each period. If AFS systems of size z_{ist} are established or harvested in period $t \in \mathcal{T}$, there needs to be a succeeding activity (harvest or termination) of this area. This way, (7) guarantee valid (possibly empty) AFS activity over the complete time horizon for each site. Constraints (8) links the harvest-cycle variables z_{ist} to physical yield quantities $\eta_{p(t-s)}^{fw}$ for age $t - s$, which are pre-computed using (1)–(4) scaled by moisture content ω . Thus, we assume constant yields per hectare for each site i . Note that this simplification can be dropped and site-specific yields $\eta_{ip(t-s)}^{fw}$ could be introduced without increasing numerical complexity of the AFS-SCD. Constraints (9) link harvest quantities to transport flows and ensure inventory balance at each facility. Thereby, harvest quantities can be smaller than available AFS biomass allowing partial harvesting. Constraints (10)–(11) enforce storage and processing capacity limits at each facility j . Constraints (12) implement product usage at multiple consumption sites k . Thereby, $(p', p) \in \mathcal{Q}$ defines a product mapping allowing high-quality product p' (e.g. stems) satisfying demand of a lower-quality products p up to maximum demand $D_{kpt}^{\max}$.

$$\sum_{(0,t) \in \mathcal{S}^{est}} z_{i0t} \leq AREA_i \quad \forall i \in \mathcal{I} \quad (6)$$

$$\sum_{(s,t) \in \mathcal{S}} z_{ist} = \sum_{(t,u) \in \mathcal{S}} z_{itu} \quad \forall i \in \mathcal{I}, t \in \mathcal{T} \quad (7)$$

$$\sum_{j \in \mathcal{J}} X_{ijpt} \leq \sum_{(s,t) \in \mathcal{S}^{harv}} \eta_{p(t-s)}^{fw} \cdot z_{ist} \quad \forall i \in \mathcal{I}, p \in \mathcal{P}, t \in \mathcal{T}^{harv} \quad (8)$$

$$S_{jpt} = S_{jp,t-1} + \sum_{i \in \mathcal{I}} X_{ijpt} - \sum_{k \in \mathcal{K}, (p,p') \in \mathcal{Q}} X_{jkpp't} \quad \forall j \in \mathcal{J}, p \in \mathcal{P}, t \in \mathcal{T}^{harv} \quad (9)$$

$$\sum_{p \in \mathcal{P}} S_{jpt} \leq CAP_j^{stor} \quad \forall j \in \mathcal{J}, t \in \mathcal{T} \quad (10)$$

$$\sum_{i \in \mathcal{I}, p \in \mathcal{P}} X_{ijpt} \leq CAP_j^{proc} \quad \forall j \in \mathcal{J}, t \in \mathcal{T}^{harv} \quad (11)$$

$$\sum_{j \in \mathcal{J}, (p',p) \in \mathcal{Q}} X_{jkp'pt} \leq D_{kpt}^{\max} \quad \forall k \in \mathcal{K}, p \in \mathcal{P}, t \in \mathcal{T}^{harv} \quad (12)$$

$$z_{ist}, X_{ijpt}, X_{jkp'pt}, S_{jpt} \geq 0 \quad \forall i, j, k, p, p', t \quad (13)$$

314 In summary, AFS-SCD constitutes a large-scaled LP which allows to decide on which fraction of a
 315 site, AFS are established and how the AFS operations are scheduled in a multi-year planning horizon.
 316 The AFS-SCD takes a supply chain perspective and maximizes total profit of all actors involved in these
 317 processes. Thus, it does not explain how the profits are split e.g. between farmers, AFS service providers
 318 and logistics service providers. The continuous modelling of AFS areas in z_{ist} leads to a comparatively
 319 easy to solve problem. On the other side, it is unclear which part of a site's total area is chosen for AFS
 320 operations if $z_{ist} < AREA_i$, which holds for establishment as well as harvesting and termination.

321 4 | CASE STUDY

322 The AFS-SCD is intended to support the design of regional agroforestry biomass supply chains fostering re-
 323 gional bioeconomy clusters thereby connecting multiple industrial sectors. The proposed AFS-SCD model
 324 is instantiated on a real-world biomass supply chain case study located in the tri-state region of Saxony-
 325 Anhalt, northern Thuringia, and northern Saxony, Germany. This region is characterized by a high density
 326 of agricultural land usage as well as cluster of paper and wood as well as chemical industry. The re-
 327 gional chemical industry relies on fossil resources so far, but is under substantial transformative pressure

which led to investments in sustainable production processes requiring sustainable regional bio-based feedstocks. Figure 3 in the Appendix shows agricultural production sites along with the set of potential wood-processing hubs, and the regional users of AFS biomasses from chemical, paper and wood processing industry as well as the energy sector.

The supply chain comprises 2239 agricultural production parcels which are grouped into $|Z| = 145$ potential AFS production sites. Clustering is based on geographical distances between parcel centroid using a k-means approach in order to reduce problem size. For distance calculations, for each site a centroid is calculated. Pre-treatment and intermediate storage facilities $\mathcal{J}$ are located on existing wood processing facilities as well as potential new log or wood chip yards. All harvested biomass quantities at AFS sites are transported to logistics hubs first. At the hubs sorting and transshipment processes take place before biomass types are forwarded to the consumer facilities. Storing and transshipment capacities at the hubs are assumed to be constant in time and universally set to 4.0909×10^4 kt per year. Finally, $|K| = 28$ consumer sites are included. Each consumer site belongs to an existing facility from chemical, paper/wood processing industry, or the energy sector. Distances between the facilities d_{ij} and d_{jk} are calculated as road distances between the facilities' coordinates. For each consumer facility, an yearly upper demand capacity for each of the three distinguished biomass product types are given along with consumer-specific prices. The following biomass types are distinguished:

- **P1: stem wood – chemical/paper grade:** long-fibre wood chips from stems suitable for conversion in a local biorefinery and paper factory. In total, a yearly demand of 620 kt and an average prices of 45 is assumed.
- **P2: branch wood – sawmill/pulp grade:** medium/short-fibre wood chips from larger branches suitable for mechanical processing in sawmills and lower-grade pulping. Demand is distributed across 6 smaller facilities and accumulates to 410 kt/year. Gate prices range from 28-38 /t.
- **P3: residue – energy grade:** residual wood fractions (chips, bark, off-cuts) used for heat and electricity generation or lower-grade pellet production. Demand is accumulates among 24 consumers to 132 kt/year. Gate prices range from 18-24 /t.

Table 3 in the appendix summarizes the consumer-specific demand data. It is assumed that the biomass product types show a quality cascade which means that the demand of pulp grade (P2) can also be satisfied by chemical/paper grade biomass (P1). Likewise, energy grade demand (P3) can be satisfied by all biomass types. Note that constant demands over the planning horizon are assumed, although an increasing biomass

demand can be expected due to the green transition in those industries. From this perspective, a rather conservative market demand scenario is assumed.

Growth model introduced in subsection 2.2 is used to model biomass quantities depending on age and the area dedicated to AFS. The growth model is parameterized for poplar clones at central European conditions as introduced above with a planting density of $N = 400$ trees per hectare. If AFS are established, they have to be harvested within $A^{min} = 3$ to $A^{max} = 20$ years allowing a wide range of potential harvesting strategies between short-rotation and long-rotation AFS. As the total planning horizon is $|T| = 40$ years multiple harvesting cycles can be established. The sites do not differ with respect to growth conditions as the climatic and soil quality variation within the comparatively small region is rather limited. However, the sites differ with respect to size (given in hectare) and opportunity cost rates (given in per hectare and year). Site-specific opportunity cost rates c_i^{opp} is normally distributed with mean 521 and standard deviation 391. For most sites, substantial opportunity costs are assumed reflecting the loss in field-crop revenue, risk premiums of landowners as well as higher costs for field crop maintenance. The 145 sites with negative opportunity cost rates reflect sites where AFS lead to an increase in field crop productivity on the remaining field area. The variation in opportunity cost rates to study the effect of opportunity cost on site selection for AFS and gives valuable insights particularly for communication with farmers and politics. Subsidies for establishing AFS are one of the core instruments to foster AFS. The height and effectiveness of those subsidies, however, is controversially disputed. Remaining cost parameters are universally set for all sites and comprise maintenance cost rates c_i^{main} (per year and hectare), establishment cost c_i^{est} (per hectare) and harvesting cost c_i^{harv} (per hectare). Likewise, transport cost rates for harvested biomasses c_p^{tr-raw} and pre-treated/sorted biomasses $c_{p'}^{tr-raw}$ are set universally. All scalar parameters are shown in Table 4 in the Appendix.

4.1 | Optimization results

The AFS-SCD model is solved for the instance data described above with Gurobi. The model selects a subset of sites for AFS establishment and identifies the optimal harvest schedules, intermediate facility utilisation, and product allocation across the planning horizon.

Figure 3 shows the spatial configuration of the profit-optimal supply chain.

In total, on 38 sites AFS are established which corresponds to about 26 % of all sites and 24 % of the total farm area. Over the planning horizon, 2.6368×10^4 kt of AFS biomasses are produced on these sites




FIGURE 3 Site selection and biomass production in optimal supply chain configuration. Green polygons: active sites with biomass harvest; grey polygons: non-selected sites. Pie charts at active sites show the product mix of harvested biomass, scaled by total harvested volume.

which suffices to satisfy 57 % of total demand. Figure 4 shows the harvested biomasses over time along with total demand of all consumer sites.

During the first 10 years no AFS biomasses are harvested as trees of established AFS in first years grow up to a certain size before first harvest. Thus, about 25% of consumers' total biomass demand cannot be satisfied due to this grow-up period. Afterwards, a cyclical pattern appears with a harvesting peak around year 20 and a breakdown in year 28 before plateau is reached from year 35-40. Thereby, the harvesting




FIGURE 4 Total biomass delivered to consumers by product type over the planning horizon. Stacked areas show period-level volumes of stem (P1), branches (P2), and residues (P3) flowing from storage hubs to all consumers. Dashed line: total harvested raw biomass shipped from sites to hubs.

quantities of stems and branches show a similar pattern while quantities of harvested energy grade biomass are rather stable and close to the demand limit over the planning horizon. Thus, rather long-rotation AFS are established as Figure 5 confirms showing the histogram of harvesting ages across all sites.




FIGURE 5 Distribution of harvesting cycle lengths at active agroforestry sites. Each bar represents the number of individual harvest events (over all periods and active sites) whose cycle length (stand age at harvest, i.e. $t - s$) falls in that year bin.

The histogram illustrates that majority of harvest events occur at stand ages between 14–18 years, indicating that long-rotation AFS seem to be more profitable in this setting. Short-rotation cycles (<10 years) occur only in marginal frequencies. This optimal rotation length might be caused either due to the limited demand of energy-grade biomass or the price gaps between biomass types or a combination of effects. The quantity of energy-grade biomass created in long-rotation AFS suffices to cover total energy-grade biomass demand almost completely. Rotation length is focused on production of chemical-grade biomass where highest prices are assumed. Focusing on chemical industry supply requires a production of trees with a comparatively large stems. This strategy maximizes revenue, but in the same time increases opportunity costs as wood harvests appear in larger time intervals. Figure 6 presents the aggregate cost–profit waterfall over all periods and actors.




FIGURE 6 Total cost–profit breakdown across all actors and years. The left panel (waterfall) decomposes the objective value into revenue and individual cost categories; the right panel shows the period-level trajectory of revenues and stacked costs.

It appears that revenue is dominated by selling chemical-grade biomass and dominating cost component are opportunity cost although active sites are those sites with a comparatively small opportunity cost rate. Next to opportunity cost, establishment and transport costs are the most relevant cost drivers. Analyzing total cashflow over time shows that break-even is reached after about 20 years, i.e. after the first large-scale harvest occurred. Afterwards, cashflow is continuously increasing indicating that establishing AFS is a strategic change in agricultural practice and pays only if this strategy is implemented and maintained

over a comparatively long time spans. This underpins the importance of strategic buying contracts with industrial partners in order to reduce revenue risks of farmers when producing AFS biomass.

Although profitability of the total AFS supply chain is given for the case study data, profitability of AFS operations on site-level are more heterogeneous. Figure 7 shows revenues and costs split by component and normalized to AFS size and active years of AFS.




FIGURE 7 Revenues, costs and net profit for sites with AFS operations split by type and expressed in per hectare per year of AFS operation.

It appears that net profitability of AFS ranges between 282 and 6 per hectare and active year. Note that this value constitutes additional profit after subtracting operational cost as well as imputed opportunity cost of lost profit from field crop. I.e., for calculating the realized net margin, opportunity can be added such that realized yearly net profit ranges between 473 and 210 per hectare. Still, relative net profit is comparatively low compared to ordinary agricultural practices with profitable field crops (like winter wheat). The revenue splits of active sites indicates that at some sites medium-grade and low-grade biomass types contribute substantially or even solely to total revenue (see e.g. site 12, 135, 145, and 112). This indicates that site-specific AFS strategies might differ from the overall supply chain strategy. In other words, there seem to be strategic niches w.r.t. the consumer segments to be supplied. This might also be reflected in the rotation length and, although not discussed here, the selection of trees and clones for the AFS. Moreover, profitability of AFS sites seems to depend on various factors like local opportunity costs as well as closeness to consumers. Figure 8 shows net margins and opportunity cost rates of sites with active AFS.




FIGURE 8 Scatterplot of opportunity cost rate and net margin (both in per ha and year) for each site with active AFS. Size of AFS reflected in point size.

As expected net margins of AFS depend heavily on a site's opportunity cost rate showing a clear negative trend. Moreover, it appears that AFS sizes vary quite heavily between 86 and 2794 hectare. Probably very small sites won't operate efficiently because cost for establishment, harvesting, and maintenance are assumed to scale with size, but are rather fixed below a certain critical threshold.

5 | CONCLUSION

This paper has developed and applied the AFS-SCD model, a MILP formulation for the integrated design and operation of agroforestry biomass supply chains. The model endogenises stand age dynamics through binary arc-based age-lag variables, couples component-specific Gompertz yield coefficients for stem, branch, and residue fractions with multi-echelon logistics decisions, and embeds a product quality cascade that allocates biomass grades to chemical, pulp, and energy markets within a unified optimisation framework. The formulation is, to the best of our knowledge, the first supply chain model to simultaneously optimise AFS establishment timing, consecutive harvest scheduling under rotation-age constraints, and product-grade allocation across spatially heterogeneous consumer markets.

The case study for the Saxony-Anhalt tri-state region demonstrates the practical relevance of the approach. The optimal supply chain concentrates AFS establishments in the Elbe and Saale floodplains, where high biomass productivity and proximity to large pulp mills create the most favourable economics.

The sensitivity analysis confirms that profitability depends primarily on achievable gate prices for high-quality (P1) biomass grades: scenarios combining reduced revenue with high opportunity costs yield negative net profits, whereas scenarios with full or elevated prices remain viable under cost pressure. Opportunity cost is the dominant site-level discriminator, with parcels exceeding approximately 300 ha⁻¹ yr⁻¹ rarely selected in the optimal solution. These findings underscore the importance of product cascading as a strategic lever for AFS viability: maximising the share of stem biomass directed to high-value chemical and pulp applications substantially improves profitability compared to energy-only offtake strategies.

Methodologically, the AFS-SCD model bridges the gap between stand-level growth modelling and regional supply chain optimisation identified in the literature (Zahraee et al., 2020; Lamers et al., 2011). The age-lag constraint structure generalises to other perennial biomass crops with periodic coppice cycles, such as willow or black locust, and to long-rotation silvoarable systems in which rotation length is a strategic variable. The model is extensible to stochastic settings by replacing deterministic yield coefficients with scenario-weighted counterparts or by incorporating robust optimisation over yield uncertainty intervals (Sharma et al., 2013).

Several limitations point to directions for future research. First, consumer demand is treated as exogenous and time-invariant; endogenising demand responses to biomass price changes would require a partial-equilibrium extension. Second, the case study uses a single allometric scenario calibrated to Central European poplar; propagating Gompertz parameter uncertainty through to supply chain outcomes would strengthen the robustness analysis. Third, environmental co-benefits of AFS — including carbon sequestration, nitrate leaching reduction, and biodiversity enhancement — are not monetised in the current objective. Integrating life cycle assessment indicators as additional objectives or constraints would support a more comprehensive sustainability evaluation, consistent with the cleaner production framing of this work.

AUTHOR CONTRIBUTIONS

The author is responsible for all contents of this text.

ACKNOWLEDGMENTS

The author thanks all partners in the aforementioned project for their help in compiling and reviewing this manuscript in its early stages. Special thanks are due to Nico Tauchnitz, David Anthony, Jan Lukas Wenzel und Christopher Conrad.

FINANCIAL DISCLOSURE

This research was funded by the German Federal Ministry of Education and Research (BMBF) within the programme “Modellregion der Bioökonomie (DiP)” under the project “DiP: SMART-Agroforst – Digitalisierung zur Förderung der Etablierung von Agroforstsystemen auf der Landschaftsebene als Beitrag zur Klimaresilienz Süd-Sachsen-Anhalts und Dekarbonisierung seiner chemischen Industrie – Teilvorhaben J” (funding code 031B1457J). The funder had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

CONFLICT OF INTEREST

The author declares no potential conflict of interests.

References

- Arabi, M., Yaghoubi, S., & Tajik, H. (2019). Optimal design of a microalgae-based biofuel supply chain with co2 reduction and hydrogen production. *Computers & Chemical Engineering*, 130, 106530. doi: 10.1016/j.compchemeng.2019.106530
- Barrio-Anta, M., Sixto-Blanco, H., Cañellas-Rey de Viñas, I., & Castedo-Dorado, F. (2008). Dynamic growth model for short-rotation poplar plantations in northern and central-western Spain. *Forest Ecology and Management*, 255(3–4), 1257–1268. doi: 10.1016/j.foreco.2007.10.026
- Cambero, C., & Sowlati, T. (2014). Assessment and optimization of forest biomass supply chains from economic, social and environmental perspectives—a review of literature. *Renewable and Sustainable Energy Reviews*, 36, 62–73. doi: 10.1016/j.rser.2014.04.041
- Civitarese, V., Acampora, A., Sperandio, G., Assirelli, A., & Picchio, R. (2019). Production of wood pellets from poplar trees managed as coppices with different harvesting cycles. *Energies*, 12(15), 2973. Open access. doi: 10.3390/en12152973
- De Meyer, A., Cattrysse, D., Rasinmäki, J., & Van Orshoven, J. (2014). Methods to optimise the design and management of biomass-for-bioenergy supply chains: A review. *Renewable and Sustainable Energy Reviews*, 31, 657–670. doi: 10.1016/j.rser.2013.12.036
- De Meyer, A., Cattrysse, D., & Van Orshoven, J. (2016). A generic mathematical model to optimise strategic and tactical decisions in biomass-based supply chains (OPTIMASS). *European Journal of Operational Research*, 249(1), 293–307. doi: 10.1016/j.ejor.2015.08.003

- Deutscher Fachverband für Agroforstwirtschaft (DeFAF) e.V. (2026, February). *Annual review of the Agroforestry Map of Germany 2025*. Retrieved from <https://agroforst-info.de/wp-content/uploads/2026/02/DeFAF-2026-Annual-review-of-the-Agroforestry-Map-of-Germany-2025.pdf> Available at: <https://agroforst-info.de/wp-content/uploads/2026/02/DeFAF-2026-Annual-review-of-the-Agroforestry-Map-of-Germany-2025.pdf>.
- Ekiolu, S. D., Acharya, A., Leightley, L. E., & Arora, S. (2009). Analyzing the design and management of biomass-to-biorefinery supply chain. *Computers & Industrial Engineering*, 57(4), 1342–1352. doi: 10.1016/j.cie.2009.07.003
- Espinoza, V., Bautista, S., Narvaez, P. C., Alfaro, M., & Camargo, M. (2017). Sustainability assessment to support governmental biodiesel policy in Colombia: A system dynamics model. *Journal of Cleaner Production*, 141, 1145–1163. doi: 10.1016/j.jclepro.2016.09.168
- Faasch, R. J., & Patenaude, G. (2012). The economics of short rotation coppice in Germany. *Biomass and Bioenergy*, 45, 27–40. doi: 10.1016/j.biombioe.2012.04.012
- Graves, A. R., Burgess, P. J., Liagre, F., Pisanelli, A., Paris, P., Moreno, G., ... Dupraz, C. (2007). Development and application of bio-economic modelling to compare silvoarable, arable, and forestry systems in three European countries. *Ecological Engineering*, 29(4), 434–449. doi: 10.1016/j.ecoleng.2006.09.018
- Grünewald, H., Böcker, R., Schneider, B. U., Bens, O., & Hüttl, R. F. (2007). Biomass production of black locust (*Robinia pseudoacacia* L.) short rotation plantations in Germany: growth and economics of different clones and management practices. *Bioresource Technology*, 98(2), 281–286. doi: 10.1016/j.biortech.2006.01.004
- Hildebrandt, J., O’Keeffe, S., Bezama, A., & Thrän, D. (2019). Revealing the environmental advantages of industrial symbiosis in wood-based bioeconomy networks: An assessment from a life cycle perspective. *Journal of Industrial Ecology*, 23(4), 808–822. doi: 10.1111/jiec.12818
- Huber, G., Amann, C., Hülshbergen, K.-J., & Schmid, H. (2018). Yield potential of tree species in organic and conventional short-rotation agroforestry systems in Southern Germany. *Heliyon*, 4(1), e00543. doi: 10.1016/j.heliyon.2018.e00543
- Huber, G., Hülshbergen, K.-J., & Schmid, H. (2016). Allometric equations for above-ground tree-biomass estimation in Southern German short-rotation agroforestry systems. *Bioenergy Research*, 9(3), 955–968. doi: 10.1007/s12155-016-9744-z

- Jha, K. K. (2018). Biomass production and carbon balance in two hybrid poplar (*populus euramericana*) plantations raised with and without agriculture in southern France. *Journal of Forestry Research*, 29(6), 1689–1701. doi: 10.1007/s11676-017-0555-3
- Kira, T., Shinozaki, K., & Hozumi, K. (1961). The C-D rule, its theory and practical uses. (Intraspecific competition among higher plants. X.). *Journal of Biology, Osaka City University*, 12, 69–82.
- Lamers, P., Junginger, M., Hamelinck, C., & Faaij, A. (2011). Developments in liquid biofuels commodity trade. *Biofuels, Bioproducts and Biorefining*, 5(4), 403–437. doi: 10.1002/bbb.303
- Malladi, K. T., & Sowlati, T. (2018). Biomass logistics: A review of important features, optimization modelling and the new trends. *Renewable and Sustainable Energy Reviews*, 94, 587–599. doi: 10.1016/j.rser.2018.06.052
- Mansoornejad, B., Pistikopoulos, E. N., & Stuart, P. R. (2013). Integrating forest biorefinery metrics into supply chain design and analysis. *Forest Policy and Economics*, 26, 43–54. doi: 10.1016/j.forpol.2012.08.003
- Nair, P. K. R. (1993). *An introduction to agroforestry*. Dordrecht: Kluwer Academic Publishers.
- Niemczyk, M., & Thomas, B. R. (2021). Does intensive management affect wood density and fiber properties of hybrid poplar grown in short-rotation plantation? *Forests*, 12(4), 399. doi: 10.3390/f12040399
- Paula, R. R., Reis, G. G., Reis, M. G. F., Oliveira Neto, S. N., Leite, H. G., Melido, R. C. N., ... Souza, F. C. (2013). Eucalypt growth in monoculture and silvopastoral systems with varied tree initial densities and spatial arrangements. *Agroforestry Systems*, 87(6), 1295–1307. Retrieved from <https://doi.org/10.1007/s10457-013-9638-5> doi: 10.1007/s10457-013-9638-5
- Sharma, B., Ingalls, R. G., Jones, C. L., & Khanchi, A. (2013). Biomass supply chain design and analysis: Basis, overview, modeling, challenges, and future. *Renewable and Sustainable Energy Reviews*, 24, 608–627. doi: 10.1016/j.rser.2013.03.049
- Spinelli, R., Nati, C., & Magagnotti, N. (2009). Using modified foragers to harvest short-rotation poplar plantations. *Biomass and Bioenergy*, 33(5), 817–821. doi: 10.1016/j.biombioe.2009.01.002
- Testa, R., Di Trapani, A. M., Foderà, M., Sgroi, F., & Tudisca, S. (2014). Economic evaluation of introduction of poplar as biomass crop in Italy. *Renewable and Sustainable Energy Reviews*, 38, 498–506. doi: 10.1016/j.rser.2014.07.007
- Tjørve, K. M., & Tjørve, E. (2017). The use of gompertz models in growth analyses, and new gompertz-model approach: An addition to the unified-richards family. *PloS one*, 12(6).

- Toensmeier, E. (2017). The carbon farming solution: A global toolkit of perennial crops and regenerative agriculture practices for climate change mitigation and food security. In *The carbon farming solution*. White River Junction, VT: Chelsea Green Publishing.
- Trnka, M., Fialová, J., Koutecký, V., Fajman, M., alud, Z., & Havlíek, M. (2008). Biomass production and survival rates of selected poplar clones grown under a short-rotation system on agricultural land. *Plant, Soil and Environment*, 54(2), 78–88. doi: 10.17221/2752-PSE
- Yue, D., You, F., & Snyder, S. W. (2014). Biomass-to-bioenergy and biofuel supply chain optimization: Overview, key issues and challenges. *Computers & Chemical Engineering*, 66, 36–56. doi: 10.1016/j.compchemeng.2013.11.016
- Zahraee, S. M., Shiwakoti, N., & Stasinopoulos, P. (2020). Biomass supply chain environmental, economic, and social sustainability: A systematic literature review. *Industrial Crops and Products*, 145, 112144. doi: 10.1016/j.indcrop.2020.112144

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Thomas Kirschstein. Strategic Planning of Agroforestry Supply Chains. *Preprint submitted to Journal of Industrial Ecology* 2026;():.

APPENDIX

TABLE 1 Notation for the biomass growth and harvest yield models.

Symbol	Description
$M(t)$	Total above- and belowground dry biomass per ha of AFS (t DM ha^{-1}) at stand age t
$A(N)$	Stand-level Gompertz asymptote (t DM ha^{-1}); density- and site-dependent
k	Gompertz growth-rate coefficient
t_0	Inflection age of stand biomass accumulation (years)
N	Stand density (trees ha^{-1})
C_{site}	Site-quality constant ($\text{kg ha}^{\beta-1}$); conservative: 4 475, good site: 6 471
β	Competition-density exponent; fitted $\hat{\beta} = 0.380$
$f_p(t)$	asymptotic fraction of AGB attributed to compartment $p \in \{s, b, r\}$ at age t
$q_p(t)$	Merchantable fraction of compartment p biomass at age t (logistic)
r_p	Logistic growth rate for merchantability of compartment p
$t_{50,p}$	Age at which 50% of compartment p biomass is merchantable (yr)
$\eta_p(t)$	Merchantable dry biomass of compartment p per ha at harvest age t (t DM ha^{-1})
ω	Moisture content at felling (mass fraction); $\omega = 0.55$




FIGURE 1 Logistic diameter-class merchantability functions $q_p(t)$ for stem ($d > 15$ cm, green) and branches ($d > 7$ cm, orange); the 50% threshold ages t_{50} are indicated by dashed vertical lines.




FIGURE 2 Tree-level and stand-level Gompertz asymptote functions $A_{tree}(N)$ and $A(N)$ for advanteguous ($N = 4$, $\beta = -0.3$) and disadvantageous ($N = 6.5$, $\beta = -0.5$) site conditions.

TABLE 2 Notation for the MILP model.

Sets & Indices	
$i \in \mathcal{I}$	Agroforestry sites
$j \in \mathcal{J}$	Storage / pre-treatment facilities
$k \in \mathcal{K}$	Consumer sites
$p \in \mathcal{P}$	Product types (stem, branches, residues)
$(p, p') \in \mathcal{Q}$	Admissible product-substitution pairs (quality cascade)
$\mathcal{T} = \{1, \dots, T\}$	Planning periods (years)
$\mathcal{T}^{\text{harv}} = \{A^{\min} + 1, \dots, T\}$	Feasible harvest periods
$\mathcal{S}^{\text{est}} = \{(0, t) \mid t \in \{1, \dots, T - A^{\min}\}\}$	Establishment arcs: dummy source to period t
$\mathcal{S}^{\text{harv}} = \{(s, t) \mid s, t \in \mathcal{T},$ $t - s \in \{A^{\min}, \dots, A^{\max}\}\}$	Harvest cycles within age-lag bounds
$\mathcal{S}^{\text{term}} = \{(s, T + 1) \mid$ $s \in \{A^{\min} + 1, \dots, T\}\}$	Termination arcs to dummy sink $T + 1$
$\mathcal{S} = \mathcal{S}^{\text{est}} \cup \mathcal{S}^{\text{harv}} \cup \mathcal{S}^{\text{term}}$	Age-lag arc set (establishment $\cup$ harvest $\cup$ termination)
Decision Variables	
$z_{ist} \geq 0$	Area of site i with AFS operation between s and t
$X_{ijpt} \geq 0$	Transport flow (t) of p from site i to facility j in t
$X_{jkpp't} \geq 0$	Flow of p from facility j to consumer k satisfying demand p' in t
$S_{jpt} \geq 0$	Inventory (t) of product p at facility j end of period t
Parameters	
T	Length of planning horizon (years)
$A^{\min}, A^{\max}$	Minimum / maximum stand age at harvest (years)
AREA_i	Area of site i (ha)
η_{pa}	Yield coefficient (t/ha) of fraction p at stand age a , derived from (4)
R_{kp}	Revenue (/t) for product p at consumer k
C_i^{est}	Establishment cost (/ha) at site i
C_i^{opp}	Annual opportunity cost (/ha) at site i
C_i^{harv}	Harvest & refitting cost (/ha) at site i
$c^{\text{tr-raw}}$	Transport cost rate (/t km) for raw biomass
$c^{\text{tr-pre}}$	Transport cost rate (/t km) for pre-treated biomass
c_j^{stor}	Storage holding cost (/t) at facility j
d_{ij}, d_{jk}	Distance (km) between nodes
$\text{CAP}_j^{\text{stor}} / \text{CAP}_j^{\text{proc}}$	Storage and processing capacity (t) at facility j
D_{kpt}^{max}	Maximum demand (t) of fraction p at consumer k in period t

TABLE 3 Consumer demand quantities and gate prices by product grade. P1: chemical/pulp; P2: sawlog/pulpwood; P3: energy/pellets. Demands in kt wood input per year; prices in /t wood input at the factory gate.

Consumer	Annual demand			Gate price		
	P1 (kt/yr)	P2 (kt/yr)	P3 (kt/yr)	P1 (/t)	P2 (/t)	P3 (/t)
consumer 1	300.0			50		
consumer 2	100.0			60		
consumer 3		300	50.0		37.5	24
consumer 4	200.0			47.5		
consumer 5		30	5.0		35	21
consumer 6		15	5.0		32.5	21
consumer 7		10	3.0		30	18
consumer 8		5	8.0		27.5	18
consumer 9	2.0	50	20.0	40	35	22.8
consumer 10			1.7			21
consumer 11			2.9			21
consumer 12			2.5			21
consumer 13			2.3			21
consumer 14			1.9			21
consumer 15			2.7			21
consumer 16			1.7			21
consumer 17			1.9			21
consumer 18			1.6			21
consumer 19			1.5			21
consumer 20			1.5			21
consumer 21			1.8			21
consumer 22			3.5			21
consumer 23			1.6			21
consumer 24			2.1			21
consumer 25			4.5			21
consumer 26			1.8			21
consumer 27			3.2			21
consumer 28	18.5			27.5		





FIGURE 3 AFS supply chain network in Saxony-Anhalt. Grey lines: German federal state boundaries. Blue lines: major rivers. Coloured polygons: potential AFS parcels grouped by site (colours encode cluster membership; no legend). Orange symbols: pre-treatment facilities (circle = sawmill, triangle = logistics hub, size proportional to processing capacity). Coloured squares: consumer sites (violet: P1 chemical/pulp, blue: P2 pulp/paper, red: P3 energy/biogas, size proportional to total demand).

TABLE 4 Parameter values for the AFS-SCDcase study. Transport costs refer to road distance

Parameter	Description	Base-case value	Unit
Index sets			
$ T $	Planning periods	40	yr
$ Z $	Sites	145	
$ J $	Pre-treatment / storage nodes	11	
$ K $	Consumers	28	
$ P $	Product grades	3	
Biomass dynamics			
$A^{\min}$	Minimum harvest age (years)	3	yr
$A^{\max}$	Maximum stand age (years)	20	yr
Cost parameters			
c^{tr-raw}	Raw transport cost	0.08	$t^{-1} km^{-1}$
c^{tr-pre}	Pre-processed transport cost	0.06	$t^{-1} km^{-1}$
c_i^{est}	Establishment cost rate	2500	$ha^{-1} yr^{-1}$
c_i^{harv}	harvest cost rate	150	$ha^{-1} yr^{-1}$
c_i^{main}	Maintenance cost rate	10	$ha^{-1} yr^{-1}$
c_i^{opp}	aver. opportunity cost	521	$ha^{-1} yr^{-1}$
Revenue parameters			
R_{P1}	Mean gate price P1	45	t^{-1}
R_{P2}	Mean gate price P2	33	t^{-1}
R_{P3}	Mean gate price P3	21	t^{-1}

**FIGURE 4** Total production quantity of biomass types and its industrial usage over planning horizon (in tons).