

# Dynamics of Fluids in Random Environments

Thomas Konincks

Master Theoretical Chemistry and  
Computational Modeling

September 2014



rijksuniversiteit  
 groningen

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Experimental works</b>	<b>3</b>
2.1	Dynamics of colloids in a random optic potential . . . . .	3
<b>3</b>	<b>Theoretical models</b>	<b>7</b>
3.1	Random walk . . . . .	8
3.2	Monte-Carlo method . . . . .	9
3.3	Molecular Dynamics . . . . .	10
3.4	Mode-Coupling theory . . . . .	11

## 1 Introduction

In the recent few years, porous materials have attracted much attention, especially because of the possibility of hydrogen storage, for which the MOF materials are the most viewed.<sup>18</sup> MOFs are an example of a porous solid that presents a very high symmetry, due to their crystalline structure. However in this work we will be concerned with an other type of porous medium: the disordered porous materials (i.e. without any symmetry or long distance order), with randomly arranged pores.

This type of materials contains most of the natural porous systems: some rocks are known to be porous (sand, gravel and volcanic rocks for example), the soil has also some degree of porosity that is extensively studied in order to make good use of natural resources, and understand geological phenomenons.<sup>16,6,5,17</sup> But porosity can also be wanted, created and controlled: the controlled porous glasses are a good example, as well as the novel aerogel materials that present an exceptionally high degree of porosity.

Dynamics of liquids in confined disordered environment is a kind of system that is difficult to capture experimentally, because of the speed of the dynamics and the fact that the system is by definition confined. Some experiments have given some interesting results on comparable systems, but the main way to study them is to develop models that are able to compute information on their dynamics. Experimentally or theoretically, there are essentially two phenomenas that people want to capture:

- First, the liquid-glass transition. If the dynamics are slow enough (i.e. the potential in which the particle moves: the matrix), the system will undergo a phase transition, meaning that the colloids will be completely trapped inside

the potential and the dynamics stopped. The glass phase is simply a liquid which dynamics are "frozen".

- The second phenomenon that has been observed in this kind of confined environments is the three step relaxation that is characteristic of supercooled liquids. High temperature and non confined fluids follows a simple relaxation scheme as shown in fig. 1 (left): when the time increases, the system goes smoothly from a non-relaxed state to a state where all the distances have been balanced, and the potential energy minimized. For confined fluids, the relaxation is done in three steps: one first simple relaxation (I) that follows the usual dynamics of the fluids (as for the non-confined case) where the particles explore their local minima, the second step (II): a plateau and a second relaxation during which the particles appear trapped in the cages formed by the matrix of the solid, this step is called the  $\alpha$ -relaxation regime. The third step is the  $\beta$ -relaxation regime, a very slow approach towards the long time diffusion regime, during which the particles will be able to hop from a site to another, their dynamics can then be compared to a random walk from site to site.




Figure 1: Left: Exponential decay for a normal liquid. Right: supercooled liquids do not have such a simple decay. Notice the logarithmic scale. D. Reichman and P. Charbonneau, *J. Stat. Mech.: Theory and Experiment*, **05**, P05013 (2005).

Studying the dynamics of fluids in disordered environments allows to make predictions on various types of systems such as supercooled liquid or glasses, but also as living cells or protein foldings.

Section 3 will make a summary of the different theories and models that have been developed in order to understand and make predictions on the dynamics and the phase transitions that these trapped system undergo. A special emphasis will be put on the mode-coupling theory that is to this date the most efficient theory, and one that has made the most accurate predictions. The next section will present some of the experimental works that have been made in the field.

## 2 Experimental works

Having experimental evidence of phenomenas that happen at the microscopic level, confined into a porous material is very difficult, not to say impossible. However scientists have found ways to observe macroscopic phenomena that have the exact same behaviour than molecules inside a porous matrix. The experiments have been made in one, then in two dimensions. Statistical mechanics is one of the few fields in which the validity of the experiments have to be compared to theory, and not the opposite. Thus, each of the experiments have been "dubbed" with a Monte-Carlo simulation to compare the results (see sec. 3.2).

### 2.1 Dynamics of colloids in a random optic potential

The diffusion of particles in a disordered environment can be seen as a diffusion of those particles in an potential energy landscape, that has a finite value if the pores are not permanent, or goes to infinity when the particles touches edges of pores in a hard matrix. Some colloids are known to have an interaction with light, either attractive or repulsive (this force is lead by the difference of refractive index, higher or lower than that of the environment, leads to respectively an attractive or a repulsive force).

That observation has lead Hanes *et al.*<sup>7</sup> to conduct an experiment in which they let polystyrene colloids suspended in heavy water (D<sub>2</sub>O) flow in a channel. The channel is then randomly hit by laser beams that creates a one dimensional random potential that the particles feels.

An other experiment, made by Evers *et al.*<sup>4</sup> focused on a two dimensional description of the dynamical phenomenons. Instead of letting the colloids flow in a channel, they sent a random light interference figure (called speckle) on polystyrene particles floating in heavy water (as argued vy Evers *et al.* the heavy water allows the particles to cream rather than sediment). The speckle simulates the random arrangement of a physical matrix. The experiment has simply been extended to one more dimension compared to the one of Hanes *et al.*.

Fig. 2 (left) shows a potential that was used by Evers *et al.* to make a Monte-Carlo simulation in order to complete their experimental work (see sec. 3.2). The potential is supposed to be very close to the potential that they produced experimentally. It clearly presents maximas, minimas and saddle points. Knowing that the colloids have a repulsive interaction with light, they will tend to stay at the minimas and avoid the maximas. The colloids will however be able to hop from one minima to another if the energy barrier is low (i.e. saddle points are low) enough or if the colloids are fast (i.e. the temperature is high) enough.

Fig. 2 (right) shows a picture of the trajectories of the colloids in the system studied by Evers *et al.*. The picture is taken with a CMOS camera (usual camera). We can see that the colloids that are trapped inside the speckle (green circle) do not move as much as the colloids outside the speckle do, which can freely explore all the space available to them. The particles inside the speckle remain trapped for a more or less long time inside the minimas of the potential, sometimes hopping from a minima to the other (that behaviour can be compared to a random walk, see sec.3.1). The presence of the random potential slows down the dynamics of the system drastically.




Figure 2: (left) Potential generated for a Monte-Carlo simulation, it is supposed to reflect the potential felt by the colloids beamed by a laser speckle. (right) Trajectories of particles undergoing diffusion in a two-dimensional plane, part of which contains a random potential (green background), which is separated by a barrier (white/green rings) from the surroundings (white background). F. Evers *et al.*, *Physical Review E*, **88**, 022125 (2013)

Fig. 3 shows the mean square displacement and the normalized diffusion coefficient for the experiment conducted by Hanes *et al.* (left) and the experiment by Evers *et al.*. The mean square displacement measures how much the particles

have moved from their initial position in function of time, it is calculated as:

$$\langle \Delta r(t)^2 \rangle = \sum_{i=1}^d \langle x_i(t)^2 \rangle - \langle x_i(t) \rangle^2 \quad (1)$$

The normalized diffusion coefficient:

$$\frac{D(t)}{D_0} = \frac{1}{2dD_0} \frac{d}{dt} \langle \Delta r(t)^2 \rangle \quad (2)$$

In both equations,  $d$  is the dimension and the  $\langle \dots \rangle$  represents the time average.




Figure 3: Top pictures: the mean square displacement in function of the time. Bottom picture: the normalized diffusion function in function of the time.

Left: R. D. L. Hanes et al., *Soft Matter*, **8**, 2714 (2012)

Right: F. Evers et al., *Physical Review E*, **88**, 022125 (2013)

The mean square displacement presents two variations of slope: at early times the behaviour is the same than for the system without external potential ( $\varepsilon = 0$

in the picture from Evers *et al.*), the slope is equal to one, indicating a normal diffusion, at intermediate times the slope becomes lower (this happens earlier when the roughness is higher), the system is then said to have a subdiffusive behaviour, and for long times it returns to its initial value and follows again the slope of the  $\varepsilon = 0$  curve.

The same kind of behaviour can be seen on the normalized diffusion coefficient: at small times the value is very close to one, meaning that the system follows a normal diffusive behaviour, then at intermediate times the value of  $D(t)/D_0$  decreases dramatically, finally at large times the system regains a diffusive behaviour, reaching a  $D_\infty$  value.

These two physical quantities give precious information about the dynamics of the confined systems in function the time: at small times the system follows the behaviour of a non-confined fluid ( $\langle \Delta r(t)^2 \rangle$  and  $D(t)/D_0$  have a diffusive behaviour), at intermediate times the system behave like it is trapped in the cages of the matrix and does not move as much as at earlier times (slopes of  $\langle \Delta r(t)^2 \rangle$  and  $D(t)/D_0$  decrease), finally at large times, the system follows again a diffusive behaviour like at the beginning. This can be put in comparison with fig. 1. Each of the observed regimes can be assigned to a relaxation regime: at early times the particles undergo a diffusive behaviour inside their local minima, then they start to hop from a minima to another until the system is under equilibrium, then they start to hop from a minima to another, which can be assimilated to a diffusive behaviour again.

The Van Hove function is the probability density of finding a particle  $i$  at the position  $r$  at a time  $t$ , knowing that a particle  $j$  is at the origin at  $t=0$ . It is expressed as:

$$G(\Delta r, t) = \frac{1}{N} \left\langle \sum_{i=1}^N \sum_{j=1}^N \delta(r - r_i(t) + r_j(0)) \right\rangle \quad (3)$$

The Van Hove correlation function can then be split into two part: the distinct and the self parts:

$$G_s(\Delta r, t) = P(\Delta r, t) = \frac{1}{N} \left\langle \sum_{i=1}^N \delta(r - r_i(t) + r_i(0)) \right\rangle \quad (4)$$

$$G_d(\Delta r, t) = \frac{1}{N} \left\langle \sum_{j \neq i}^N \delta(r - r_j(t) + r_j(0)) \right\rangle \quad (5)$$

The physical interpretation of the self part of the Van Hove function is the probability density of finding a particle  $i$  at the time  $t$ , knowing that this particle was at the origin at  $t=0$ . In other words, it the probability that this particle has

moved of a distance  $r$  during the time  $t$ . The distinct part of the Van Hove function is the probability that a particle  $j$  different from  $i$  at a distance  $r$  at the time  $t$ , knowing that the particle  $i$  was at the origin at  $t=0$ .

The self part of the Van Hove function is often calculated during experiments or simulations because it gives precious information about the dynamics of the system, as we can see in fig. 4.




Figure 4: Self part of the Van Hove function *i*-for different waiting times (left) and for different roughness (right).

*F. Evers et al., Physical Review E, 88, 022125 (2013)*

From fig. 4, we can see that the presence of a potential changes qualitatively the dynamics of the particles: the black curve at the right shows the self part of the Van Hove function without any potential and the green curve shows the same quantity with a roughness of 2.8. The behaviour is qualitatively different: when there is an external potential the particle explores less space, obviously because it remains trapped inside the minimas. Comparing the other curves we can say that, the less the waiting time, the less the particle can explore space, and the higher the roughness, the more the particle is trapped in the minima and the less it can move from its original position (this is clearly seen from the narrowing of the function for small times or high roughness).

### 3 Theoretical models

This section will make a brief summary of the different models that have been developed in order to tackle the problem of the diffusion of particles in a disordered environment. The aim here is not to derive exactly every equation that makes



the model, but simply to give a brief overview of the idea, problems and eventual results of these models.

### 3.1 Random walk

A simple way to see the diffusion in a disordered matrix is to model the dynamics of the particles by a random walk with constraints. This method is efficient when system where the particles are allowed to cross barriers (biological, magnetic systems...) Two models have emerged from this idea: the random trap model<sup>9</sup> and the random barrier model.<sup>2,11</sup> The two models describe the diffusion from a one dimensional point of view, a simple model indeed, but precise enough to understand how diffusion works. The two models are not in concurrence but present a duality relation through their master operators.

These two models are based on the assumption that the particle is Brownian, that is it has a random path through space. This type of behaviour is naturally very well described with a random walk.

Consider a single particle hopping on a chain of sites. The particle on site  $n$  may hop either to the right  $n + 1$ , or to the left  $n - 1$ . In the random trap model, each hop happen with a site-dependent rate  $W_n$ . Thus, the equation describing the dynamics of the random trap model is:

$$\frac{dp_n^T(t)}{dt} = W_{n+1}p_{n+1}^T(t) + W_{n-1}p_{n-1}^T(t) + 2W_n p_n^T(t) \quad (6)$$

In the random barrier model, the hopping rate is not site-dependent but barrier-dependent, that is dependent on the link between the sites:

$$\frac{dp_n^B(t)}{dt} = W_n [p_{n+1}^B(t) - p_n^B(t)] + W_{n-1} [p_{n-1}^B(t) - p_n^B(t)] \quad (7)$$

The two models use the same set of rates  $W_n$ , then the fluctuation of the barrier model  $j_n^B = W_n (p_N^B - p_{N+1}^B)$  and the rescaled probability density of the trap model  $W_n p_N^T$  obey the same differential rules. It follows that when using the operator notation of  $W_n$ , the spectrum of eigenvalues of both models will be the same. This is called the trap-barrier duality. A simple way to understand it is to identify the regions between large barriers in the so-called model model with the deep traps in the trap model. This vision is not perfectly exact but has the advantage of clearly showing the kind of duality that links these two models. One can refer to the publication of Jack and Sollich about the barrier-trap duality<sup>10</sup> for the exact derivation.

RESULTS!!!!!!!!!!!!!!!!!!!!!!

### 3.2 Monte-Carlo method

The Monte-Carlo method is a stochastic method that focuses on computing physical quantities by making a large number of calculations using random numbers, eventually averaging the results. This method is particularly suited in the kind of systems that were studied experimentally by Hanes *et al.* and Evers *et al.* (resumed in sec. 2, which precisely made MC calculations to complete their work.

The particle evolves in a computer-generated random potential. The simulations were made either by putting the particles randomly distributed in the potential energy surface (instantaneous quench of the system), or letting the particles relax and reach an equilibrium position (potential occupied following Boltzmann statistics). At each iteration a particle is randomly chosen, along with a direction. Then, the corresponding energy for a step in that direction is calculated from the particle's current position on the energy landscape. The move is accepted if the energy needed to make that step is low enough. It can also be put into a probability that depends on the energy of the step.

Fig. 5 shows the mean square displacement (left) and the normalized diffusion coefficient (right) as obtained by MC simulations by Hanes *et al.* in the simulation study of the particles in a one dimensional random potential.<sup>8</sup>




Figure 5

From fig. 5 and referring to sec. 2 we can see that the results are in good agreement with the experiments. In particular, the mean square displacement (left) shows three different regimes: at first the slope is close to one, meaning that the particle has a diffusive behaviour. When the roughness is high enough, the slope decreases, the system has a subdiffusive behaviour: the particles undergo diffusion inside their local minima without escaping it. When the waiting time is long enough, the diffusive behaviour reappears and the particles start to hop from a minima to another. The same behaviour can be seen on the normalized diffusion coefficient (right): after a short stagnation at  $D_0$ , the slope decreases strongly until

it reaches a plateau value  $D_\infty$  (horizontal lines). These three behaviours again describe the three relaxation processes: diffusion,  $\alpha$ -relaxation and  $\beta$ -relaxation.

### 3.3 Molecular Dynamics

Molecular dynamics is one of the most used methods when making calculations about large systems of molecules or macromolecules. It is often used in drug design when it comes to model large proteins (often in addition with quantum mechanical methods, the so-called QM/MM). Molecular mechanics (MM), as expected, models the behaviour of molecular systems using the classical equations of motion (instead of quantum mechanical equations). This makes MM a very fast method by which thousands of atoms can be treated. It is also very adapted for thermodynamical and statistical systems, like for example liquids in confined matrices.

A study from Chang *et al.*<sup>3</sup> published in The American Physical Society presents a MM calculation of a binary mixture of hard spheres. One component is mobile and will undergo the dynamics, while the other component is fixed in space and time (similar to the QA mixture in sec. 3.4). The algorithm calculates the velocity of each particle, and the new velocity after each collision.. Fig.6 shows the mean square displacement in function of time, and the normalized diffusion coefficient in function of the matrix/fluid ratio, as calculated with the MM algorithms.

But before any analysis we have to define some quantities that are used in this publication. Since the matrix and the fluid are made of hard sphere particles, we can define  $\sigma_m m$  as the matrix particle diameter,  $\sigma_f f$  as the fluid particle diameter and  $\sigma_{fm} = \sigma_{mf}$  as the typical particle-matrix distance. The matrix and fluid volume fractions are defined as:

$$\phi_{f/m} = \frac{\pi N \sigma_{ff/mm}^3}{6L^3} \quad (8)$$

As stated in sec. 2, a system with a mean square displacement that follows a line with a slope equal to one has a diffusive behaviour. If the slope is lower than one, the system has a subdiffusive behaviour, and that is what happens when  $\phi_m$  has a value around 0.25.

Fig. 6 shows the normalized diffusion coefficient  $D(t)/D_0$  as a function of  $\phi_f$  for different values of  $\sigma_{ff}$  for a fixed value of  $\sigma_{mm} = 1.0$ . Therefore  $\phi_m$  decreases as  $\phi_f$  increases.

Something apparently strange happens: the curve has a qualitatively different behaviour when the ratio  $\phi_{ff}/\phi_{mm}$  changes. When  $\sigma_{ff}$  is greater than  $\sigma_{mm}$ , the diffusion decreases as the size of the fluid particle increases. But when  $\sigma_{ff}/\sigma_{mm} < 1.0$ ,




Figure 6: (left) Mean square displacement in function of time, the different curves show different matrix volume fractions  $\phi_m$ . (right) Normalized diffusion coefficient in function of the fluid volume fractions, the different curves show different  $\phi_f/\phi_m$  ratios.

the diffusion decreases when  $\sigma_{ff}$  increases.

This can be understood as: when  $\sigma_{ff}$  is greater than  $\sigma_{mm}$ , replacing a large fluid particle by several smaller immobile particles will create what is called a free volume. Free volume appears when matrix particles are so close to each other that some of the space is not available for fluid to flow in. This is why the diffusion decreases. On the other hand, when  $\sigma_{ff}$  is smaller than  $\sigma_{mm}$ , replacing several small fluid particles with a large matrix immobile one will tend to reduce the free space, and increasing the diffusion.

### 3.4 Mode-Coupling theory

The Mode-Coupling Theory (MCT), developed by Wolfgang Gotze *et al.*,<sup>1</sup> has emerged as one of the most accurate models for the dynamics of liquids, such that results from experiments or other simulations are often compared to MCT. The success of MCT lies in its ability to reproduce reliably important phenomena such as the slowing down of the dynamics of a liquid when the temperature is decreased or when density increases.

The MCT theory expresses the equation of motion, by projecting its terms on the slow modes of the system. The "slow modes" is a physical quantity that varies very slowly in times, and in the case of the confined systems, when approaching the liquid-glass transition, the density fluctuation varies less and less until it becomes stationary. Thus, in the case of the MCT applied to confined systems, the slow

modes will be the density fluctuations.  
The final MCT equation has the form:

$$\frac{\partial F(q, t)^2}{\partial^2} + \frac{q^2 k_B T}{m S(q)} F(q, t) + \int_0^t d\tau K(q, t - \tau) \frac{\partial F(q, t)}{\partial \tau} = 0 \quad (9)$$

Where  $F(q, t)$  is the correlation function,  $K(q, t)$  is the memory function and  $S(q)$  is the structure factor.

The correlation function is the function that connects the value of a the density fluctuation  $t$  with its value at time  $t+t'$ . In eq. 9, the equation of motion has been totally rewritten in function of the slow modes of the system. The memory function is the function that contains all the remaining non-slow modes, and as we can see it is contained inside an integral over time. This integral takes account for a delayed influence of the fast modes on the slow modes. The structure factor is the function containing the correlation function between the physical quantity with itself at time  $t=0$ .

The MCT has helped to make quality predictions of the liquid-glass transition on many bulk liquids, thus the idea emerged that MCT could be applied to systems presenting certain constraints. Thus, an extension for fluids confined in rough potentials has been developed by V. Krakoviack.<sup>12, 13, 14, 15</sup> The theory accounts for the presence of the disordered modeling the system by a binary mixture (called quenched-annealed mixture, QA mixture) in which one of the species would be immobilized (the quenched component), and the other component (the annealed component) would follow the MCT dynamics (see fig. 7).




Figure 7: Sketch of a QA system. In black, the immobile matrix particles. In white and gray, with arrows symbolizing their movement, the fluid particles and the tagged particle, respectively.

V. Krakowiack, *Phys. Rev. E* **79**, 061501 (2009)

## References

- <sup>1</sup> U Bengtzelius and Wolfgang Gotze. Dynamics of supercooled liquids and the glass transition. *Journal of Physics C: solid ...*, 5915, 1984.
- <sup>2</sup> J Bernasconi, HU Beyeler, S Strässler, and S Alexander. Anomalous Frequency-Dependent Conductivity in Disordered One-Dimensional Systems. *Physical Review Letters*, (March), 1979.
- <sup>3</sup> Rakwoo Chang, Kamakshi Jagannathan, and Arun Yethiraj. Diffusion of hard sphere fluids in disordered media: A molecular dynamics simulation study. *Physical Review E*, 69(5):051101, May 2004.
- <sup>4</sup> Florian Evers, Christoph Zunke, Richard D. L. Hanes, Jörg Bewerunge, Imad Ladadwa, Andreas Heuer, and Stefan U. Egelhaaf. Particle dynamics in two-dimensional random-energy landscapes: Experiments and simulations. *Physical Review E*, 88(2):022125, August 2013.
- <sup>5</sup> Tomasz Glab and Wojciech Szewczyk. Influence of simulated traffic and roots of turfgrass species on soil pore characteristics. *Geoderma*, 230-231:221–228, October 2014.
- <sup>6</sup> Claire Grosbellet, Laure Vidal-Beaudet, Virginie Caubel, and Sylvain Charpentier.

- Improvement of soil structure formation by degradation of coarse organic matter. *Geoderma*, 162(1-2):27–38, April 2011.
- <sup>7</sup> Richard D. L. Hanes, Cécile Dalle-Ferrier, Michael Schmiedeberg, Matthew C. Jenkins, and Stefan U. Egelhaaf. Colloids in one dimensional random energy landscapes. *Soft Matter*, 8(9):2714, 2012.
- <sup>8</sup> Richard D L Hanes and Stefan U Egelhaaf. Dynamics of individual colloidal particles in one-dimensional random potentials: a simulation study. *Journal of physics. Condensed matter : an Institute of Physics journal*, 24(46):464116, November 2012.
- <sup>9</sup> JW Haus, KW Kehr, and JW Lyklema. Diffusion in a disordered medium. *Physical Review B*, 25(4):4–6, 1982.
- <sup>10</sup> Robert L. Jack and Peter Sollich. Duality between random trap and barrier models. pages 1–15, October 2007.
- <sup>11</sup> Robert L Jack and Peter Sollich. Duality symmetries and effective dynamics in disordered hopping models. *Journal of Statistical Mechanics: Theory and Experiment*, 2009(11):P11011, November 2009.
- <sup>12</sup> V. Krakoviack. Liquid-glass transition of a fluid confined in a disordered porous matrix: A mode-coupling theory. *Physical Review Letters*, 94(6):5, September 2005.
- <sup>13</sup> V. Krakoviack. Liquid-glass transition of confined fluids: Some insights from a mode-coupling theory. page 7, June 2005.
- <sup>14</sup> V. Krakoviack. Mode-coupling theory for the slow collective dynamics of fluids adsorbed in disordered porous media. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, 75(3):19, October 2007.
- <sup>15</sup> V. Krakoviack. Tagged-particle dynamics in a fluid adsorbed in a disordered porous solid: Interplay between the diffusion-localization and liquid-glass transitions. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, 79(6):18, January 2009.
- <sup>16</sup> Paolo Nasta, Tamir Kamai, Giovanni B. Chirico, Jan W. Hopmans, and Nunzio Romano. Scaling soil water retention functions using particle-size distribution. *Journal of Hydrology*, 374(3-4):223–234, August 2009.
- <sup>17</sup> R.T. Walczak, F. Moreno, C. Slawinski, E. Fernandez, and J.L. Arrue. Modeling of soil water retention curve using soil solid phase parameters. *Journal of Hydrology*, 329(3-4):527–533, October 2006.

- <sup>18</sup> Yong Yan, Xiang Lin, Sihai Yang, Alexander J Blake, Anne Dailly, Neil R Champness, Peter Hubberstey, and Martin Schröder. Exceptionally high H<sub>2</sub> storage by a metal-organic polyhedral framework. *Chemical communications (Cambridge, England)*, (9):1025–7, March 2009.