Building a Design-Based Matching Pipeline: From Principles to Practical Implementation in R

Thomas Leavitt and Luke W. Miratrix

Abstract

Matching, a canonical design for observational studies, takes many forms that rest on distinct — yet often implicit — statistical principles. We construct a matching pipeline for practitioners that makes these principles explicit and integrates stages often treated separately within a coherent design-based framework. The pipeline begins from the conceptual ideal of a randomized experiment, traces how observational studies depart from it, and then employs matching to approximate that ideal. The next stage is inference under the as-if randomization assumption that matched sets are equivalent to a collection of randomized experiments within blocks, where each block has a fixed number of treated units equal to the number observed in that set. Under this assumption, we consider inference on all individual effects in the "sharp" framework and on the average effect in the "weak" framework. The final stage is a sensitivity analysis to assess, under either framework, how inferences change under departures from as-if randomization. Each step includes extensively commented `R` code that equips practitioners to implement both established and newly developed procedures, including several not yet available in existing `R` packages. We illustrate the full workflow through an application examining the effect of United Nations peacekeeping interventions on the duration of post-conflict peace.

1 Design-Based Foundations of the Matching Pipeline

1.1 The Randomized Experimental Ideal

Imagine a randomized experiment in which a researcher flips a fair coin independently for each individual in the study. Heads means assignment of the individual to control, while tails means assignment to treatment. After assignments to treatment and control, the researcher administers the conditions and then compares outcomes between treated and control groups.

Why is this procedure effective? Randomization is a *fair lottery*, in that every individual has the same chance of being assigned to treatment. This equality of chances means that individuals who would respond more strongly to treatment are no more likely to receive it than those who would respond less strongly, and the same is true for control. Because each individual's assignment is determined by a toss of the same coin (with the same probability of landing heads or tails), randomization leaves only two possibilities: (1) the difference in outcomes between treatment and control groups reflects the true causal effect, or (2) chance variation produced a misleading difference. Although misleading differences can occur by chance, randomization is valuable because it enables us to use statistical tools to quantify and limit the chance of such errors. As a result, randomized experiments yield especially credible causal conclusions.

Randomization is useful not only as a procedure, but also as an idea. It helps us understand when statistical tools will (and will not) yield credible conclusions, even when we have not directly randomized. In an observational study, the researcher does not control who receives the treatment and instead observes units after they have been assigned to treatment and control groups. In such settings, the idea of randomization can guide how we design studies so that they yield more credible causal conclusions.

1.2 Bridging Randomized Experiments and Observational Studies

A useful framework for connecting randomized experiments to observational studies is what Rubin (1977) calls "assignment to treatment group on the basis of a covariate," where a covariate is a pre-treatment characteristic of a study's units. To make this idea concrete, suppose treatment is assigned by independent coin tosses for each individual in which the probability of heads or tails depends on that individual's value of a single covariate. Consequently, all individuals with the same covariate value share the same

chance of receiving treatment, while those with a different covariate value may have a different chance.

In this setting, we can envision forming groups (i.e., matched sets) so that all individuals within a group share the same covariate value. Each group then functions as a miniature randomized experiment in that random chance alone explains why some individuals in the group received treatment while others did not. Importantly, to justify this interpretation, we do not need to know each individual's actual probability of treatment. It is enough to know that the probability of treatment depends only on the covariate, which ensures that all individuals with the same covariate value have the same chance of treatment.

1.3 Matching to Approximate the Randomized Experimental Ideal

The same intuition applies to matching when the probability of treatment depends on many baseline covariates. The underlying idea is that individuals with similar covariates have similar treatment assignment probabilities. Thus, in an effort to recreate a randomized experiment, we use the covariates we observe and believe determine treatment chances in order to divide individuals into groups, with each group containing both treated and control subjects who are homogeneous in those covariates. We are, in effect, constructing a new single variable: group (i.e., matched set) membership. We can think of this membership variable as the single covariate in the framework of "assignment to treatment group on the basis of a covariate" (Rubin, 1977) discussed in Section 1.2. Although we still do not know each individual's treatment probability, the hope is that all individuals within a group share the same probability, whatever it may be.

We call the condition in which all individuals within each matched set share the same treatment probability, strictly between 0 and 1, *as-if randomization*. As-if randomization is closely related to the conditions known as *selection on observables* and *strong ignorability*, although those conditions are typically stated directly in terms of covariates rather than in terms of matched sets built from those covariates. The term as-if randomization is especially fitting because, when units are assigned independently and all units within each matched set share the same individual treatment probability, all assignments within each set — holding fixed the set's observed number of treated units — are equally likely (treating the matched sets themselves as fixed; see Section 3.1). In this case, the probability distribution over these assignments is equivalent to that induced by a randomized experiment in which a researcher first forms blocks corresponding to the matched sets and then randomly assigns a fixed number of units to treatment

within each block. Thus, when as-if randomization holds, we can apply the same statistical tools that we would use under such a completely randomized experiment within blocks to draw credible causal inferences from the observational study.

1.4 Why Sensitivity Analysis Matters

Unlike randomized experiments, even the best matched designs rely on the strong assumption of as-if randomization. When this assumption holds, we can draw causal conclusions by analyzing the data as if they came from a randomized experiment. However, if the assumption is wrong, our causal claims are no longer guaranteed to be credible.

How can this assumption fail? First, individuals within a group may be similar on observed covariates, but not similar enough to have the same treatment probabilities. Second, treatment assignment may depend on covariates we did not measure. If so, even if individuals within groups appear comparable on observed covariates, those individuals may still differ on hidden covariates that determine the probability of treatment.

For these reasons, it is important to assess the sensitivity of our causal conclusions to departures from as-if randomization. Conclusions are especially convincing when they hold not only under this assumption, but also under moderate violations of it. Conclusions that collapse under only mild departures are much less convincing.

1.5 Roadmap of the Design-Based Matching Pipeline

Building on these design-based foundations, we now outline a pipeline that starts with matching and then proceeds to inference and sensitivity analysis:

- **Construct and evaluate matched sets**. Given the specified covariates to be balanced, we begin with the mechanics of optimal matching (Hansen, 2004; Hansen and Klopfer, 2006): choosing a distance measure that defines similarity on the covariates, setting calipers — maximum allowable distances between treated and control units for inclusion in the same matched set — and imposing structural constraints (e.g., requiring matches to be pairs). We then show how to evaluate the resulting design in terms of both covariate balance and effective sample size. For the former, we

focus on tests proposed by Hansen and Bowers (2008) that compare the matched design's covariate balance to what one would expect under an equivalent completely randomized experiment within blocks (that is, random assignment with a fixed number of treated units per block).

- **Draw causal inferences**. Once a matched design is chosen, practitioners can conduct inference under as-if randomization, either within a "sharp" causal framework, which pertains to individual-level effects for all units, or a "weak" framework, which pertains to a summary quantity of the unit-level causal effects, typically the average treatment effect (ATE). For the sharp framework, we focus on how researchers can use both simulation- and Normal-based approximations to the exact randomization distribution to perform hypothesis tests, which can be inverted to obtain confidence sets and point estimates. For the weak framework, we cover estimation of the ATE and hypothesis tests about it. We emphasize exposition and code for recent variance estimators tailored to designs with only 1 treated or 1 control unit per matched set (Fogarty, 2018; Pashley and Miratrix, 2021) — an important case, since the closest matches that discard the fewest units can always be achieved with such designs (Gu and Rosenbaum, 1993; Rosenbaum, 1991; Hansen, 2004, 2011).
- **Assess sensitivity**. Finally, we turn to sensitivity analyses under both inferential frameworks. We review established methods for conducting sensitivity analysis for tests of sharp nulls under possible violations of as-if randomization (Rosenbaum and Krieger, 1990; Gastwirth et al., 2000; Rosenbaum, 2018). We then describe and implement new methods that extend sensitivity analysis to tests of weak null hypotheses (Fogarty, 2023).

Figure 1 presents a flow diagram summarizing the overall pipeline, partitioned into three stages that correspond to Sections 2.2–2.4. The diagram displays each step and decision point, the relevant `R` tools (whether existing packages or custom functions included herein), and core references. All relevant code and replication materials are publicly available in the manuscript's companion GitHub repository, https://github.com/tl2624/matching-guide.




Figure 1: Flow diagram of the design-based matching pipeline.

2 Implementation of the Matching Pipeline

We provide a high-level overview of the ideas behind matching and include code that demonstrates how to implement those ideas with various `R` packages. We also work through some of these steps "by hand" to underscore the underlying conceptual issues. Doing so also gives practitioners more flexibility to adapt the matching pipeline to their own needs.

We break the implementation into specific decision points that practitioners commonly face, and present the pipeline in three main stages (see Figure 1):

First Stage: Making a Matched Dataset (toward Justifying As-If Randomization)

(a) How Do I Measure Similarity on My Chosen Covariates?

(b) How Can I Apply Rules for Matches to Ensure Comparability?

(c) How Can I Apply Rules for Matches to Improve Effective Sample Size?

(d) How Do I Actually Form the Matches?

(e) How Do I Decide Whether to Move Ahead with My Matched Design?

Second Stage: Causal Inference (under As-If Randomization)

(a) How Do I Draw Inferences under the Sharp Framework?

(b) How Do I Draw Inferences under the Weak Framework?

Third Stage: Sensitivity Analysis (Departures from As-If Randomization)

(a) How Do Inferences under the Sharp Framework Change under these Departures?

(b) How Do Inferences under the Weak Framework Change under these Departures?

Before turning to First Stage, we introduce a running example taken from Gilligan and Sergenti (2008), which we use throughout this manuscript.

2.1 Running Example: United Nations Peacekeeping and Post-Conflict Peace

We introduce matching through an example that examines the causal effect of United Nations (UN) peacekeeping missions on the durability of post-conflict peace, a question of central importance for both

academic research and policy. Our example draws on data from Gilligan and Sergenti (2008), whose title includes the phrase "Matching to Improve Causal Inference," underscoring the value of applying matching to study the UN's causal impact on post-conflict peace. These data are publicly available in the supplementary information of the article's webpage in the *Quarterly Journal of Political Science* (DOI: 10.1561/100.00007051).

The dataset we use from Gilligan and Sergenti (2008) includes 87 observations, each corresponding to a country's peace period episode following a civil war, with episodes beginning as early as January 1989 and data extending through December 2003. In some episodes, UN peacekeepers intervened (e.g., Sierra Leone, Jan 2001 – Dec 2003), while in others they did not (e.g., Macedonia, Sep 2001 – Dec 2003). The treatment variable is UN intervention (`UN`), coded as 1 if a UN mission was present during the peace period and 0 otherwise. The outcome variable, `ldur`, is the log of the duration of the peace spell in months, which Gilligan and Sergenti (2008) measure from the start of peace until the outbreak of a new conflict. Duration is thus measured from the onset of peace rather than from the start of a UN intervention, but the two nearly coincide: "Of the 19 post-conflict UN interventions, the United Nations was present within the first month for 16 of them" (Gilligan and Sergenti, 2008, p. 118). For 43 of the 87 episodes, peace had not ended by December 2003, so duration is right-censored at that date. Our analyses therefore concern the log of the observed, possibly censored, duration rather than the log of the full duration of peace.

As Gilligan and Sergenti (2008) state, "UN missions are not randomly assigned" (p. 89). Whether peacekeepers are present during a given post-conflict peace period presumably depends on baseline covariates measured by Gilligan and Sergenti (2008), including the logged number of battle deaths (`lwdeaths`), the duration of the previous war in months (`lwdurat`, which, despite its name, is not logged), ethnic fractionalization (`ethfrac`, a 0–100 index for which dividing by 100 is intended to represent the probability that two randomly chosen individuals belong to different ethnic groups), logged population size (`pop`), and others. We implement matching using these same covariates, but emphasize that our exercise is expository and not intended as a replication of the original findings.

2.1.1 Loading the Data for the Running Example

To load the data, you could first download the replication files from the supplementary information on the article's webpage (DOI: 10.1561/100.00007051), save the files to your working directory, and then use a package such as `haven` to load the Stata (.dta) file, `peace_pre_match.dta`, into `R`. However, for our purposes, we recommend loading our pre-created .rds file (`peace_pre_match.rds`), in which Stata's monthly numeric dates have been converted to `R`'s `Date` class and the geographic region indicators recoded into a single factor variable (`region`). The following command loads this pre-created dataset (`peace_pre_match.rds`) into the `R` environment as an object named `data`.

```
# Define base URL for the Matching Guide GitHub repository

base_url <- "https://raw.githubusercontent.com/tl2624/matching-guide/main"

# Load the cleaned dataset
data <- readRDS(
  url(
    paste0(
      base_url, "/data/peace_pre_match.rds"
    )
  )
)
```

2.2 First Stage: Making a Matched Dataset (toward Justifying As-If Randomization)

In an observational setting, treatment is not assigned by the flip of a coin but depends on individuals' covariates. The goal of matching is to compare treated and control units that have the same chances of treatment based on those covariates — i.e., the same *propensity scores*. If we could observe propensity scores, it would be straightforward to compare treated and control units by matching them directly on their propensity scores. Because propensity scores are not directly observed, we instead aim to create a collection of matched sets that is *balanced* — meaning that treated and control observations are similar in their covariates.

Before turning to questions of covariate similarity and matching, it is important to note that substantive transformations of covariates play a central role, as they determine the inputs on which subsequent

notions of similarity between units are based. These transformations reflect substantive judgments and are often among the most important decisions in practice. In this manuscript, however, we do not address these substantive choices. Instead, we take the transformations used by Gilligan and Sergenti (2008) as given. For example, they measure ethnic fractionalization (`ethfrac`) on a 0–100 scale (rather than a 0–1 scale) and apply logarithmic transformations to covariates such as the number of deaths (`lwdeaths`) and population size (`pop`), among others. Our focus is on design and analysis decisions conditional on these substantively chosen scales.

2.2.1 How Do I Measure Similarity on My Chosen Covariates?

In the simplest terms, matching is about ensuring apples-to-apples, rather than apples-to-oranges, comparisons between treated and control observations (see Rubin and Waterman, 2006). To create matched sets in which treated and control groups are similar in their covariates, we first need a distance measure that quantifies how close any two observations are. With such a measure in hand, we can then construct sets of treated and control observations that are close on this measure — i.e., apples-to-apples in their pre-treatment characteristics.

We record the distances between treated and control units in a distance matrix, in which the rows correspond to treated units and the columns to control units. Each cell of the matrix records the distance between a specific treated unit and a specific control unit, as defined by a distance measure. This distance measure takes the baseline covariates of the two units and maps them to a single nonnegative number, with smaller values indicating greater similarity.

In our setting, we are interested in similarity across 9 covariates, named in the object `covs`.

```
# Define character vector of the 9 covariate names in the dataset
covs <- c("lwdeaths", "lwdurat", "ethfrac", "pop", "lmtnest", "milper",
          "bwgdp", "bwplty2", "region")
```

The first four — `lwdeaths`, `lwdurat`, `ethfrac`, and `pop` — were introduced earlier. The others are a logged measure of the percentage of a country's terrain that is mountainous (`lmtnest`), the logged number of military personnel in a country, in thousands (`milper`), logged GDP per capita before the last civil war (`bwgdp`), the Polity score (a −10 to 10 scale, from autocratic to democratic institutions) before the last civil war (`bwplty2`), and a region factor (`region`) with categories for Eastern Europe,

Latin America, Asia, Sub-Saharan Africa, and North Africa/Middle East.

All of these covariates are measured before treatment and presumably determine the chance of a UN intervention during a country's peace period. Our interest in them stems primarily from their role in determining those intervention probabilities. Yet many of these covariates may also be prognostic; that is, they help predict the outcome of interest in Gilligan and Sergenti (2008), the log duration of the peace period (`ldur`) that countries would potentially experience with or without a UN intervention. This prognostic value of covariates can provide an additional reason to match on them (Hansen, 2008a) or to adjust for them after matching (Sales et al., 2018).

There are many ways to measure the distance between a treated and a control unit. For example, we might compare units using the Euclidean distance — i.e., the square root of the sum of squared differences — across all baseline covariates. To do so, we first convert the factor variable `region`, which stores categorical labels, into an indicator variable for each region, since distance measures require numeric variables rather than categorical labels. Following the convention of `match_on()`, introduced below, each indicator equals $1/\sqrt{2}$ for units in that region and 0 otherwise, so that belonging to different regions adds exactly 1 to the squared distance.

```
# Install "dplyr" package (only run if not already installed)
# install.packages("dplyr")

# Load package for data manipulation (mutate, group_by, summarize, etc.)
library(dplyr)

# Convert "region" into indicators equal to 1/sqrt(2) or 0
data <- data |> # Pipe (|>) to pass left-hand result into next function call
  mutate(
    # Use mutate() to create new variables by transforming existing columns
    # ifelse() returns one value if condition is TRUE and another if FALSE
    eeurop   = ifelse(test = region == "eeurop",   yes = 1/sqrt(2), no = 0),
    lamerica = ifelse(test = region == "lamerica", yes = 1/sqrt(2), no = 0),
    asia     = ifelse(test = region == "asia",     yes = 1/sqrt(2), no = 0),
    ssafrica = ifelse(test = region == "ssafrica", yes = 1/sqrt(2), no = 0),
    nafrme   = ifelse(test = region == "nafrme",   yes = 1/sqrt(2), no = 0)
  )

# Covariate names with "region" replaced by indicator variables
expanded_covs <- c(
  # setdiff() returns elements in 'x' that are not in 'y'
  setdiff(
```

```
    x = covs,
    y = "region" # Drops "region" from the covariate list
  ),
  "eeurop", "lamerica", "asia", "ssafrica", "nafrme"
)
```

As an example, we calculate the Euclidean distance between post-conflict Liberia, where the UN did intervene, and post-conflict Guinea-Bissau, where the UN did not.

```
# Extract covariates for Liberia and Guinea-Bissau (cname = country name)
liberia <- data[data$cname == "Liberia", expanded_covs]

guinea_bissau <- data[data$cname == "Guinea-Bissau", expanded_covs]

# Compute Euclidean distance between the two countries on these covariates
sqrt(
  sum(
    (liberia - guinea_bissau)^2
    )
  )  # Display the Euclidean distance
```

```
[1] 57.44263
```

We can obtain the full matrix of pairwise distance values using `match_on()` from the `optmatch` package. We do not need to manually recode factor variables into indicator variables because `match_on()` handles this conversion automatically. (Like our hand calculation, `match_on()` scales each region indicator by $1/\sqrt{2}$, so the resulting distances equal the manual calculation.)

```
# Create a formula: UN (treatment indicator) ~ covariates
# Note: we keep "region" in covs as a factor
cov_fmla <- reformulate(
  termlabels = covs,
  response = "UN"
)

# Install optmatch if not already installed
# install.packages("optmatch")

# Load optmatch, which provides the match_on() function
library(optmatch)

# Euclidean distance matrix: treated (UN = 1) vs control (UN = 0)
dist_mat_euc <- match_on(
  x = cov_fmla,                 # Formula for covariates
```

```r
  data = data,                    # Dataset used
  method = "euclidean"            # Use Euclidean distance
)

# Add country names (cname) as row/column labels for clarity
dimnames(dist_mat_euc) <- list(
  data$cname[data$UN == 1], data$cname[data$UN == 0]
)

# Display submatrix of distances:
round(        # Round distances for display
  # treated units in rows 11-15 vs. control units in columns 27-30
  x = dist_mat_euc[11:15, 27:30],
  digits = 2  # Number of decimal places to round
)
```

```
              Niger Guinea   Togo Central African Republic
Sierra Leone  94.21 101.18 116.32                    116.94
Zaire         26.68  29.87  45.63                     46.57
Rwanda        66.75  71.57  77.04                     76.03
Mozambique   133.47 140.60 155.40                    155.80
Namibia      246.44 253.23 268.35                    268.08
```

In this subset of the full distance matrix, the first entry is the Euclidean distance of Sierra Leone (treated) from Niger (control). The last listed entry is the distance between Namibia (treated) and the Central African Republic (control).

One concern with Euclidean distance is that it depends on the scale of the covariates. For example, Guatemala (treated) belongs to Latin America and Afghanistan (control) to Asia, a substantively large difference; yet belonging to different regions adds only 1 to the squared Euclidean distance between the two countries, because they differ by $1/\sqrt{2}$ on each of two region indicators (`lamerica` and `asia`). The two countries' ethnic fractionalization scores (`ethfrac`), 64.37 and 65.83 on the 0–100 scale, differ by only 1.46 points, a substantively small difference; yet this difference adds 2.13, more than twice as much as belonging to different regions. Intuitively, we would not want such a small difference in ethnic fractionalization to count for more than belonging to different regions of the world. More generally, we want differences across variables to be placed on a comparable scale, so that a meaningful difference in one variable counts about the same as a difference of similar importance in another.

Another concern with Euclidean distance is that it ignores correlations among covariates. For example, countries with larger populations (`pop`) usually have more military personnel (`milper`), if only because a

larger population provides a greater pool of potential recruits. Therefore, differences in both covariates (`pop` and `milper`) may largely reflect the same underlying factor — population size (`pop`). Yet Euclidean distance adds these differences separately, as if they were unrelated, which can exaggerate the overall distance between two observations.

The Mahalanobis distance (Mahalanobis, 1936) addresses both of these concerns. First, it standardizes covariates so that differences are placed on a comparable scale. Second, the Mahalanobis distance adjusts for correlations among covariates, ensuring that highly related variables are not effectively counted twice. We can construct a distance matrix based on Mahalanobis rather than Euclidean distances as follows.

```
# Mahalanobis distance matrix: treated (UN = 1) vs control (UN = 0)
dist_mat_mah <- match_on(
  x = cov_fmla,
  data = data,
  method = "mahalanobis" # Use Mahalanobis distance
)
```

The standardization in the Mahalanobis distance is distinct from researchers' substantive choices about how to transform covariates. Substantive transformations — such as logging certain covariates — reflect subject-matter considerations and precede the matching procedure. The standardization in the Mahalanobis distance is a statistical device that operates on the pre-specified covariates to compute distances comparably across covariates. Ultimately, the purpose of this device is to facilitate matches that are similar on the original covariate scales.

2.2.1.1 Similarity on the Estimated Propensity Score

So far, we have focused on ways to measure distance between treated and control units across many covariates in order to identify which units are most similar and group them together to achieve covariate balance. However, when there are many covariates, it becomes difficult to find treated and control units that are similar on all covariates. This challenge is often referred to as the "curse of dimensionality."

A common way to address this problem is to reduce the information from many covariates into a lower-dimensional form. The *estimated propensity score* does this by collapsing information from all covariates into a single number. This number represents a transformation of an estimated linear index of covariates that accounts for how strongly each covariate predicts treatment.

Consider, for example, a logistic model for the estimated propensity score of an individual unit i. We write this model as

$$\hat{\lambda}(\boldsymbol{x}_i) = \frac{1}{1 + \exp(-\hat{\boldsymbol{\beta}}^\top \boldsymbol{x}_i)}, \tag{1}$$

where $\boldsymbol{x}_i$ is the covariate vector, $\hat{\boldsymbol{\beta}}$ is the vector of estimated coefficients, and the superscript $^\top$ denotes the transpose, turning the column vector $\hat{\boldsymbol{\beta}}$ into a row vector, so that $\hat{\boldsymbol{\beta}}^\top \boldsymbol{x}_i$ is a scalar. The quantity $\hat{\boldsymbol{\beta}}^\top \boldsymbol{x}_i$ is the estimated linear index, and the logistic (inverse-logit) function, $1/(1 + \exp(-a))$, maps any real number, a, onto the interval (0, 1). The estimated linear covariate index for unit i, $\hat{\boldsymbol{\beta}}^\top \boldsymbol{x}_i$, is simply the logit, i.e., log odds, transformation of $\hat{\lambda}(\boldsymbol{x}_i)$ in (1).

This quantity, $\hat{\lambda}(\boldsymbol{x}_i)$, is a simple transformation of a linear index of covariates that best "separates" treated from control units. The estimated coefficients ($\hat{\boldsymbol{\beta}}$) "separate" treated from control units on the linear index because the coefficients are chosen to maximize the likelihood of the logistic model in (1), and this likelihood is larger when the fitted probabilities are close to 1 for treated units and close to 0 for controls. When treated and control units have little or no covariate overlap, the estimated linear indices can diverge substantially, very positive for treated units and very negative for controls. With greater overlap, the estimated linear indices for treated and control units are more similar, clustering more closely around a common value.

This estimation process reflects how predictive each covariate is of treatment, holding the other covariates fixed. A covariate that strongly predicts treatment, given the other covariates, receives a coefficient that turns typical differences in that covariate into large differences in the estimated linear index. A covariate with little predictive power receives a coefficient that turns typical differences in that covariate into small differences in the estimated linear index. Consequently, when we assess similarity on the estimated linear index of covariates in (1), the estimated coefficients assign greater importance to covariates that strongly predict treatment and less importance to those that do not.

To see this logic in action, we first estimate a logistic propensity score model using all covariates except `region`, which we exclude because all episodes in one region (Asia) are controls, a case of quasi-complete separation in logistic regression (Albert and Anderson, 1984).

```r
# Formula for UN ~ covariates (excluding "region")
psm_cov_fmla <- reformulate(
  termlabels = setdiff(
    x = covs,
    y = "region"
  ),
  response = "UN"
)

# Fit logistic regression for propensity score model
psm <- glm(
  formula = psm_cov_fmla,             # Treatment ~ covariates
  family = binomial(link = "logit"), # Logistic regression (logit link)
  data = data
)
```

Using this fitted propensity score model, we can extract each unit's estimated linear covariate index, $\hat{\boldsymbol{\beta}}^\top \boldsymbol{x}_i$, as follows.

```r
# Extract logit propensity scores (linear predictors from fitted model)
lin_cov_inds <- psm$linear.predictors  # = model.matrix(psm) %*% coef(psm)
```

The following code shows that the estimated linear covariate indices from the model `psm` correspond exactly to the logits (i.e., the log-odds transformations) of the model's predicted probabilities of treatment, which range between 0 and 1.

```r
# Extract estimated propensity scores (predicted probabilities of UN = 1)
p_scores <- psm$fitted.values

# Check that lin_cov_inds equals log odds (propensity scores on logit scale)
all.equal(
  lin_cov_inds, log(p_scores / (1 - p_scores))
)

# Also check that p_scores equals logistic(lin_cov_inds)
all.equal(
  p_scores, 1 / (1 + exp(-lin_cov_inds))
)
```

To see how two observations that differ greatly on a covariate can still have similar estimated propensity scores, consider post-conflict Haiti (treated; peace beginning in 1992) and Papua New Guinea (control; peace beginning in 1991). The two countries differ greatly in ethnic fractionalization (`ethfrac`), about 1

versus 45 on the 0–100 scale, but have the same logged number of battle deaths (`lwdeaths`) and similar logged numbers of military personnel (`milper`). Despite this large difference in ethnic fractionalization, Haiti's estimated linear index exceeds Papua New Guinea's by only about 0.26. The gap in `ethfrac`, 43 points or about 1.7 standard deviations of `ethfrac` across all 87 episodes, contributes only about 0.13 to this difference because the estimated coefficient on `ethfrac` is close to 0 (about -0.003 per point). By contrast, the gap in `milper`, 0.29 or about 0.18 standard deviations of `milper` across all 87 episodes, contributes about 0.26 because the estimated coefficient on `milper` is large (about -0.92). The remaining covariates contribute positive and negative amounts that net to about -0.13. The estimated linear index therefore gives little weight to the covariate on which the two countries differ most and more weight to covariates that strongly predict treatment.

To illustrate these broader patterns, Figure 2 compares the distributions of the estimated linear covariate index for treated and control groups. There is some, but limited, overlap between treated and control groups. In accordance with our earlier discussion of how the estimated linear covariate index "separates" treated from control units, many control observations have very negative values (as low as -7.52), far from the treated units' range of -2.24 to 2.21. Nevertheless, a sufficient number of treated and control observations have estimated linear covariate indices that cluster around 0, indicating covariate overlap for at least a subset of treated and control observations.




Figure 2: The empirical distributions of the estimated linear covariate index for treated and control groups.

2.2.2 How Can I Apply Rules for Matches to Ensure Comparability?

The discussion in Section 2.2.1 on measuring covariate distances between treated and control observations helps identify which treated–control pairs are similar. The goal of identifying these similar treated–control pairs is to form matched sets that are closely aligned on their covariates, thereby improving balance between treatment and control groups. In practice, we do this by excluding potential matches that are "too dissimilar." There are two common ways to do this:

- **Exact matching**: Require that units be identical on some subset of important covariates, typically those that are categorical or coarse enough for units to take the same values.
- **Calipers**: Impose a threshold for the maximum allowable distance so that no matched set may include a treated and a control unit that are farther apart than this caliper.

As a simple example to build intuition, we will impose the following constraints:

- Observations can belong to the same matched set only if they are in the same geographic region.
- Treated and control observations more than two points apart on the Polity score cannot be in the same matched set.

We now impose the first constraint, requiring an exact match on region. Doing so produces a sparse distance structure, `em_region`, that stores a distance of 0 for every treated–control pair in the same region and stores no entry for pairs in different regions. The `optmatch` package treats these missing entries as forbidden (infinite) distances, so treated and control units can be matched only within a region.

```
# Create distance structure: 0 if units are in same region, Inf otherwise
em_region <- exactMatch(
  x = UN ~ region,
  data = data
)
```

Now we impose the second constraint by constructing a distance matrix based on the Polity score, `bwplty2`, with a caliper of 2. This matrix records the absolute difference in Polity scores when the difference is 2 or less, and assigns a value of ∞ (denoted in `R` as `Inf`) when the difference exceeds 2. The ∞ entries are crucial because `optmatch` minimizes the sum, across all matched sets, of the distances between each treated–control pair within the set. Consequently, any treated–control pair differing by more than 2 points on the Polity score is assigned an overall distance of ∞, and this infinite distance prevents the two units from being matched.

```
# Euclidean distance on Polity score (bwplty2) with caliper = 2
# Pairs differing by > 2 are set to Inf
euc_dist_polity_cal_2 <- match_on(
  x = UN ~ bwplty2,
  caliper = 2,  # Set caliper
  data = data,
  method = "euclidean"
)
```

Note that we used Euclidean distance here because, after exactly matching on geographic region, matching proceeds on only 1 covariate (Polity score), so we do not have to worry about covariates' relative scales.

Finally, we combine the two constraints into distance matrices defined within each region by "adding" the objects `em_region` and `euc_dist_polity_cal_2`, as the following code shows.

```
# Overall distance matrix by element-wise addition of two distance matrices
overall_dist_mat <- em_region + euc_dist_polity_cal_2
```

2.2.3 How Can I Apply Rules for Matches to Improve Effective Sample Size?

Beyond comparability in covariates, we also care about the matched study's size. The effective sample size depends not only on the total number of units included in our matches but also on how those units are arranged across the matched sets.

The `optmatch` package will always produce matches with a particular arrangement of units across sets. In particular, all sets contain either 1 treated unit or 1 control unit. Rosenbaum (1991) shows that any grouping of treated and control units into matched sets can be subdivided into sets of this form without increasing the quantity that `optmatch` minimizes, namely the sum of within-set treated–control distances, and without discarding any additional units (see also Gu and Rosenbaum, 1993; Hansen, 2004). In practice, optimal full matching can yield lopsided sets, with either 1 treated matched to many controls or 1 control matched to many treated units, which has implications for the effective sample size.

In the `optmatch` package, the effective sample size is defined as the sum, across matched sets, of the harmonic mean of the numbers of treated and control units within each set. The formal definition is given by

$$\text{Effective sample size} = \sum_{s=1}^{S} \left[\left(m_s^{-1} + (n_s - m_s)^{-1} \right) / 2 \right]^{-1}, \tag{2}$$

where the index $s \in \{1, \ldots, S\}$ runs over the S matched sets, with m_s denoting the number of treated units in set s and $n_s - m_s$ the number of control units. With n_s denoting the number of units in set s, the total number of individuals included in the matched study is $n = \sum_{s=1}^{S} n_s$.

From the formula in (2), we can see how the effective sample size depends on the arrangement of units across sets. For example, in a study with 4 total units, the effective sample size would be 2 if the units were arranged into 2 matched pairs. By contrast, in a study of the same total size but arranged as a single set with 1 treated unit and 3 controls, the effective sample size would be 1.5. The effective sample size is larger in the former arrangement because it provides two distinct treated-versus-control comparisons, whereas the latter provides only one.

This definition of effective sample size anticipates subsequent outcome analysis under as-if randomization, as effective sample size bears directly on the precision of estimators and the power of hypothesis tests.

Assuming constant, additive treatment effects within a set, the variance of the Difference-in-Means — the average outcome among treated units minus the average outcome among control units — in that set is proportional to the set's outcome variance divided by the set's harmonic mean (Hansen and Bowers, 2008; Hansen, 2011). For a fixed set size, the harmonic mean is largest when the numbers of treated and control units are as equal as possible. Thus, all else equal, matched pairs and other sets with equal numbers of treated and control units provide more information about causal effects than lopsided sets with unequal treated-to-control ratios. If outcome variances are also equal across sets, then among weighted averages of the sets' Differences-in-Means, weighting by the sets' harmonic means gives the smallest variance (Hansen, 2011, p. 164, fn. 3), and the effective sample size in (2) then measures the precision of that weighted average.

One straightforward way to increase effective sample size is to relax restrictions on which units can be matched. Relaxing calipers can admit additional units into the matched design by allowing matches for units that would otherwise be discarded. For instance, we might widen the caliper on Polity score from 2 to 3 and then rebuild the distance matrix.

```
# Apply a caliper of width 3 to the Polity Euclidean distance matrix
euc_dist_polity_cal_3 <- match_on(
  x = UN ~ bwplty2,
  caliper = 3,
  data = data,
  method = "euclidean"
)

# Combine regional exact match distance with Polity distance
em_region + euc_dist_polity_cal_3
```

In this case, doing so increases the effective sample size, but such gains need not be substantial. When newly admitted units are absorbed into existing matched sets — each of which contains only one treated or one control unit — rather than generating additional treated-versus-control comparisons, the resulting contribution to effective sample size is limited. More generally, as the formula in (2) makes clear, the effective sample size depends on how units are distributed across sets. When units, including newly admitted ones, are concentrated in only a few sets, the resulting gain in effective sample size is modest.

Instead of relying solely on caliper width, researchers can also shape the effective sample size by controlling the structure of matched sets via the `min.controls` and `max.controls` arguments in `optmatch`'s

`fullmatch()` function. These arguments specify lower and upper bounds, respectively, on the ratio of control to treated units within each matched set. By default, `min.controls = 0` and `max.controls = Inf`, which impose no restrictions on matched set composition. Changing these defaults allows researchers to control how lopsided matched sets may be, thereby shaping the effective sample size.

To illustrate, suppose we want to restrict matches using the `overall_dist_mat` introduced earlier. We know that `optmatch` will divide the data into matched sets containing one treated unit and any positive number of controls, or one control unit and any positive number of treated units. However, we can impose additional constraints on this full matching, such as requiring a minimum control-to-treated ratio of 1:2 — that is, at least one control for every two treated units (`min.controls = 0.5`) — and no more than two controls per treated unit (`max.controls = 2`). Under these restrictions, allowable matched sets could include 2 treated units with 1 control, 1 treated unit with 1 control (a matched pair), or 1 treated unit with 2 controls.

```
# Full matching using overall distance matrix
fullmatch(
  x = overall_dist_mat,
  min.controls = 0.5,    # At least 0.5 controls per treated unit
                         # (i.e., no more than 2 treated per control)
  max.controls = 2,      # At most 2 controls per treated unit
  omit.fraction = NULL, # Fraction of controls (if < 0, treated) to discard
  mean.controls = NULL, # Governs average controls per treated unit
  # At most one of omit.fraction or mean.controls can be non-NULL
  data = data
)
```

Note, in addition, that the `fullmatch()` arguments of `omit.fraction` and `mean.controls` provide explicit levers for controlling how many treated and control units are discarded, though at most one of `omit.fraction` or `mean.controls` may be set to a value other than `NULL`.

Imposing ratio constraints can have mixed consequences for balance and effective sample size. In some cases, balance may worsen if a control is forced to match with a less similar treated unit — though still within the specified calipers — in order to satisfy the minimum and maximum ratio rules. In other cases, such constraints may improve effective sample size by redistributing how units are grouped without hurting balance. However, if the ratio constraints are too stringent, they can reduce effective sample size by forcing too many units to be discarded.

In applied settings, final choices of calipers and ratio constraints typically follow iterative checks of both covariate balance and effective sample size. Practitioners often compare several specifications and select among those that satisfy diagnostics for both goals. Hansen and Sales (2015) outline how to conduct this process in a structured way, drawing on the stepwise intersection–union principle (SIUP) of hypothesis testing.

2.2.4 How Do I Actually Form the Matches?

The simple matching example in Sections 2.2.2 and 2.2.3 — based on an exact match on geographic region and a caliper on Euclidean distance for a single covariate (Polity score) — serves to illustrate the basic ideas. When matching on many covariates, however, we will often prefer some combination of Mahalanobis distance and propensity score matching, sometimes adding calipers on specific covariates. In what follows, we use *rank-based* Mahalanobis distance, which has the advantage of being less sensitive than the ordinary Mahalanobis distance to outliers and to covariates with many tied values, such as rare binary indicators (Rosenbaum, 2010, pp. 171–172). We further constrain the matching by imposing a caliper on the estimated propensity score, requiring treated and control observations to come from the same geographic region, and applying additional calipers directly to two covariates: ethnic fractionalization (`ethfrac`) and logged GDP per capita (`bwgdp`).

We impose a caliper on the logit of the estimated propensity score, that is, on the estimated linear covariate index in (1). The caliper is the largest difference on this logit scale that a treated and a control unit in the same matched set may have. Narrower calipers yield closer matches on the estimated propensity score but leave more units without a permissible match. We set the caliper to 0.5 times the population standard deviation of the logit across all 87 episodes. Because this standard deviation is approximately 1.96, treated and control units in the same set may differ by up to about 0.98 on the logit scale. This caliper is wider than the commonly recommended 0.20 standard deviations, a guideline that traces to Cochran and Rubin (1973, Table 2.3.1, p. 421).

```
# Add logistic regression linear predictors to dataset
data$logit_p_score <- lin_cov_inds

# Population standard deviation of logit_p_score (divides by n, not n - 1)
pop_sd_logit <- sqrt(
  mean(
```

```
    (data$logit_p_score - mean(data$logit_p_score))^2
  )
)

# Distance matrix for logit of estimated propensity scores
ps_mat <- match_on(
  x = UN ~ logit_p_score,
  caliper = 0.5 * pop_sd_logit,
  data = data,
  method = "euclidean"
)
```

We next construct the distance matrix for rank-based Mahalanobis distance.

```
# Rank-based Mahalanobis distance on covariates
# covs defined earlier as covariate names; here, drop "region"
rank_mah_mat <- match_on(
  x      = reformulate(
    termlabels = setdiff(
      x = covs,
      y = "region"),
    response   = "UN"),
  data   = data,
  method = "rank_mahalanobis" # Use rank-based Mahalanobis distance
)
```

Finally, we construct Euclidean distance matrices for ethnic fractionalization (`ethfrac`) and logged GDP per capita (`bwgdp`) using calipers of 35 and 2, respectively. The exact-match constraint on region has already been defined through the object `em_region` in Section 2.2.2.

```
# Compute Euclidean distance matrix for ethnic fractionalization
eth_mat <- match_on(
  x = UN ~ ethfrac,
  caliper = 35,
  data = data,
  method = "euclidean"
)

# Compute Euclidean distance matrix for logged GDP per capita
bwgdp_mat <- match_on(
  x = UN ~ bwgdp,
  caliper = 2,
  data = data,
  method = "euclidean"
)
```

Next, we combine the `ps_mat`, `rank_mah_mat`, `eth_mat`, `bwgdp_mat`, and `em_region` objects to form the overall distance structure. We then pass the combined object to `optmatch`'s `fullmatch()` function, imposing a constraint that no more than 4 control units may be matched to any treated unit.

```
# Full matching:
# PS + rank Mahalanobis + Euclidean (ethfrac, bwgdp) + region exact
fm <- fullmatch(
# x specifies the distance structure:
# match_on() formula, distance matrix, or sum of match_on() distances
  x           = ps_mat + rank_mah_mat + eth_mat + bwgdp_mat + em_region,
  data        = data,
  max.controls = 4 # <= 4 controls per treated (min.controls = 0 by default)
)
```

To calculate the effective sample size of the matched observations, we use the following function from `optmatch`.

```
# Effective sample size of matched sets
effectiveSampleSize(
  fm
)
```

```
[1] 15.26667
```

Referring back to (2), this reported effective sample size of approximately 15.27 equals the sum across sets of the within-set harmonic mean of the number of treated and control subjects. This effective sample size reflects the matched structure in which no set contains more than 4 controls per treated observation.

```
# Summarize matched sets and report effective sample size
summary(
  fm
)
```

```
Structure of matched sets:
1:0 2:1 1:1 1:2 1:3 1:4 0:1
  6   1   4   4   2   1  45
Effective Sample Size:  15.3
(equivalent number of matched pairs).
```

The notation in the `summary()` output of `optmatch` indicates the ratio of treated to control observations within each matched set. For example, `1:0` indicates 1 treated unit and no controls (effectively an unmatched treated unit). Similarly, `1:2` indicates 1 treated unit and 2 controls, and `0:1` indicates no

treated units and 1 control (effectively an unmatched control). For each label, the output shows how many matched sets have that particular structure. Here, 6 of the 19 treated episodes are left unmatched, so the inferences that follow concern only the 13 matched UN episodes and their 23 matched controls.

If we examine the object returned by the matching call (`fm`), we see that `optmatch` labels each observation according to its matched set, assigning `NA` to those not included. Because we performed exact matching by region, `optmatch` labels each matched set using the name of the exact-match stratum followed by a set index within that stratum. For example, the label `lamerica.1` denotes the first matched set within the Latin America stratum.

To see which units were matched together, we can add the `fm` object to the data frame and then list the units in a chosen set. For example, to view the `ssafrica.3` set, we run the following.

```
# Add matched set ID to data for each unit
data$fm <- fm

# Look at one matched set ("ssafrica.3") for illustration
data |>
  filter(
    fm == "ssafrica.3" # Keep only units in set "ssafrica.3"
  ) |>
  # Display selected variables
  select(
    cname, UN, region, logit_p_score, ethfrac, bwgdp, bwplty2
  )
```

```
# A tibble: 4 x 7
  cname      UN region   logit_p_score ethfrac bwgdp bwplty2
  <chr>   <dbl> <fct>            <dbl>   <dbl> <dbl>   <dbl>
1 Burundi     0 ssafrica         0.508    3.55  5.35      -7
2 Rwanda      1 ssafrica         0.148   12.9   5.68      -7
3 Somalia     0 ssafrica         0.209    7.67  6.61      -7
4 Lesotho     0 ssafrica         0.558   22.2   6.28       0
```

We can see that the exact match on geographic region holds, since all 4 countries are located in Sub-Saharan Africa. The logit of the estimated propensity score is similar across units, though not identical. The treated country, Rwanda, is matched to three controls — Burundi, Somalia, and Lesotho — and in each case the distance falls within 0.5 standard deviations of the estimated logit propensity scores (approximately 0.98), the caliper we specified. Likewise, Rwanda's distances to each of the three control units fall within the calipers of 35 for ethnic fractionalization (`ethfrac`) and 2 for logged GDP per capita

(bwgdp). By contrast, distances between control countries can technically exceed these thresholds, since optmatch enforces calipers only between treated and control units, not among controls.

We can also see that some covariates used in the propensity score model and the rank-based Mahalanobis distance — such as Polity score (bwplty2) — still show modest imbalance within this matched set. This imbalance is unsurprising. We applied a caliper on the logit of the estimated propensity score and matched on the rank-based Mahalanobis distance including bwplty2, but we did not apply a caliper directly to bwplty2, though such a caliper could easily be added if desired.

Had we wanted pair matching instead, we could call pairmatch() or set min.controls = 1 and max.controls = 1 in fullmatch(). The two calls give the same match when every treated unit can be paired. Otherwise, the two calls can differ, because fullmatch() requires every treated unit with a permissible match to be paired, whereas pairmatch() uses the group sizes in each exact-match stratum (here, each region) to fix how many units may go unmatched. When calipers rule out so many pairs that more units would have to go unmatched than pairmatch() allows, pairmatch() issues a warning and leaves that stratum entirely unmatched, as fullmatch() does when it cannot pair every treated unit that has a permissible match.

2.2.5 How Do I Decide Whether to Move Ahead with My Matched Design?

Once we have constructed our matched sets, we want to evaluate the overall quality of the matched design. Covariate balance is an important aspect of this evaluation. To assess covariate balance, we compare the balance in our matched observational study with the balance we would expect to see in a completely randomized experiment within blocks (Hansen and Bowers, 2008).

We next calculate adjusted covariate means for the treated and control groups by averaging set-specific means across matched sets, with each set weighted by its number of treated units, as RItools does; the adjusted treated mean is therefore the mean among matched treated units. We also report standardized differences, defined as the treated-minus-control difference in these adjusted means divided by a pooled standard deviation of the covariate — the square root of the average of the treated-group and control-group variances — computed in the overall data (including both matched and unmatched units). The significance tests, by contrast, weight each set by its contribution to the effective sample size (Hansen and Bowers, 2008).

```r
# Install "RItools" package (only run if not already installed)
# install.packages("RItools")

# Load RItools package for balance diagnostics (balanceTest)
library(RItools)
# Covariate balance test
cov_bal <- balanceTest(
  # Formula: treatment ~ covariates
  # update(): keep the original formula (. ~ .)
  # and add stratification by matched set, + strata(fm)
  fmla = update(
    cov_fmla, . ~ . + strata(fm)
  ),
  data = data,
  p.adjust.method = "none" # Method of p-value adjustment (none here)
)
```

	Before matching			After matching		
Covariate	Control mean	Treated mean	Std. diff	Control mean	Treated mean	Std. diff
Log Cumulative Battle Deaths from Last War	6.65	8.98	0.84*	8.34	8.57	0.08
Duration of Last War	50.28	80.53	0.39	60.51	73.23	0.16
Ethnic Fractionalization	56.50	49.21	−0.28	51.75	57.50	0.22
Log Population Size	9.51	8.75	−0.64*	8.89	8.91	0.02
Log Mountainous	2.22	2.80	0.43	2.69	2.82	0.09
Log Military Personnel	3.87	3.25	−0.42	3.63	3.54	−0.06
Log GDP per Capita Before Last War	6.56	6.59	0.03	6.73	6.55	−0.17
Democracy (Polity Score) Before Last War	−0.84	−2.58	−0.32	−2.19	−2.54	−0.07
Asia	0.19	0.00	−0.68*	0.00	0.00	0.00
Eastern Europe	0.15	0.37	0.51*	0.46	0.46	0.00
Latin America	0.12	0.21	0.25	0.08	0.08	0.00
North Africa/Middle East	0.12	0.11	−0.04	0.08	0.08	0.00
Sub-Saharan Africa	0.43	0.32	−0.23	0.38	0.38	0.00

Table 1: Adjusted covariate means for treated and control groups and standardized treated–control differences, before and after matching. Stars denote two-sided p-values less than or equal to 0.05.

Table 1 reports the results before and after matching. The adjusted means offer a direct description of balance. The standardized differences place all covariates on a common scale, enabling at-a-glance comparisons of imbalance across covariates. Stars indicate covariates whose harmonic-mean-weighted Difference-in-Means, which weights each set's treated-minus-control difference by the set's contribution to the effective sample size, would be unusually extreme under a completely randomized experiment within blocks, with the matched sets as blocks after matching and with all 87 episodes as a single block before matching. The harmonic-mean-weighted Difference-in-Means is not the difference between the adjusted means in Table 1, which weight each set by its number of treated units, although the two coincide before matching. The stars are based on a Normal approximation to the randomization distribution of

the harmonic-mean-weighted Difference-in-Means, and, because no adjustments are made for multiple comparisons, these stars are conservative for the purpose of judging balance. That is, the chance that at least one covariate is starred under a completely randomized experiment within blocks exceeds 0.05, so the stars err toward flagging imbalance. Alternative adjustments for testing multiple covariates are available, such as the Holm procedure (Holm, 1979), which is the default option in `RItools`'s `balanceTest()` function.

One concern with balance tests is that high p-values may arise not from improved covariate balance but from the reduction in effective sample size that typically accompanies the matching process (Austin, 2008; Imai et al., 2008). As Hansen (2008b) notes, however, this possibility is less troubling than it first appears. The same increase in standard errors that produces high p-values for covariate balance tests will also be present in subsequent outcome analyses, meaning that those high p-values remain informative. They suggest that any bias that the remaining imbalances in observed covariates could introduce into those analyses is small relative to those standard errors. We are therefore, if anything, less likely to overstate our causal conclusions — for example, by rejecting a true hypothesis about causal effects — than if the balance tests had been significant.

Regardless of whether the balance tests are statistically significant, there are also established guidelines for what constitutes sufficient balance. While precise thresholds depend on context and substantive judgment about each covariate's importance, two rules of thumb appear in Austin (2009) and Stuart (2010). Austin (2009) reports a proposal that standardized differences of 0.1 or greater indicate meaningful imbalance on a covariate, whereas Stuart (2010, p. 11) gives a more lenient threshold of 0.25. In our case, all covariates meet this latter standard, but three (duration of the last war, ethnic fractionalization, and GDP per capita, with standardized differences of 0.16, 0.22, and -0.17) exceed the 0.1 threshold reported by Austin (2009, p. 3090). None show statistically significant differences, indicating that observed imbalances would not be unusual under a completely randomized experiment within blocks (i.e., under as-if randomization).

In addition to assessing balance on each covariate individually, Hansen and Bowers (2008) also propose an omnibus, chi-squared (χ^2) test that evaluates balance across all covariates and their linear combinations simultaneously. The following code conducts this test.

```r
# Extract overall chi-square balance test by matched set (fm)
cov_bal$overall["fm", ]
```

```
   chisquare df   p.value
fm  2.631007  8 0.9553417
```

This χ^2 balance test yields an observed test statistic of 2.63 and a corresponding p-value of 0.96. This high p-value indicates that the observed level of covariate balance is consistent with what we would expect in a completely randomized experiment within blocks. The test has 8 rather than 13 degrees of freedom because the degrees of freedom equal the rank of the covariance matrix of the adjusted differences (Hansen and Bowers, 2008, p. 229). Because we matched exactly on region, the five region indicators are constant within every matched set, so their adjusted differences equal zero under every assignment that holds fixed the number of treated units in each matched set. These five differences therefore have zero variance and zero covariance with the other adjusted differences, and the covariance matrix has rank 8, the number of remaining covariates.

Despite the high p-value, there is no guarantee that balance is sufficient for as-if randomization. Residual imbalance on observed covariates and hidden imbalance on unobserved ones may undermine the as-if randomization assumption. For now, we proceed under the as-if randomization assumption, but we will later assess how sensitive our inferences are to violations of it due to such imbalances.

2.3 Second Stage: Causal Inference (under As-If Randomization)

In both the sharp and weak frameworks, the causal targets of inference are defined through potential outcomes. Under what is known as SUTVA, the stable unit treatment value assumption (Rubin, 1980, 1986; Cox, 1958), each unit has two potential outcomes: a value the outcome would take if that unit were assigned to treatment and a value the outcome would take if that unit were assigned to control. Let $y_{si}(1)$ and $y_{si}(0)$ denote these potential outcomes, respectively, for unit i in set s, where the index $i \in \{1, \ldots, n_s\}$ runs over the n_s units in set s. The individual treatment effect is $\tau_{si} = y_{si}(1) - y_{si}(0)$. With $n = \sum_{s=1}^{S} n_s$ total units, let $\boldsymbol{\tau} = (\tau_{11}, \tau_{12}, \ldots, \tau_{Sn_S})^\top$ collect all n unit-level effects, indexed first by matched set and then by unit within set, and write the ATE as $\tau = (1/n) \sum_{s=1}^{S} \sum_{i=1}^{n_s} \tau_{si}$.

Both the vector of unit-level effects, $\boldsymbol{\tau}$, and the ATE, τ, are defined only for the subset of units retained after matching, which often differs from the original set of units prior to matching (Rosenbaum and

Rubin, 1985; Rosenbaum, 2012). This difference is especially important when retention is driven by overlap in the estimated propensity score, as the resulting study population can be difficult to interpret substantively. In such settings, it is often preferable to define the study population directly in terms of a small number of substantively meaningful covariates, reflecting the argument in Rosenbaum (2010) that it is "usually better to go back to the covariates themselves ... perhaps redefining the population under study to be a subpopulation of the original population" (p. 86), as also emphasized by Stuart (2010). Building on this insight, subsequent approaches provide formal procedures for defining the matched study population in this way (Fogarty et al., 2016; Traskin and Small, 2011).

The breadth of the study population retained after matching depends in part on the relative abundance of treated and control units across covariate space. When one treatment arm is sparsely represented in regions dominated by the other, overlap may be limited to a narrow region of covariate space and include relatively few units. All else equal, a larger pool of units available for matching supports overlap over a wider region of covariate space and a larger, more interpretable study population over which the target of inference is defined.

Neither target of inference, $\boldsymbol{\tau}$ nor τ, defined among the matched units, can be directly calculated. Even in a randomized experiment, we cannot assign a country emerging from civil war to receive a UN peacekeeping mission, observe how long the ensuing peace lasts, and then rewind time to observe how long peace would have lasted in the absence of a UN peacekeeping mission. In other words, for each unit we observe only one of its two potential outcomes. Let y_{si} denote this observed outcome, which equals the treated potential outcome when unit i in set s receives the treatment and the control potential outcome otherwise. Because the individual-level causal effect can never be directly observed — only one potential outcome is realized for each unit — we must rely on statistical inference to draw conclusions about $\boldsymbol{\tau}$ and τ.

We consider inference under the sharp and weak frameworks, targeting $\boldsymbol{\tau}$ and τ, respectively. A line of work shows how a single test can be exact for a sharp null and asymptotically valid for the corresponding weak null under as-if randomization (Wu and Ding, 2021; Cohen and Fogarty, 2022; Fogarty, 2023; see Chung and Romano, 2013 for the precursor with independent samples, and Ding, 2017 on conflicts between the two types of tests). The unification of sharp and weak tests does not generally extend to sensitivity analyses outside matched-pair designs (see Fogarty, 2020a, 2023). When researchers must

choose between the two frameworks, the decision depends on both statistical properties and substantive goals.

- Statistically, the sharp framework specifies all missing potential outcomes, allowing exact randomization inference under minimal assumptions. The weak framework leaves some outcomes unspecified and instead relies on variance estimation and a Normal approximation, which can perform poorly in small samples or when outcomes are skewed with extreme outliers. In such cases — or whenever one wants exact p-values under minimal assumptions — permutation inference under the sharp framework may be preferable.
- Substantively, researchers usually test a constant effect in the sharp framework. Such a hypothesis may be unrealistic or of limited scientific interest (see Gelman, 2011, p. 961; Caughey et al., 2023). The weak framework, by contrast, accommodates heterogeneous effects across units, making the ATE a more relevant target in many settings. Nevertheless, tests of a constant effect can provide a useful approximation to a more complex hypothesis with heterogeneous effects (Rosenbaum, 2010, pp. 44–46), and such tests remain valid for bounded null hypotheses, under which heterogeneous effects all lie on one side of the hypothesized constant, for a broad class of test statistics that includes the Difference-in-Means (Caughey et al., 2023, Theorems 1–2).

2.3.0.1 The Assignment Process as the Basis for Inference Regardless of the framework, inference is based on the treatment assignment process. Let z_{si} denote an indicator for whether unit i in matched set s is treated ($z_{si} = 1$) or not ($z_{si} = 0$). We collect these indicators for all units in set s into the vector $\boldsymbol{z}_s = (z_{s1}, \ldots, z_{sn_s})^\top$. Stacking these vectors across all sets gives the full assignment vector $\boldsymbol{z} = (z_{11}, \ldots, z_{Sn_S})^\top$, again indexed first by matched set and then by unit within set. For inference, we condition on the number of treated units within each set, m_s, even if the actual assignment mechanism were to consist of n_s independent assignments. For further discussion of this form of "conditional as-if analysis," see Pashley et al. (2021) and Rosenbaum (2017, pp. 289–290, fn. 15). Forming matched sets from the realized assignment complicates conditional as-if analysis (Pashley et al., 2021, Section 5), a problem that Pimentel and Huang (2024) call $\boldsymbol{Z}$-dependence and that Pimentel and Yu (2026) also examine (see Section 3.1).

We denote by Ω_s the set of all possible treatment assignments in set s, holding fixed the observed

number of treated units in that set. Formally, Ω_s includes every possible way the n_s units in set s could be assigned to treatment and control such that exactly m_s units are treated. The number of possible assignments in Ω_s is denoted by $|\Omega_s|$, where the notation $|\cdot|$ indicates the number of elements in a set. This quantity equals $|\Omega_s| = \binom{n_s}{m_s} = n_s!/[m_s!(n_s - m_s)!]$, where "!" denotes the factorial operator (e.g., $4! = 4 \times 3 \times 2 \times 1$).

In the `ssafrica.3` set, for example, there are 4 observations — 1 treated and 3 control — so $|\Omega_s|$ for the `ssafrica.3` set is $\binom{4}{1} = 4$. The corresponding set of possible assignments with this treated count is shown in Table 2. The column labeled Assignment 3 shows the assignment that actually occurred. The other possible assignments — Assignment 1, Assignment 2, and Assignment 4 — represent cases in which Lesotho, Somalia, or Burundi is treated instead of Rwanda. The set Ω_s for `ssafrica.3` excludes every assignment that does not treat exactly one unit.

	Assignment 1	Assignment 2	Assignment 3	Assignment 4
Burundi	0	0	0	1
Rwanda	0	0	1	0
Somalia	0	1	0	0
Lesotho	1	0	0	0

Table 2: All possible treatment assignments within matched set `ssafrica.3`, holding fixed the observed number of treated units in that set.

The set of possible treatment assignments across all matched sets, given the number treated in each, is $\Omega = \Omega_1 \times \cdots \times \Omega_S$, which consists of all combinations formed by choosing one assignment from each Ω_s. Although the assignment itself can vary, the underlying causal quantities of interest — $\boldsymbol{\tau}$ and τ — remain fixed across all possible assignments. What changes from one assignment to another is which potential outcomes we would actually observe. In the observed Assignment 3, we see Rwanda's treated potential outcome and the control potential outcomes of Burundi, Somalia, and Lesotho, but not Rwanda's control potential outcome or the treated potential outcomes of the others. Under a different assignment, such as Assignment 1 in Table 2, a different set of treated and control potential outcomes would have been observed. No matter which assignment occurs, we observe only partial information about our causal targets.

Inference from the partial information contained in the data to our causal targets is predicated on a probability distribution defined over the set of assignments, Ω. This distribution constitutes the

uncertainty underlying our inferences — what Fisher (1935, p. 14) famously called the "reasoned basis" for inference — but, unlike in a randomized experiment, this distribution is unknown in an observational study. Because this distribution is unknown, we must make assumptions about it when drawing inferences from observational data. Under the assumption of as-if randomization, every possible assignment within each Ω_s is equally likely, independently across sets, making each overall assignment in Ω equally likely as well. In this case, all individuals within the same matched set have the same probability of treatment, equal to m_s/n_s. When as-if randomization does not hold, however, assignments are no longer equally likely, and some individuals within the same matched set have a higher probability of treatment than others.

2.3.1 How Do I Draw Inferences under the Sharp Framework?

To set the stage for inference under the sharp framework, consider a thought experiment. Suppose we were to subtract the true individual effects from the outcomes of whichever units happen to be treated. Doing so would yield, for each unit, the outcome it would have had under control. In other words, we may imagine reconstructing the dataset so that, under any possible treatment assignment, the outcomes would appear exactly as they would have if no one had been treated. In this reconstructed world, there would be no effect since every unit's outcome would reflect what it would have been without treatment.

Of course, we do not know the true collection of individual effects, $\boldsymbol{\tau}$, but we can test hypotheses about it, such as the hypothesis that all units share a common effect, whose hypothesized value we denote by τ_h. We do so by evaluating whether the data would look consistent with no treatment effect when that hypothesized value is subtracted from the treated outcomes. If, after this reconstruction, the data still show a positive effect, then the hypothesized value is presumably too small; if they show a negative effect, then the hypothesized value is presumably too large.

From a hypothesis testing perspective, when outcomes are reconstructed under a null hypothesis, the observed test statistic — computed on these reconstructed outcomes — will tend to fall in the upper tail of the null distribution if the hypothesized effect is too small, and in the lower tail if it is too large. To generate this null distribution, we reconstruct the outcomes under the hypothesized effect. Under the null, these reconstructed outcomes would remain fixed across assignments, so we hold them constant and recalculate the test statistic for every possible assignment. We then use this null distribution to

determine where the observed test statistic falls for the calculation of p-values.

A canonical choice of test statistic in this setting is a *sum statistic*, which first adds up the treated outcomes within each set, and then adds those set-level sums across all sets. Many familiar test statistics can be expressed in this form by applying to the outcomes scale and shift transformations that do not depend on the treatment assignments. One such useful test statistic that can be expressed as a sum statistic is the Difference-in-Means, computed within sets and then averaged across sets, with each set weighted by its contribution to the effective sample size.

Under as-if randomization, the harmonic-mean-weighted Difference-in-Means is useful because, as noted in Section 2.2.3, harmonic-mean weights minimize the variance of the weighted average of within-set Differences-in-Means when treatment effects are homogeneous and outcome variances are equal across sets. Under these conditions, a test has greater power when the test places more weight on sets with large harmonic means — that is, sets in which the Difference-in-Means is most informative. Weighting sets solely by their share of units, by contrast, can overweight large but highly lopsided sets and dilute information from more balanced comparisons. When effects are heterogeneous, the choice of weights has no bearing on the validity of a test of a sharp null, but the choice may bear on the test's power against heterogeneous alternatives and on the interpretation of the Hodges–Lehmann estimate at the end of this section. Under as-if randomization, the expected value of the harmonic-mean-weighted Difference-in-Means is the harmonic-mean-weighted average of set-level effects, the matched-set analogue of the overlap-weighted average effect of Li et al. (2018), rather than the ATE targeted in the weak framework of Section 2.3.2.

The harmonic-mean-weighted Difference-in-Means also has the practical advantage of coinciding with two other common approaches to analyzing matched data:

- First, the harmonic-mean-weighted Difference-in-Means equals the coefficient on the treatment indicator from a fixed-effects regression that includes matched-set indicators (Hansen and Bowers, 2008, p. 229), an approach commonly used in practice after matching.
- Second, the harmonic-mean-weighted Difference-in-Means coincides with the overlap-weighted Difference-in-Means (Li et al., 2018) when overlap weights are constructed using the treatment assignment probabilities implied by as-if randomization. Overlap weighting assigns each control

unit a weight equal to the unit's probability of treatment and each treated unit a weight equal to 1 minus that probability, so that the weighted treated group and the weighted control group place the same total weight on each matched set. Because the assignment probabilities implied by as-if randomization are constant within each matched set, the resulting overlap weights are constant within the treated units and within the control units of each set, and these overlap weights aggregate to weights proportional to the harmonic-mean weights implied by the matched sets.

To implement a sum statistic equivalent to the harmonic-mean-weighted Difference-in-Means for hypothesis testing, we first reconstruct the outcomes that treated units would have exhibited under the null hypothesis $\tau_h = 0$.

```
# Keep only rows assigned to a matched set (drop NA in fm)
data_matched <- filter(
  .data = data, !is.na(fm)
)

# Null hypothesis value
tau_h <- 0

# Reconstruct outcomes under sharp null (tau_h = 0)
data_matched <- data_matched |>
  mutate(
    ldur_tilde = ldur - tau_h * UN
  )
```

We now source a helper function from the companion GitHub repository; this function rescales the outcome variable so that the sum of the rescaled values among treated units equals the harmonic-mean-weighted Difference-in-Means. After sourcing the function, we apply it to the matched dataset to generate a new column containing the rescaled outcomes. We then use this rescaled outcome to compute the observed test statistic.

```
# Load hm_stat_rescale() from GitHub repository
# base_url (defined earlier) points to repository URL
source(
  paste0(
    base_url, "/R/hm_stat_rescale.R"
  )
)

# Apply rescaling: adds ldur_tilde_hm_scaled and returns matched dataset
```

```r
data_matched <- hm_stat_rescale(
  data = data_matched,
  outcome = ldur_tilde, # Set outcome to be rescaled within matched sets
  treat = UN,           # Set name of treatment indicator variable
  strata = fm           # Set name of matched strata (block) variable
)

# Observed HM-weighted diff-in-means statistic
obs_stat <- sum(
  data_matched$ldur_tilde_hm_scaled[data_matched$UN == 1]
)
```

We can verify in R that this observed sum statistic coincides with the harmonic-mean-weighted Difference-in-Means, the coefficient on the treatment indicator from a fixed-effects regression that includes matched-set indicators, and the overlap-weighted Difference-in-Means constructed from the assignment probabilities implied by as-if randomization.

First, we show that the observed sum statistic equals the harmonic-mean-weighted average of the within-matched-set Differences-in-Means, where each matched set is weighted by its contribution to the effective sample size.

```r
# Harmonic-mean-weighted difference in means (weights computed within sets)
dim_hm <- data_matched |>
  group_by(
    fm
  ) |>
  summarize(
    n_treated = sum(UN == 1),
    n_control = sum(UN == 0),
    # Within-set Difference-in-Means
    dim_set   = mean(ldur_tilde[UN == 1]) - mean(ldur_tilde[UN == 0]),
    # Harmonic-mean weight (set's contribution to effective sample size)
    w_hm      = 2 * n_treated * n_control / (n_treated + n_control),
    .groups   = "drop" # Drop grouping after summarize
  ) |>
  summarize(
    # Harmonic-mean-weighted average of within-set Differences-in-Means
    dim_hm = sum(
      w_hm * dim_set) / sum(w_hm)
  ) |>
  pull( # Pull column out of data frame
    dim_hm
  )
```

```r
# Approach coincides with the sum statistic
all.equal(
  dim_hm, obs_stat
)
```

```
[1] TRUE
```

Next, we verify that the same quantity is obtained as the coefficient on the treatment indicator from a fixed-effects regression that includes matched-set indicators.

```r
# FE coefficient on UN (matched-set FE)
fe_fit <- lm(
  formula = ldur_tilde ~ UN + fm,
  data = data_matched
)

# Drop name so comparison is purely numeric
treat_coef_fe_fit <- unname(
  coef(fe_fit)["UN"]
)

# Approach coincides with the sum statistic
all.equal(
  treat_coef_fe_fit, obs_stat
)
```

```
[1] TRUE
```

Finally, we confirm that the observed sum statistic also coincides with the overlap-weighted Difference-in-Means. The overlap weights are constructed from the treatment assignment probabilities implied by as-if randomization within matched sets, with treated units weighted by one minus their treatment probability and control units weighted by their treatment probability.

```r
# Install "PSweight" package (only run if not already installed)
# install.packages("PSweight")
library(PSweight) # For overlap weights from Li et al. (2018)

# Treatment probabilities within matched sets: n_treated / n
ps_fit <- PSmethod(
  ps.formula = UN ~ factor(fm), # Reproduces n_treated / n in each set
  method     = "glm",           # Estimate using generalized linear model
  # Use as.data.frame() for PSmethod() compatibility
  data       = as.data.frame(data_matched),
  ncate      = 2                # Binary treatment (treated vs. control)
)
```

```r
assign_prob <- ps_fit$e.h[, "1"]  # Column for treatment level "1"

# Overlap weights from treatment probabilities
# Controls: treatment probability; Treated: 1 - treatment probability
w_ow <- ifelse(
  test = data_matched$UN == 1,
  yes  = 1 - assign_prob,
  no   = assign_prob
)

# Overlap-weighted difference in means
dim_ow <- data_matched |>
  summarize( # Aggregate rows into summary values
    # Overlap-weighted mean outcome for treated units
    treated_weighted_mean =
      sum(w_ow[UN == 1] * ldur_tilde[UN == 1]) / sum(w_ow[UN == 1]),

    # Overlap-weighted mean outcome for control units
    control_weighted_mean =
      sum(w_ow[UN == 0] * ldur_tilde[UN == 0]) / sum(w_ow[UN == 0]),

    # Difference between overlap-weighted treated and control means
    dim_ow = treated_weighted_mean - control_weighted_mean
  ) |>
  pull(
    dim_ow
  )

# Approach coincides with the sum statistic
all.equal(
  dim_ow, obs_stat
)
```

```
[1] TRUE
```

Now, to generate the distribution to which we refer our observed sum statistic, we can hold the reconstructed and rescaled outcomes fixed and then calculate the sum statistic over all possible assignments in Ω. In our application, with 12 sets ranging in size from 2 to 5, the total number of assignments is 311,040.

```r
# For each matched set (fm), record:
# n = total units in the set
# m = number treated (UN == 1)
block_ns <- data_matched |>
  group_by(          # Group results by key variables
```

```
    fm
  ) |>
    summarize(          # Aggregate to one row per group
    n = n(),          # Row count per group
    m = sum(UN),
    .groups = "drop"
  )

# Total possible treatment assignments = product of binomial coefficients
# (choose m_s of the n_s units in each set, then multiply across sets)
exact_n_assigns <- prod(
  choose(
    n = block_ns$n,
    k = block_ns$m
  )
)
```

Because our matched study is relatively small, we can enumerate all possible treatment assignments exactly.

```
# Install "randomizr" (only run if not already installed)
# install.packages("randomizr")

# Load randomizr for generating random assignments
library(randomizr)

exact_assigns <- obtain_permutation_matrix(
  declaration = declare_ra(    # Declare assignment procedure
    N = nrow(data_matched),    # Total number of units
    blocks = data_matched$fm, # Matched set membership
    block_m = block_ns$m       # Number treated in each set
  ),
  # Total number of assignments with treated counts fixed within sets
  maximum_permutations = exact_n_assigns
)
```

However, in most applications, exactly enumerating all possible assignments is computationally infeasible. Instead, we typically draw a random sample of assignments (e.g., 10,000) to approximate the exact randomization distribution, as the following code illustrates.

```
# Set RNG seed for reproducibility
set.seed(11242017)

# Generate permutation matrix of treatment assignments
# Each column = one possible assignment consistent with block structure
```

```r
sim_assigns <- obtain_permutation_matrix(
  declaration = declare_ra(
    N = nrow(data_matched),
    blocks = data_matched$fm,
    block_m = block_ns$m
  ),
  maximum_permutations = 10^4  # Cap at 10,000 random draws
)
```

Now, to generate the null distribution of the sum statistic, we apply the statistic to each of these 10,000 assignments while holding the reconstructed and rescaled outcomes fixed under the null.

```r
# Randomization distribution under sharp null
# Apply sum statistic to each assignment in sim_assigns
# Outcome transformed so sum = harmonic-mean-weighted difference-in-means
sim_sharp_null_dist <- apply(
  X = sim_assigns,     # Matrix of treatment assignments
  MARGIN = 2,          # Iterate over columns (assignments)
  FUN = function(x) {
    # Sum transformed outcomes among treated
    sum(
      data_matched$ldur_tilde_hm_scaled[x == 1]
    )
  }
)
# Faster equivalent computation via matrix multiplication
# as.numeric(
#   t(data_matched$ldur_tilde_hm_scaled) %*% sim_assigns
# )
```

From this null distribution, we compute an upper one-sided p-value, which, under as-if randomization, is simply the proportion of null test statistics greater than or equal to the observed test statistic.

```r
# One-sided, upper p-value:
# Proportion of simulated randomization stats >= observed
round(
  x = mean(
    sim_sharp_null_dist >= obs_stat
  ),
  digits = 4
)
```

```
[1] 0.0346
```

The upper p-value of a test of the sharp null of no effect against the alternative of a larger effect is 0.0346.

In this particular case, unlike in most applications, the matched study is small enough to compute the exact p-value and assess the accuracy of the simulation-based p-value approximation.

```
# Transformed outcomes under sharp null (tau_h = 0)
q_tau_h_0 <- data_matched$ldur_tilde_hm_scaled

# Fast computation of exact null distribution using matrix multiplication
exact_sharp_null_dist <- as.numeric(
  t(q_tau_h_0) %*% exact_assigns
)

# Slower apply()-based computation (for reference)
# apply(
#  X = exact_assigns,
#  MARGIN = 2,
#  FUN = function(x) {
#    sum(data_matched$ldur_tilde_hm_scaled[x == 1])
#  }
#)

# Exact one-sided, upper p-value
round(
  x = mean(
    exact_sharp_null_dist >= obs_stat
  ),
  digits = 4
)
```

```
[1] 0.0363
```

The exact p-value of 0.0363 is nearly identical to the simulation-based p-value of 0.0346. At the conventional significance level of $\alpha = 0.05$, we would reject the null hypothesis in favor of the alternative, regardless of whether we use the exact or simulation-based p-value.

As an alternative to randomly sampling assignments and computing the test statistic for each one, we can use a much faster Normal approximation to the null distribution when the matched study contains many matched sets. The approximation relies on closed-form expressions for the expected value and variance of a sum statistic derived in Rosenbaum and Krieger (1990). Using these expressions, we standardize the observed test statistic and then compare it to the standard Normal distribution, which gives us a corresponding p-value. Although primarily designed for sensitivity analyses under violations of as-if randomization, the `senstrat` package can also be used to compute the null expected value and variance

of a sum statistic under as-if randomization.

```r
# Install "senstrat" package (only run if not already installed)
# install.packages("senstrat")

# Load senstrat for stratum-level null moments and sensitivity analysis
# (Rosenbaum & Krieger 1990)
library(senstrat)

# Compute per-block null expectations and variances
per_block_moms <- data_matched |>
  group_by(
    fm
  ) |>
  summarize(
    expect   = ev(
      sc = ldur_tilde_hm_scaled, # Transformed outcomes for the stratum
      z = UN,                    # Treatment indicator
      m = 1,                     # Count of 1s in hidden confounder
                                 # Irrelevant here since Gamma = 1
      g = 1,                     # Sensitivity parameter Gamma
      method = "RK"              # Use Rosenbaum and Krieger (1990) formula
    )$expect,                    # Null expectation of sum statistic by set
    variance = ev(
      sc     = ldur_tilde_hm_scaled,
      z      = UN,
      m = 1,
      g      = 1,
      method = "RK"
    )$vari,                      # Null variance of sum statistic by set
    .groups  = "drop"
  )

# Sum across blocks to get overall null expectation
null_ev  <- sum(
  per_block_moms$expect
)

# Sum across blocks to get overall null variance
null_var <- sum(
  per_block_moms$variance
)

# Standardized test statistic and one-sided Normal p-value (upper tail)
norm_upper_p_value <- pnorm(
  q = (obs_stat - null_ev) / sqrt(null_var), # Standardized statistic
  lower.tail = FALSE                         # Compute upper-tail prob
)
```

```r
# By default in pnorm(): mean = 0 and sd = 1 (standard normal distribution)

# Display the Normal-approximation p-value
round(
  x = norm_upper_p_value,
  digits = 4
)
```

```
[1] 0.0343
```

This Normal-approximation p-value of 0.0343 is close to the exact and simulation-based p-values of 0.0363 and 0.0346, respectively. All three p-values lead us to reject the sharp null of no effect in favor of a larger effect.

To test not only a single null hypothesis but a grid of null hypotheses under the sharp framework, we repeat the Normal-approximation test for a range of hypothesized constant effects, τ_h. For each value of τ_h, we reconstruct the outcomes under the null, rescale them so that the sum statistic equals the harmonic-mean-weighted Difference-in-Means, compute the observed test statistic and the statistic's null expectation and variance, and obtain the corresponding p-value using the Normal approximation. To form a one-sided confidence set, we retain all null values that are not rejected by the upper-tail test at the chosen α level. To form a two-sided confidence set, we proceed analogously, allocating $\alpha/2$ to each tail — reflecting that each null value is tested against departures in both directions — and using the corresponding upper- and lower-tail rejection regions.

```r
# Significance level
alpha <- 0.05

# Grid of constant-effect null values for the Normal approximation
tau_h_grid_norm <- seq(
  from = -0.5,
  to = 2,
  by = 0.001
)

# Repeat the Normal-approximation test for each tau_h
# sapply() computes and stacks the resulting p-values
p_mat_norm <- sapply(X = tau_h_grid_norm,
                     FUN = function(tau_h) {

                       # Reconstruct and rescale outcomes under null tau_h
                       dat_tau_h <- hm_stat_rescale(
```

```
                    data    = transform(
                      data_matched,
                      ldur_tilde_shift = ldur - tau_h * UN),
                    outcome = ldur_tilde_shift,
                    treat   = UN,
                    strata  = fm
                  )

                  # Observed statistic under null tau_h
                  obs_stat_tau_h <- sum(
                    dat_tau_h$ldur_tilde_shift_hm_scaled[
                      dat_tau_h$UN == 1]
                  )

                  # Null expectation and variance under null tau_h
                  moms_tau_h <- dat_tau_h |>
                    group_by(
                      fm
                    ) |>
                    summarize(
                      expect   = ev(sc = ldur_tilde_shift_hm_scaled,
                                    z = UN, m = 1, g = 1,
                                    method = "RK")$expect,
                      variance = ev(sc = ldur_tilde_shift_hm_scaled,
                                    z = UN, m = 1, g = 1,
                                    method = "RK")$vari,
                      .groups  = "drop"
                    )

                  # Standardized statistic under null tau_h
                  std_stat_tau_h <- (obs_stat_tau_h -
                                       sum(moms_tau_h$expect)) /
                    sqrt(sum(moms_tau_h$variance))

                  # Upper- and lower-tail Normal p-values
                  c(
                    upper_tail = pnorm(q = std_stat_tau_h,
                                       lower.tail = FALSE),
                    lower_tail = pnorm(q = std_stat_tau_h)
                  )
                })

# Upper-tail confidence set: retain all tau_h with upper-tail p >= alpha
cs_sharp_upper_tail_norm <- c(
  lower = min(
    tau_h_grid_norm[p_mat_norm["upper_tail", ] >= alpha]
  ),
  upper = Inf
```

```
)

# Two-sided confidence set: retain tau_h only if neither one-sided test
# rejects at level alpha / 2
cs_sharp_two_sided_norm_vals <- tau_h_grid_norm[
  p_mat_norm["upper_tail", ] >= alpha / 2 &
    p_mat_norm["lower_tail", ] >= alpha / 2
]

# Two-sided confidence set summarized by its bounds
cs_sharp_two_sided_norm <- c(
  lower = min(
    cs_sharp_two_sided_norm_vals
  ),
  upper = max(
    cs_sharp_two_sided_norm_vals
  )
)
```

An analogous procedure applies when constructing confidence sets without relying on a Normal approximation, instead using a simulation-based approximation to the null randomization distribution.

```
# Grid of constant-effect null values (sharp framework)
tau_h_grid <- seq(
  from = -0.5,
  to = 1.5,
  by = 0.0001
)

# Repeat the randomization test for each tau_h
# sapply() computes and stacks the resulting p-values
p_mat <- sapply(X = tau_h_grid,
                FUN = function(tau_h) {

                  # Shift outcomes under null: ldur_i - tau_h * UN_i
                  dat_tau_h <- hm_stat_rescale(
                  # transform(): copy data_matched adding shifted outcome
                    data    = transform(
                      data_matched,
                      ldur_tilde_shift = ldur - tau_h * UN),
                    outcome = ldur_tilde_shift,
                    treat   = UN,
                    strata  = fm
                  )

                  # Transformed outcomes under sharp null tau_h
                  q_tau_h <- dat_tau_h$ldur_tilde_shift_hm_scaled
```

```
                    # Observed statistic under null tau_h
                    obs_stat_tau_h <- sum(
                      dat_tau_h$UN * q_tau_h
                    )

                    # Randomization distribution via simulated assignments
                    # Matrix multiplication used for speed (vs apply())
                    sim_null_dist_tau_h <- as.numeric(
                      t(q_tau_h) %*% sim_assigns
                    )

                    # Upper-tail and lower-tail randomization p-values
                    p_upper_tau_h <- mean(
                      sim_null_dist_tau_h >= obs_stat_tau_h
                    )

                    p_lower_tau_h <- mean(
                      sim_null_dist_tau_h <= obs_stat_tau_h
                    )

                    c(
                      upper_tail = p_upper_tau_h,
                      lower_tail = p_lower_tau_h
                    )
                  })

# Extract vector of upper-tail p-values
p_upper <- p_mat["upper_tail", ]

# Extract vector of lower-tail p-values
p_lower <- p_mat["lower_tail", ]

# Simulation-based confidence sets

# Upper-tail confidence set: retain all tau_h with upper-tail p >= alpha
cs_sharp_upper_tail_sim <- tau_h_grid[p_upper >= alpha]

# Lower bound of the upper-tail confidence set
cs_sharp_upper_tail_sim_bound <- min(
  cs_sharp_upper_tail_sim
)

# Two-sided confidence set (inversion using alpha/2 in each tail):
# retain tau_h only if neither one-sided test rejects at level alpha/2
cs_sharp_two_sided_sim <- tau_h_grid[
  p_upper >= alpha / 2 & p_lower >= alpha / 2
]
```

```r
# Two-sided confidence set summarized by its bounds
cs_sharp_two_sided_sim_bounds <- c(
  lower = min(
    cs_sharp_two_sided_sim
  ),
  upper = max(
    cs_sharp_two_sided_sim
  )
)
```

With these simulated assignments, the one-sided 95% confidence set for a constant effect is $[0.069, \infty)$ and the two-sided 95% confidence set is $[-0.066, 1.358]$. The Normal approximation gives $[0.077, \infty)$ and $[-0.064, 1.364]$, respectively.

Likewise, for a point estimate, one could follow Hodges and Lehmann (1963) and Rosenbaum (1993) by identifying the null value that makes the observed sum statistic equal to its null expectation. With our test statistic under as-if randomization, the null expectation is 0 for every τ_h, and the observed statistic equals 0 exactly when τ_h equals the harmonic-mean-weighted Difference-in-Means computed from the observed outcomes (roughly 0.673). This harmonic-mean-weighted Difference-in-Means is therefore the Hodges–Lehmann estimate.

2.3.2 How Do I Draw Inferences under the Weak Framework?

When the target is the ATE, it can be written as $\tau = \sum_{s=1}^{S}(n_s/n)\tau_s$, a weighted average of the set-level ATEs, $\tau_s = (1/n_s)\sum_{i=1}^{n_s}\tau_{si}$, with weights equal to each set's share of the total study size. Thus, a straightforward way to estimate the ATE is to compute the Difference-in-Means within each matched set, and then combine these set-level estimates using the same weights. Formally, the overall Difference-in-Means is $\hat{\tau} = \sum_{s=1}^{S}(n_s/n)\hat{\tau}_s$, where $\hat{\tau}_s$ is the Difference-in-Means in set s.

These weights are appealing because, under as-if randomization, the resulting Difference-in-Means is an unbiased estimator of the ATE. Formally, the estimator's expected value — i.e., the average of $\hat{\tau}$ taken over all treatment assignments in Ω, each of which is equally likely under as-if randomization — equals the true ATE. Although correct in expectation, the Difference-in-Means can vary substantially across assignments. In any given assignment, the estimate may lie far from the target ATE.

Because the estimator can fluctuate across different treatment assignments, it is important to quantify the

typical squared distance between an estimate and the true ATE. To this end, Neyman (1923) introduced a canonical variance estimator. Under as-if randomization, this estimator is exactly unbiased when individual treatment effects are homogeneous; otherwise, it is conservative, meaning its expected value is greater than or equal to the true variance of the Difference-in-Means, although subsequent refinements can reduce this conservativeness while maintaining validity (see Robins, 1988; Aronow et al., 2014; Harshaw et al., 2026). In principle, we could apply this approach by estimating the variance of the Difference-in-Means within each matched set and then taking a weighted sum of these estimates (see, e.g., Miratrix et al., 2013).

In our setting, however, this approach is infeasible because each matched set contains only 1 treated or 1 control unit. The usual Neyman formula relies on computing sample variances separately within the treated and control groups of each set. The sample variance requires at least two observations because `var()` divides by the number of observations minus 1. As a result, we cannot compute the sample variance when there is only a single treated or control unit.

How, then, can we estimate the variance of the Difference-in-Means in a matched observational study? Pashley and Miratrix (2021) and Fogarty (2018) offer distinct solutions. Pashley and Miratrix (2021) propose two approaches, each valid under different conditions on the matched structure, while Fogarty (2018) develops a single method that applies more broadly to finely stratified designs.

The first approach in Pashley and Miratrix (2021), `hybrid_m`, requires at least two matched sets of each unique size among the sets with one treated or one control unit. The second approach, `hybrid_p`, permits variation in set sizes, provided that no single set with one treated or one control unit contains half or more of the units in all such sets. In our design, every set has one treated or one control unit, so the condition requires only that no set contain half or more of the 36 matched units, and our largest set contains 5. Our matched design meets the condition for the second approach, not the first. We therefore estimate the variance using `hybrid_p` through the `blkvar` package, the companion software for Pashley and Miratrix (2021).

```
# Install blkvar (only run if not already installed)
# install.packages("remotes")
# remotes::install_github("lmiratrix/blkvar")

# Load blkvar for block randomization variance estimators
```

```r
library(blkvar)

# Compute results with hybrid_p method
res <- block_estimator(
  Yobs = ldur,          # Observed outcomes
  Z = UN,               # Treatment indicator
  B = fm,               # Block (matched set) membership
  data = data_matched, # Dataset
  method = "hybrid_p"  # Variance estimation method
)

# Extract variance estimate
res$var_est
```

```
[1] 0.1187282
```

We can also estimate the variance of the Difference-in-Means in a finely stratified design using the approach of Fogarty (2018), with set sizes incorporated as in Fogarty (2023, p. 2199). To do so, we load a custom `R` function, `fine_strat_var_est()`, from the companion GitHub repository.

```r
# Load the fine_strat_var_est() function from the GitHub repository

source(
  paste0(
    base_url, "/R/fine_strat_var_est.R"
  )
)
```

The function takes as inputs the set-specific treatment effect estimates and the number of units in each set, and returns a single scalar representing the estimated variance. We compute the set-specific estimates and corresponding set sizes, and then pass these two vectors as inputs to `fine_strat_var_est()`.

```r
# Compute matched set sizes and matched-set-specific differences-in-means
set_stats <- data_matched |>
  group_by(
    fm
  ) |>
  summarize(
    n = n(),                                # Matched set size
    diff_in_means = mean(                   # Treated mean
      ldur[UN == 1L]
    ) -                                     # minus
      mean(                                 # control mean
        ldur[UN == 0L]
```

```
      ),
    .groups = "drop"
  )

# Apply Fogarty (2018/2023) variance estimator
fine_strat_var_est(
  strat_ns = set_stats$n,                # Vector of stratum sizes
  strat_ests = set_stats$diff_in_means  # Vector of stratum estimates
)
```

```
[1] 0.1163934
```

This variance estimate is nearly identical to the one we obtained using the `hybrid_p` approach of Pashley and Miratrix (2021).

Now that we have estimates of both the ATE and the variance of the ATE estimator, we can form a standardized test statistic by subtracting the expected Difference-in-Means implied by the null and dividing the result by the estimated standard error (the square root of the estimated variance). We then compare this statistic to a standard Normal distribution to calculate p-values. The following code calculates the upper one-sided p-value for a test of the null hypothesis that the ATE is 0 against the alternative that the ATE is positive.

```
pnorm(
  q = (res$ATE_hat - 0) / sqrt(res$var_est),
  lower.tail = FALSE
)
```

```
[1] 0.03722157
```

Here, we reject the weak null hypothesis that the ATE equals 0 in favor of the alternative that the ATE is greater than 0, using the conventional significance level of $\alpha = 0.05$.

As in the construction of confidence sets for a homogeneous treatment effect under the sharp framework, we form 95% confidence sets by inverting the corresponding hypothesis tests. We first construct a one-sided confidence set by inverting the upper-tail test and retaining all null values that are not rejected at the $\alpha = 0.05$ level. We then construct a two-sided confidence set by allocating $\alpha/2 = 0.025$ to each tail.

```
# One-sided (upper-tail) confidence set
cs_weak_upper_tail <- c(
```

```
  lower = res$ATE_hat - qnorm(p = 1 - alpha) * sqrt(res$var_est),
  upper = Inf
)

# Two-sided confidence set
cs_weak_two_sided <- c(
  lower = res$ATE_hat - qnorm(p = 1 - alpha / 2) * sqrt(res$var_est),
  upper = res$ATE_hat + qnorm(p = 1 - alpha / 2) * sqrt(res$var_est)
)
```

The estimated ATE (`res$ATE_hat`) is 0.615, with an estimated standard error of 0.345. The resulting one-sided 95% confidence set is $[0.048, \infty)$. The two-sided 95% confidence set, $[-0.061, 1.290]$, includes 0 because the one-sided p-value of 0.037 exceeds $\alpha/2 = 0.025$.

2.4 Third Stage: Sensitivity Analysis (Departures from As-If Randomization)

Up to this point, we have assumed a framework in which each observation's chance of a UN intervention is like flipping a weighted coin. Each coin flip is independent across observations, but the probability of landing tails (i.e., receiving treatment) can differ from one unit to another depending on each unit's baseline characteristics. With matching, we make our crucial assumption that all units within a set are similar enough on these characteristics that their coins have the same probability of landing tails.

This as-if randomization assumption may fail because of imbalances on hidden covariates (or residual imbalances on observed covariates, though the sensitivity analysis to follow subsumes both within the same framework). To represent such imbalances, consider a single hidden covariate, $\boldsymbol{u} = (u_{11}, \ldots, u_{Sn_S})^\top$, indexed first by matched set and then by unit within set, with each u_{si} constrained to lie in the interval from 0 to 1. Although the restriction that each element of $\boldsymbol{u}$ is between 0 and 1 may seem strong, Rosenbaum (2017, p. 300, fn. 33) shows that any departure from complete random assignment within blocks that leaves every unit's probability of treatment strictly between 0 and 1 can be represented by such a $\boldsymbol{u}$ in the sense that as-if randomization would hold if we had access to and exactly matched on this $\boldsymbol{u}$.

Rosenbaum (1987a) and Rosenbaum and Krieger (1990) then propose a model in which each unit's

independent probability of treatment assignment is given by

$$\pi_{si} = \frac{\exp\left[\kappa_s + \log(\Gamma)u_{si}\right]}{1 + \exp\left[\kappa_s + \log(\Gamma)u_{si}\right]}. \tag{3}$$

The parameter κ_s is a set-specific intercept that captures the baseline propensity for treatment shared by all units in matched set s before accounting for any differences in the hidden covariate. In other words, κ_s reflects the central idea in matching that, after forming matched sets homogeneous in observed covariates, all individuals within a set share the same treatment propensity based on those covariates. The parameter $\Gamma \geq 1$, by contrast, quantifies how strongly the hidden covariate u_{si} can alter treatment odds. When $\Gamma = 1$, all individuals in matched set s have the same probability of treatment, $\exp[\kappa_s]/(1 + \exp[\kappa_s])$. However, when $\Gamma > 1$, two individuals in the same set who differ in the hidden covariate may differ in their odds of treatment by as much as a factor of Γ.

An important point to reiterate is that we condition on assignments that belong to the set Ω, meaning that the number of treated units is fixed within each matched set. This conditioning removes dependence on the set-specific baseline κ_s in the probability distribution over assignments in Ω (Rosenbaum, 1984; Rosenbaum and Krieger, 1990). Eliminating this dependence is crucial because, without the set-specific intercepts, we can characterize both as-if randomization and departures from as-if randomization using only a single sensitivity parameter, Γ.

To build intuition for the sensitivity parameter Γ, note that the model in (3) implies the following restriction:

$$\frac{1}{\Gamma} \leq \frac{\pi_{si}\left(1 - \pi_{sj}\right)}{\pi_{sj}\left(1 - \pi_{si}\right)} \leq \Gamma \text{ for all } i,\ j \text{ and } s. \tag{4}$$

This restriction states that, within any set, no two units can differ in their odds of treatment by more than a factor of Γ. When $\Gamma = 1$, all units share the same odds of treatment, corresponding to as-if randomization. By contrast, larger values of Γ represent increasingly severe departures from this assumption.

Rosenbaum (1995, pp. 1424–1425) shows that the converse also holds, that is, that the restriction in (4) implies a model of the form in (3). For any collection of treatment probabilities satisfying the bound in

(4), it is always possible to find values of $u_{si} \in [0, 1]$ for all sets s and units i, together with set-specific intercepts κ_s, such that (3) holds with the same Γ and therefore yields the same probability distribution over assignments in Ω. In other words, (4) describes a general restriction on treatment probabilities that does not assume a particular functional form. The logistic model in (3), by contrast, provides one convenient parametric representation of that general restriction.

Thus far, we have conducted inference assuming $\Gamma = 1$, meaning that all units within a matched set share the same treatment probability. Under this as-if randomization assumption, tests of sharp null hypotheses based on the exact randomization distribution are exactly valid, while tests of weak null hypotheses are valid asymptotically, as the number of matched sets grows with set sizes bounded (Fogarty, 2018, 2023). In both cases, validity means that the probability of falsely rejecting the null hypothesis does not exceed the nominal level α.

When we allow departures from as-if randomization, governed by the sensitivity parameter $\Gamma \geq 1$, we seek new p-values that remain valid in the same sense. The probability of a false rejection should not exceed α, regardless of the hidden covariate $\boldsymbol{u}$ or the potential outcomes consistent with the null. The way we ensure this validity, however, differs between the sharp and weak frameworks, which we consider in turn.

2.4.1 How Do Inferences under the Sharp Framework Change under These Departures?

For a sharp null, if the hidden $\boldsymbol{u}$ were known, we could calculate from (3) the exact distribution of assignments consistent with the matched design for any given value of $\Gamma \geq 1$. This distribution would provide the correct reference for computing p-values under the null. In particular, the upper one-sided p-value could be obtained by summing the probabilities of all assignments whose test statistic under the null is at least as large as the observed statistic.

In practice, however, $\boldsymbol{u}$ is unknown. Thus, to ensure validity of our tests, we compute p-values under the worst-case choice of $\boldsymbol{u}$ — the one that makes the p-value as large as possible. Rejection under the worst-case choice of $\boldsymbol{u}$ guarantees rejection under the actual (but hidden) $\boldsymbol{u}$. Consequently, we will not reject the null more often than we would if the true assignment probabilities — governed by the actual but unobserved $\boldsymbol{u}$ — were known under the model in (3).

2.4.1.1 Finding the Worst-Case Scenario of Hidden Bias for Valid Inference

Finding this worst-case $\boldsymbol{u}$ exactly is difficult. As a first step, Rosenbaum and Krieger (1990) show that in the unmatched (i.e., two-sample) case, for a broad class of test statistics that includes sum statistics, the p-value under any fixed $\Gamma \geq 1$ is maximized by a vector $\boldsymbol{u}$ in a restricted set of possibilities, denoted by $\mathcal{U}_+$. Once the subjects are ordered from the largest to the smallest outcome, all the candidate vectors in $\mathcal{U}_+$ look the same: a sequence of 1s at the top followed by 0s below. We do not know *how many* 1s should precede the 0s, and this number determines which configuration of $\boldsymbol{u}$ yields the worst-case p-value. In an unmatched study, it is straightforward to enumerate all $n-1$ such candidate vectors in $\mathcal{U}_+$, and then identify which one yields the largest p-value for a fixed $\Gamma \geq 1$.

Unfortunately, in matched designs, the overall worst-case vector $\boldsymbol{u}$ cannot be obtained by simply stitching together the worst-case vectors from each matched set. That is, the global worst case is not simply the collection of local worst cases. Instead, $\mathcal{U}_+$ consists of $\prod_{s=1}^{S}(n_s - 1)$ total candidate vectors for the worst-case $\boldsymbol{u}$ (Rosenbaum and Krieger, 1990, p. 496) — a quantity that quickly becomes infeasible to enumerate directly. For example, with only 20 matched sets of 4 units each, the number of candidate vectors is already nearly 3.5 billion ($3^{20} = 3{,}486{,}784{,}401$).

2.4.1.1.1 Separable Approximation

To address this challenge, Gastwirth et al. (2000) propose a practical shortcut called the *separable approximation*. The idea is to select, within each matched set, the configuration $\boldsymbol{u}_s$ (from the set-specific candidate sequences of 1s and 0s) that maximizes the expected value of the test statistic in that set under the null. If more than one candidate yields the same expectation, the choice goes to the $\boldsymbol{u}_s$ that produces the larger variance of the test statistic in that set under the null. The method is called "separable" because it then stitches together the choices of $\boldsymbol{u}_s$ made separately within each matched set, rather than searching over all possible combinations across sets. The resulting $\boldsymbol{u}$ may not give the worst-case p-value over $\mathcal{U}_+$; yet as the number of matched sets grows, with set sizes bounded, the error becomes negligible under regularity conditions (Gastwirth et al., 2000, Proposition 1).

2.4.1.1.2 Taylor Series Approximation

The separable approximation is a useful shortcut, but the separable approximation's p-value is never larger than, and typically slightly smaller than, the worst-case p-value over $\mathcal{U}_+$ computed with the same Normal approximation (Rosenbaum, 2018, pp. 2321–2322). No

result guarantees that the difference between these two p-values is negligible in designs with relatively few matched sets or with highly unbalanced set sizes. Rosenbaum (2018) therefore introduces a refinement that is never less conservative than this worst case, so that the resulting test is valid whenever the Normal approximation is adequate.

The key idea is to reframe hypothesis testing in terms of a function that, for each configuration of hidden bias $\boldsymbol{u} \in \mathcal{U}_+$, combines the gap between the null expectation and the observed test statistic with a corresponding Normal critical buffer. Concretely, for $\alpha = 0.05$ and corresponding critical value 1.64, this function is the expected value of the test statistic under $(\boldsymbol{u}, \Gamma)$ minus the observed test statistic, plus 1.64 times the standard deviation of the test statistic under $(\boldsymbol{u}, \Gamma)$. The function is nonpositive exactly when the observed test statistic lies at least 1.64 standard deviations above its null expectation, and positive otherwise. Therefore, for a given $(\boldsymbol{u}, \Gamma)$, the Normal-approximation test rejects the null hypothesis if and only if this function is nonpositive, and the sensitivity analysis at Γ rejects if and only if the function is nonpositive for every $\boldsymbol{u} \in \mathcal{U}_+$, that is, if and only if the function's maximum over $\mathcal{U}_+$ is nonpositive.

We can code this function in `R` as follows.

```r
# Rosenbaum (2018) objective: expectation - observed + kappa * SD
# Nonpositive values imply rejection at level alpha
sens_objective <- function(null_expect,
                           null_variance,
                           obs_stat,
                           alpha = 0.05) {
  # null_expect   : overall null expectation of the test statistic
  # null_variance : overall null variance of the test statistic
  # obs_stat      : observed value of the test statistic
  # alpha         : test size (e.g., 0.05 gives kappa about 1.64)

  # Normal critical value (not the set intercept kappa_s in (3))
  kappa  <- qnorm(
    p = 1 - alpha
  )

  # Null standard deviation
  null_sd <- sqrt(
    null_variance
  )

  # Gap between null expectation and observed statistic
```

```
  # (null_expect - obs_stat) + Normal buffer (kappa * null_sd)
  obj_val <- (null_expect - obs_stat) + kappa * null_sd

  return(obj_val)
}
```

To build intuition, we next evaluate this function at $\Gamma = 1$.

```
# Evaluate under Gamma = 1 using null_ev, null_var, and obs_stat
sens_objective(
  null_expect   = null_ev,
  null_variance = null_var,
  obs_stat      = obs_stat,
  alpha         = 0.05
)
```

```
[1] -0.06500212
```

The resulting value is negative, indicating rejection of the null at the chosen significance level. This negative value is exactly what we would expect. Because we previously rejected the sharp null hypothesis that $\tau_h = 0$ for all units at $\Gamma = 1$, the function evaluated at $\Gamma = 1$ is indeed nonpositive.

The crucial property of this function is concavity. For a configuration $\boldsymbol{u} \in \mathcal{U}_+$, let μ and ν denote the null expectation and null variance of the test statistic, and let t denote the observed test statistic. For $\alpha = 0.05$, the function is $\mu - t + 1.64\sqrt{\nu}$, which is concave in (μ, ν) because the function is linear in μ and $\sqrt{\nu}$ is concave in ν. Hence the tangent plane to this function at any point (μ^*, ν^*) with $\nu^* > 0$ lies on or above the function at every (μ, ν), and in particular at the (μ, ν) of every configuration in $\mathcal{U}_+$. Rosenbaum (2018, pp. 2322–2323) obtains this linear bound as a first-order Taylor approximation at the configuration chosen by the separable approximation. If the maximum of the tangent plane over $\mathcal{U}_+$ is nonpositive, then the function itself is nonpositive for every $\boldsymbol{u} \in \mathcal{U}_+$. Consequently, rather than maximizing a nonlinear function over $\mathcal{U}_+$, we can instead maximize a linear function, which greatly simplifies the optimization.

The tangent plane is easy to work with because it is linear in μ and ν, which are sums of set-specific null expectations and variances, so the tangent plane decomposes additively across matched sets. We can therefore determine, within each matched set separately, which hidden-bias pattern makes the tangent plane as large as possible, and then add up these set-specific contributions. The resulting sum is the

maximum value of the tangent plane over all $\boldsymbol{u} \in \mathcal{U}_+$.

Because the function lies on or below the tangent plane, if the maximum value of the tangent plane over $\boldsymbol{u} \in \mathcal{U}_+$ is at or below zero, then the function itself must be nonpositive for every $\boldsymbol{u} \in \mathcal{U}_+$. Therefore, provided that the Normal approximation is adequate, we can safely base the decision to reject the null hypothesis or not on the Taylor series approximation. Although valid, this procedure can be conservative, because the tangent plane may remain above zero even when the function would be below zero at the true worst-case $\boldsymbol{u}$. By contrast, the separable approximation can be liberal, as it may reject even when the worst-case value of the function is positive.

2.4.1.2 Conducting Worst-Case Sensitivity Analysis

We next report the upper one-sided p-value obtained under the separable approximation, along with an indication of whether the resulting decision to reject or not reject the null hypothesis agrees with the decision implied by the Taylor series approximation. The decision implied by the Taylor series approximation is determined by the sign of the Taylor series bound on the objective function that `sens_objective()` codes. The `senstrat()` function reports this bound, stored as `cvA` in the following code. Disagreement can occur only when the separable approximation yields a p-value at or below α (in practice, typically only slightly below), indicating rejection, while the Taylor series approximation yields a positive value, leading to non-rejection.

```
# Grid of Gamma values
Gamma_vals <- seq(
  from = 1,
  to = 1.5,
  by = 0.0001
)

# lapply() runs sensitivity analysis at each Gamma, returns list of results
sens_results <- lapply(
  X = Gamma_vals,
  FUN = function(g) {
    out <- senstrat(
      sc          = data_matched$ldur_tilde_hm_scaled, # outcome
      z           = data_matched$UN,                   # treatment indicator
      st          = data_matched$fm,                   # matched set
      gamma       = g,                                 # Gamma
      alternative = "greater",                         # upper-tail test
      level       = alpha,                             # test level for cvA
      detail      = TRUE # return intermediate quantities
    )
```

```
    # Separable p-value
    p_sep <- as.numeric(
      out$Separable["P-value"]
    )

    # Taylor series margin (cvA)
    cvA <- as.numeric(
      out$lambda["Linear Taylor bound"]
    )

    # Decisions
    sep_reject <- p_sep <= alpha          # separable rejects?
    tay_reject <- cvA <= 0                # Taylor rejects?

    # Store results for this value of Gamma
    data.frame(
      Gamma     = g,         # sensitivity parameter
      p_sep     = p_sep,     # separable p-value
      cvA       = cvA,       # Taylor bound
      agree_tay = (sep_reject == tay_reject) # agreement indicator
    )
  }
)

# Bind into a single data frame
sens_df <- do.call(
  what = rbind,
  args = sens_results
)

# Smallest Gamma where the separable p-value is >= alpha
sens_value_sep <- min(
  sens_df$Gamma[sens_df$p_sep >= alpha]
)

# Taylor series sensitivity value:
# smallest Gamma at which Taylor margin becomes positive (non-rejection)
sens_value_tay <- min(
  sens_df$Gamma[sens_df$cvA > 0]
)
```

For this matched dataset, the conclusions from the separable and Taylor series approximations agree for nearly all values of Γ, differing over only a narrow interval from 1.1361 to 1.1399. As these results indicate, we can reject the sharp null of no effect under as-if randomization (i.e., $\Gamma = 1$), but this conclusion is sensitive to departures from that assumption. Using the Normal approximation together with the more

conservative Taylor series approximation, we can no longer reject the sharp null of no effect against the alternative of a larger effect at the $\alpha = 0.05$ level for values of Γ of approximately 1.14 or larger.

Figure 3 illustrates this comparison and clarifies how the two approaches relate across values of Γ. The results in the shaded vertical band — corresponding to values of Γ for which the two approaches yield different conclusions (between 1.1361 and 1.1399) — show that the two methods diverge in exactly the way one would expect. Over this range of Γ values, the Taylor series margin (the bound `cvA`) lies just above zero, indicating no rejection of the sharp null, while the separable approximation yields p-values just below 0.05, indicating rejection. This pattern reflects the relative conservatism of the Taylor series approximation, since rejection under the Taylor series approximation implies rejection under the separable approach, but not vice versa.




Figure 3: Sensitivity analysis for the sharp null of no individual treatment effect for all units. The top panel shows upper one-sided p-values under the separable approximation across values of Γ, with the horizontal line at 0.05; the bottom panel reports, across the same values of Γ, the corresponding Taylor series margin, with the zero line marking the rejection boundary. The shaded vertical band highlights values of Γ for which the separable approximation and the Taylor series approximation lead to different rejection decisions.

Both approaches to sensitivity analysis rely on multiple approximations. For example, the separable approximation is used to approximate the worst-case configuration of hidden bias $\boldsymbol{u} \in \mathcal{U}_+$, and inference

additionally relies on a Normal approximation to the null randomization distribution of the test statistic. If the exact randomization distribution were used instead, the resulting sensitivity conclusions could differ for certain values of Γ. In principle, one could also assess sensitivity at a given Γ using the exact randomization distribution. For each configuration of hidden bias $\boldsymbol{u} \in \mathcal{U}_+$, one would derive from (3) the implied probability distribution on Ω, combine that distribution with the previously defined `exact_sharp_null_dist` to compute the corresponding p-value, and take the largest of these p-values as the worst-case p-value (Rosenbaum and Krieger, 1990). A single configuration, such as the separable choice, gives only a lower bound on the worst-case p-value. Our study is small enough to compute the p-values for all 1,152 configurations in $\mathcal{U}_+$. The exact worst-case p-value first exceeds 0.05 at $\Gamma \approx 1.12$, compared with about 1.14 under the Normal approximation, so with only 12 matched sets the Normal approximation slightly overstates the robustness of our conclusion.

2.4.2 How Do Inferences under the Weak Framework Change under These Departures?

Compared with tests of the sharp null hypothesis of no effect, constructing valid tests of an analogous weak null hypothesis is more challenging. A sharp null hypothesis specifies all missing potential outcomes, so we need only consider p-values over configurations of $\boldsymbol{u} \in \mathcal{U}_+$, rather than also over multiple configurations of potential outcomes. By contrast, a weak null permits many possible configurations of potential outcomes, so identifying the worst-case p-value requires optimization over both $\boldsymbol{u}$ and the potential outcomes consistent with the null. With binary outcomes, this joint optimization can be solved by integer programming under a Normal approximation (Fogarty et al., 2017). In matched-pair designs, the joint optimization is unnecessary because the worst-case expectation computed under the sharp null also bounds the expectation under the weak null, so a studentized sensitivity analysis is valid for the weak null in large samples (Fogarty, 2020a).

Fogarty (2023) proposes a more generally applicable approach, which avoids the need for an explicit search for the worst-case p-value over all configurations of $\boldsymbol{u}$ and potential outcomes consistent with the null hypothesis about τ. To build intuition for this approach, note that the following inverse probability weighted (IPW) version of the Difference-in-Means is unbiased for the ATE, τ:

$$\sum_{s=1}^{S} (n_s/n) \left(\frac{1}{|\Omega_s|} \right) \left(\frac{\hat{\tau}_s}{p(\boldsymbol{z}_s)} \right), \tag{5}$$

where $p(\boldsymbol{z}_s)$ denotes the probability of assignment $\boldsymbol{z}_s$ among all treatment assignments in set s that hold fixed the realized number of treated units, with $|\Omega_s|$ denoting, as defined earlier, the total number of such assignments. Both $\hat{\tau}_s$ and $p(\boldsymbol{z}_s)$ are evaluated at the same realized treatment assignment $\boldsymbol{z}_s$ within matched set s. In other words, $p(\boldsymbol{z}_s)$ denotes the probability of the assignment under which the treated and control outcomes in set s — and hence the Difference-in-Means $\hat{\tau}_s$ — are realized.

When $\Gamma = 1$, this probability $p(\boldsymbol{z}_s)$ is $1/|\Omega_s|$, and the IPW Difference-in-Means reduces to the usual Difference-in-Means, $\hat{\tau} = \sum_{s=1}^{S}(n_s/n)\hat{\tau}_s$. When $\Gamma > 1$, however, $p(\boldsymbol{z}_s)$ is no longer known exactly. In matched designs in which every set has either a single treated unit and one or more controls or a single control unit and one or more treated units, Γ restricts the assignment probability to lie within bounds consistent with the specified degree of departure from as-if randomization:

$$\frac{1}{\Gamma(n_s-1)+1} \leq p(\boldsymbol{z}_s) \leq \frac{\Gamma}{(n_s-1)+\Gamma}, \tag{6}$$

for all $\boldsymbol{z}_s \in \Omega_s$ and for all matched sets s.

Since the IPW Difference-in-Means in (5) cannot be directly computed when $\Gamma > 1$, Fogarty (2023) instead uses the bounds in (6) to construct a worst-case version of this IPW Difference-in-Means. For a test of the weak null hypothesis $H_0\colon \tau = \tau_0$ against the alternative of a larger ATE, the worst-case IPW Difference-in-Means is

$$T_\Gamma(\tau_0) = \sum_{s=1}^{S}(n_s/n)\left(\frac{1}{|\Omega_s|}\right)\left(\frac{\hat{\tau}_s - \tau_0}{\tilde{p}_s}\right), \quad \text{where} \quad \tilde{p}_s = \begin{cases} \dfrac{\Gamma}{(n_s-1)+\Gamma} & \text{if } \hat{\tau}_s \geq \tau_0, \\ \dfrac{1}{\Gamma(n_s-1)+1} & \text{if } \hat{\tau}_s < \tau_0. \end{cases} \tag{7}$$

That is, the statistic replaces the unknown probability $p(\boldsymbol{z}_s)$ in (5) with whichever bound in (6) works against rejecting the null. Each set's Difference-in-Means is first centered at the null ATE. When the centered value $\hat{\tau}_s - \tau_0$ is positive, and so counts as evidence against the null in favor of a larger ATE, dividing the centered value by the largest probability that Γ allows makes the set's contribution as small as possible. When the centered value is negative, dividing it by the smallest probability makes the set's contribution as negative as possible. When testing the null against the alternative of a smaller ATE, the two substitutions are reversed. At $\Gamma = 1$, both bounds equal $1/|\Omega_s|$, and $T_\Gamma(\tau_0)$ reduces to $\hat{\tau} - \tau_0$.

Technically, Fogarty (2023, Section 4.1) works with a rescaled version of this statistic, $\tilde{K}_{\Gamma}^{(\tau_0)}$, which is exactly twice $T_{\Gamma}(\tau_0)$. The rescaling alone does not change the test, because the estimated standard error of $\tilde{K}_{\Gamma}^{(\tau_0)}$, computed with the same version of the variance estimator, is exactly twice that of $T_{\Gamma}(\tau_0)$. The worst-case IPW interpretation comes from the preprint version of Fogarty (2023). The preprint describes each matched set's quantity $(1/|\Omega_s|)(\hat{\tau}_s - \tau_0)/\tilde{p}_s$ in (7) as a worst-case inverse probability weighted estimator (Fogarty, 2020b, Sections 5.2–5.4).

To see the value of this procedure, consider testing a null hypothesis about the ATE against the alternative of a larger ATE. The construction of $T_{\Gamma}(\tau_0)$ has two features. First, each set's Difference-in-Means is centered at τ_0. Second, each set uses whichever bound in (6) works against rejection. As a consequence, Fogarty (2023, Section 4.1 and Lemma 3) shows that, for any $\Gamma \geq 1$, the expected value of this worst-case IPW Difference-in-Means — defined under whatever the true distribution on Ω consistent with that Γ happens to be — is always less than or equal to zero when the null is true. Conversely, when testing against the alternative of a smaller ATE, this expectation is greater than or equal to zero when the null is true. Importantly, these properties hold for all possible values of $\boldsymbol{u}$ and all configurations of potential outcomes that are consistent with the null hypothesis.

How does this property of the expected value ensure a valid test in sufficiently large studies? Focusing on the case of testing against a larger ATE, the intuition is straightforward. If the expected value of the worst-case IPW Difference-in-Means is less than or equal to zero when the null is true, then the worst-case IPW Difference-in-Means is centered at or below zero rather than above it. In other words, the procedure intentionally "tilts" the worst-case IPW Difference-in-Means in the direction opposite the alternative, making the procedure conservative.

In addition to this property of the expected value, a second key ingredient for valid hypothesis testing concerns variance estimation. For any fixed $\Gamma \geq 1$, an analogue of the variance estimator from Fogarty (2018) (discussed in Section 2.3.2) is conservative for the true variance of the worst-case IPW Difference-in-Means in two respects. The variance estimator's expected value is at least as large as the true variance, and, as the number of matched sets grows, the variance estimator's ratio to the true variance converges in probability to a value of at least 1 (Fogarty, 2023, Lemma 2 and p. 2202). As a result, in large studies, using this variance estimator tends to reduce the magnitude of the standardized test statistic — the worst-case IPW Difference-in-Means, which is already centered at the null ATE, divided by the square

root of the estimated variance — relative to what would be obtained using the true (but unknown) variance.

Together, these two properties work in the same direction:

- the expected value is at or below zero, keeping the center of the distribution at zero or shifted away from the upper tail, and
- the variance estimator tends to overestimate the true variance, which pulls the standardized statistic closer to zero.

As a result, in large studies, when the null is true, the probability that the standardized statistic falls in the upper tail of the standard Normal distribution — that is, the probability of falsely rejecting the null — remains at or below the nominal significance level. Hence, the test is conservative (valid) asymptotically, as the number of matched sets grows, under the regularity conditions of Fogarty (2023, Theorem 2). Analogous reasoning applies when testing against the alternative of a smaller effect. When the null hypothesis is true, the probability that the standardized statistic falls in the lower tail of the standard Normal distribution also stays at or below the nominal significance level. With few matched sets, as in our application with 12, the Normal approximation may be inaccurate, the variance estimator may be imprecise, and the guarantees above, which are asymptotic, need not hold. The test can then over-reject, so readers should regard the resulting p-values as approximate.

To implement this approach from Fogarty (2023), we first source from the GitHub repository an `R` function that, for a fixed $\Gamma \geq 1$, computes the quantity $(1/|\Omega_s|)(\hat{\tau}_s - \tau_0)/\tilde{p}_s$ in (7) for a single matched set s.

```
# Load the worst_case_IPW() function from GitHub repository

source(
  paste0(
    base_url, "/R/worst_case_IPW.R"
  )
)
```

We now use this function to calculate the worst-case IPW version of the Difference-in-Means within each matched set. To form the overall statistic $T_\Gamma(\tau_0)$ in (7), we average the set-specific values using weights equal to each set's share of the total number of units, n_s/n. Because `worst_case_IPW()` already centers

each set's Difference-in-Means at the null ATE before applying the bounds in (6) on the assignment probability, no further centering is needed. Finally, we standardize this statistic using the conservative variance estimator of Fogarty (2018) from Section 2.3.2, and compare the resulting standardized test statistic to the standard Normal distribution to obtain p-values. For the rescaled statistic $\tilde{K}_{\Gamma}^{(\tau_0)}$, Fogarty (2023, p. 2202) uses a slightly different version of this variance estimator, one that depends on Γ as well as on set sizes. Both versions are conservative (Fogarty, 2018, Proposition 1), and on our data the two give nearly identical sensitivity values, that is, smallest values of Γ at which the test no longer rejects (1.1756 and 1.1753).

```
# One-sided sensitivity analysis for the weak null (ATE = 0)

# Null and alternative for the weak-null test
null_ATE    <- 0                    # Null: ATE = 0

alternative <- "greater"            # Alt: ATE > 0 (upper-tail test)

# Data frame to store results for each Gamma
weak_sens_df <- data.frame(
  Gamma   = Gamma_vals,
  p_value = NA_real_
)

for (g in Gamma_vals) { # Loop over Gamma values

  # Worst-case IPW estimate within each matched set at Gamma = g
  set_stats <- data_matched |>
    group_by(
      fm
    ) |>
    summarize(
      n      = n(),                        # Set size n_s
      weight = n / nrow(data_matched),     # Weight n_s / n
      est    = worst_case_IPW(             # Worst-case IPW in set s
        z           = UN,                  # Treatment indicator
        y           = ldur,                # Outcome
        Gamma       = g,                   # Sensitivity parameter (Gamma)
        tau_h       = null_ATE,            # Weak null: ATE = 0
        alternative = alternative          # Direction of alternative
      ),
      .groups = "drop"
    )

  # Overall worst-case IPW statistic (weighted average across sets)
```

```
  wc_ipw_stat <- sum(
    set_stats$weight * set_stats$est
  )

  # Variance estimate for the worst-case IPW statistic
  var_hat <- fine_strat_var_est(
    strat_ns   = set_stats$n,
    strat_ests = set_stats$est
  )

  # Standardized worst-case IPW statistic, centered at the null
  se_hat   <- sqrt(
    var_hat
  )

  std_stat <- wc_ipw_stat / se_hat

  # One-sided p-value, determined by the alternative
  if (alternative == "greater") {
    # Upper-tail test: ATE > 0
    p_val <- 1 - pnorm(std_stat)
  } else if (alternative == "less") {
    # Lower-tail test: ATE < 0
    p_val <- pnorm(std_stat)
  } else {
    stop("Only one-sided alternatives ('greater' or 'less').")
  }

  # Store result for this Gamma
  weak_sens_df$p_value[weak_sens_df$Gamma == g] <- p_val
}

# Sensitivity value: smallest Gamma where one-sided test no longer rejects
weak_sens_value <- min(
  weak_sens_df$Gamma[weak_sens_df$p_value >= alpha]
)
```

Figure 4 plots the upper one-sided p-values corresponding to Γ values from 1 to 1.5. Under as-if randomization ($\Gamma = 1$), we find evidence of a positive average effect; the p-value at $\Gamma = 1$ in Figure 4 is 0.036, rather than the 0.037 reported in Section 2.3.2, because the sensitivity analysis estimates the variance with the estimator of Fogarty (2018) instead of `hybrid_p`. This rejection of the null hypothesis of no average effect in favor of a positive average effect is sensitive to departures from as-if randomization. Our qualitative conclusion about the null hypothesis that the ATE equals zero changes at $\Gamma \approx 1.18$ (1.1756). This sensitivity value is similar to the value of about 1.14 for the sharp null of no effect under

the Normal approximation, but the two analyses use different test statistics, variance calculations, and approximations.




Figure 4: Upper one-sided p-values for the weak null hypothesis of zero average effect across values of the sensitivity parameter Γ; the horizontal line marks 0.05.

This sensitivity analysis guarantees (in large studies) validity across *all* configurations of potential outcomes consistent with the weak null. The price of this guarantee is conservativeness. If effects are in fact constant and $\Gamma > 1$, the test's limiting probability of rejection is strictly below α, except in matched-pair designs (Fogarty, 2023, Theorem 2). By contrast, Fogarty (2023) also describes, and recommends, an alternative sensitivity analysis that is asymptotically sharp when effects are constant. The validity of this alternative analysis for the weak null is guaranteed under an additional condition that jointly restricts the potential outcomes and the pattern of hidden bias, a condition that researchers may regard as plausible (Fogarty, 2023, Theorem 1, Condition 1, and Section 8). The alternative sensitivity analysis can produce smaller p-values, but at the cost of forfeiting validity for certain (possibly unrealistic) combinations of potential outcomes and hidden bias.

3 Conclusion

We have now completed the full matching pipeline. We began by constructing matched sets and then conducted inference under the as-if randomization assumption, treating the matched design as a completely randomized experiment within blocks. We examined inference under both the sharp framework, which concerns the effect for every unit, and the weak framework, which concerns the average effect. We then assessed the sensitivity of these inferences to potential violations of the as-if randomization assumption. The embedded code, accompanied by extensive comments and illustrated through the running example from Gilligan and Sergenti (2008), is intended to be readily adapted by practitioners to their own datasets.

3.1 Limitations and Related Topics Beyond Our Scope

For clarity and focus, we have excluded several important topics, though we have pointed to relevant references where appropriate. Some of these omissions are not specific to matching or observational studies. In particular, we have not addressed imperfect compliance with treatment assignment, missing outcomes, clustered (rather than individual-level) assignment, or interference settings in which a subject's outcome depends on others' treatment assignments.

Our analysis has focused on inference for two causal targets only: a constant, additive treatment effect and the average treatment effect. We have not considered alternative effect models, such as dilated effects (Rosenbaum, 1999), multiplicative effects, or Tobit effects (Rosenbaum, 2010, pp. 46–48). Throughout, we have treated causal targets as fixed quantities (over the set of treatment assignments consistent with the matched design), rather than as random variables. As a result, we have excluded attributable effects (Rosenbaum, 2001, 2003; Hansen and Bowers, 2009) and the average treatment effect on the treated (Sekhon and Shem-Tov, 2021). We have also set aside methods for joint inference on sharp and weak causal hypotheses (Wu and Ding, 2021; Cohen and Fogarty, 2022; see also Ding, 2017), as well as strategies that improve the power of hypothesis tests by rescaling outcomes or choosing alternative test statistics, including regression-assisted approaches (Lin, 2013; Cohen and Fogarty, 2024; Guo and Basse, 2023).

Several omissions are specific to matching methodology itself:

- First, we have not covered a range of **alternative matching and balancing approaches**, including nonbipartite matching for multivalued treatments (Lu et al., 2001, 2011; Rosenbaum, 2010, pp. 207–221), template matching (Silber et al., 2014), multilevel matching (Zubizarreta and Keele, 2017; Pimentel et al., 2018), risk-set matching (Li et al., 2001), cardinality matching (Zubizarreta et al., 2014), coarsened exact matching (Iacus et al., 2012), exterior matching (Rosenbaum and Silber, 2013), or generalized full matching (Sävje et al., 2021), among others. We have further assumed that researchers specify *ex ante* the set of covariates to be balanced, thereby excluding screening procedures that identify additional covariates for adjustment based on observed imbalances (Cochran, 1965), as in Yu et al. (2021), who use a tapered match (Daniel et al., 2008). Finally, we have not considered balance constraints such as fine and near-fine balance (Rosenbaum et al., 2007; Yang et al., 2012; Yu, 2023; Pimentel et al., 2015a,b), which enforce exact or nearly exact equality of the marginal distributions of selected nominal covariates (categorical covariates with unordered categories).
- Second, we have not discussed the use of **prognostic covariates** identified using pilot or external data. Using such covariates can substantially improve efficiency by prioritizing covariates that strongly predict potential outcomes. For theoretical foundations, see Hansen (2008a), with complementary theoretical results and practical insights in Sales et al. (2018).
- Third, we have considered only matching strategies aimed at justifying as-if randomization, not those designed explicitly to improve **design sensitivity** — that is, robustness to moderate violations of as-if randomization (see Rosenbaum, 2010, Part III, pp. 257–311). Beyond match construction, researchers can also enhance design sensitivity by selecting particular test statistics, among other strategies (Rosenbaum, 2004; Heller et al., 2009; Hsu et al., 2013; Small et al., 2013).
- Fourth, we have excluded **extensions to covariate balance testing** beyond the framework developed in Hansen and Bowers (2008), including subsequent contributions by Gagnon-Bartsch and Shem-Tov (2019) and Branson (2021), as well as related approaches based on the stepwise intersection–union principle (Hansen and Sales, 2015).
- Fifth, we have not directly addressed **residual imbalance on observed covariates**. Specifically, we have proceeded under the assumption that matched-set membership can justify as-if randomization, even when small imbalances remain within matched sets. Under this approach, concerns about residual imbalance are absorbed into the sensitivity analysis for hidden bias. This

approach is not entirely satisfying, however, because imbalances on observed covariates are not, in fact, hidden. Several methods therefore address such imbalances directly, either in approaches targeting individual effects (Rosenbaum, 1988; Pimentel and Huang, 2024; Chen et al., 2023; Heng et al., 2025) or average effects (Zhu et al., 2025).

- Sixth, we have not incorporated into the overall pipeline an explicit approach for defining an **interpretable study population** in matched designs. Common practice excludes units based on limited overlap, whether in the estimated propensity score (Crump et al., 2009) or in the covariate space itself (King and Zeng, 2006), but study populations defined in this way can be difficult to characterize substantively. Hence, alternative approaches define the study population directly in terms of a small number of covariates. For example, Traskin and Small (2011) use a classification tree to define a study population that approximates one defined by trimming on the estimated propensity score, while Fogarty et al. (2016) use discrete optimization to define a maximal, rectangular region of covariate space with adequate overlap on key covariates.
- Seventh, we have not considered **quasi-experimental devices** (Campbell, 1957; Campbell and Stanley, 1963; Shadish et al., 2002), specifically how matched designs can use such devices (see Rosenbaum, 2025, Part IV). Two prominent quasi-experimental devices are an additional control group (Rosenbaum, 1987b) and outcomes with known effects (Rosenbaum, 1989), each of which has a range of valuable applications. In the context of evidence factors (Rosenbaum, 2021), one use of a second control group is to yield stronger causal evidence that is less susceptible to hidden bias (Rosenbaum, 2023a). For known effects, one illustrative use is to support tests for hidden bias that inform sensitivity analyses by ruling out certain violations of as-if randomization (Rosenbaum, 2023b).
- Eighth and finally, we have not addressed ***Z*-dependence** (Pashley et al., 2021; Pimentel and Huang, 2024; Pimentel and Yu, 2026). Our exposition has relied on an analogy to Rubin's framework of "assignment to treatment group on the basis of a covariate" (Rubin, 1977), where matched-set membership plays the role of the covariate. This analogy treats set membership as fixed, but because matching uses the realized treatment assignment, other assignments, including those in Ω that permute treatment within the realized sets, can produce different matched sets. In our application, permuting treatment within any single matched set and re-running our matching code produces a different matched structure. Strictly, assignments that would have produced a different matched

structure do not belong in the reference distribution. Pashley et al. (2021, Section 5) show that exact matching on the covariates that determine treatment justifies the usual permutation analysis by a conditioning argument and that approximate matching does not. Pashley et al. (2021, p. 276) add, however, that this distinction may matter more in theory than in practice. In the simulations of Pimentel and Huang (2024) and Pimentel and Yu (2026), permutation tests that treated the matched sets as fixed showed excess Type I error, which propensity score calipers reduced but did not always remove. We thank Samuel Pimentel for helpful discussions of $\boldsymbol{Z}$-dependence.

Despite these limitations, the pipeline and design-based perspective developed here provide a coherent framework for understanding how matched designs support causal inference. The emphasis on design-based foundations clarifies the logic underlying the construction of matched sets and the subsequent steps of inference and sensitivity analysis under both sharp and weak frameworks. We present this work not as exhaustive, but as a foundation on which researchers can build when incorporating more specialized methods tailored to their substantive and inferential goals.

References

Donald B. Rubin. Assignment to treatment group on the basis of a covariate. *Journal of Educational Statistics*, 2(1):1–26, 1977.

Ben B. Hansen. Full matching in an observational study of coaching for the SAT. *Journal of the American Statistical Association*, 99(467):609–618, 2004.

Ben B. Hansen and Stephanie Olsen Klopfer. Optimal full matching and related designs via network flows. *Journal of Computational and Graphical Statistics*, 15(3):609–627, 2006.

Ben B. Hansen and Jake Bowers. Covariate balance in simple, stratified and clustered comparative studies. *Statistical Science*, 23(2):219–236, 2008.

Colin B. Fogarty. On mitigating the analytical limitations of finely stratified experiments. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(5):1035–1056, 2018.

Nicole E. Pashley and Luke W. Miratrix. Insights on variance estimation for blocked and matched pairs designs. *Journal of Educational and Behavioral Statistics*, 46(3):271–296, 2021.

Xing Sam Gu and Paul R. Rosenbaum. Comparison of multivariate matching methods: Structures, distances, and algorithms. *Journal of Computational and Graphical Statistics*, 2(4):405–420, 1993.

Paul R. Rosenbaum. A characterization of optimal designs for observational studies. *Journal of the Royal Statistical Society. Series B (Methodological)*, 53(3):597–610, 1991.

Ben B. Hansen. Propensity score matching to extract latent experiments from nonexperimental data: A case study. In Neil J. Dorans and Sandip Sinharay, editors, *Looking Back: Proceedings of a Conference in Honor of Paul W. Holland*, volume 202 of *Lecture Notes in Statistics*, chapter 9, pages 149–181. Springer, New York, NY, 2011.

Paul R. Rosenbaum and Abba M. Krieger. Sensitivity of two-sample permutation inferences in observational studies. *Journal of the American Statistical Association*, 85(410):493–498, 1990.

Joseph L. Gastwirth, Abba M. Krieger, and Paul R. Rosenbaum. Asymptotic separability in sensitivity analysis. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 62(3):545–555, 2000.

Paul R. Rosenbaum. Sensitivity analysis for stratified comparisons in an observational study of the effect of smoking on homocysteine levels. *Annals of Applied Statistics*, 12(4):2312–2334, 2018.

Colin B. Fogarty. Testing weak nulls in matched observational studies. *Biometrics*, 79(3):2196–2207, 2023.

Michael J. Gilligan and Ernest J. Sergenti. Do UN interventions cause peace? Using matching to improve causal inference. *Quarterly Journal of Political Science*, 3(2):89–122, 2008.

Donald B. Rubin and Richard P. Waterman. Estimating the causal effects of marketing interventions using propensity score methodology. *Statistical Science*, 21(2):206–222, 2006.

Ben B. Hansen. The prognostic analogue of the propensity score. *Biometrika*, 95(2):481–488, 2008a.

Adam C. Sales, Ben B. Hansen, and Brian Rowan. Rebar: Reinforcing a matching estimator with predictions from high-dimensional covariates. *Journal of Educational and Behavioral Statistics*, 43(1): 3–31, 2018.

Prasanta C. Mahalanobis. On the generalized distance in statistics. *Proceedings of the National Institute of Sciences of India*, 2(1):49–55, 1936.

A. Albert and J.A. Anderson. On the existence of maximum likelihood estimates in logistic regression models. *Biometrika*, 71(1):1–10, 1984.

Ben B. Hansen and Adam Sales. Comment on Cochran's "Observational Studies". *Observational Studies*, 1(1):184–193, 2015.

Paul R. Rosenbaum. *Design of Observational Studies*. Springer, New York, NY, 1st edition, 2010.

William G. Cochran and Donald B. Rubin. Controlling bias in observational studies: A review. *Sankhyā: The Indian Journal of Statistics, Series A*, 35(4):417–446, 1973.

Sture Holm. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(2):65–70, 1979.

Peter C. Austin. A critical appraisal of propensity-score matching in the medical literature between 1996 and 2003. *Statistics in Medicine*, 27(12):2037–2049, 2008.

Kosuke Imai, Gary King, and Elizabeth A. Stuart. Misunderstandings between experimentalists and

observationalists about causal inference. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 171(2):481–502, 2008.

Ben B. Hansen. The essential role of balance tests in propensity-matched observational studies: Comments on 'A critical appraisal of propensity-score matching in the medical literature between 1996 and 2003' by Peter Austin. *Statistics in Medicine*, 27(12):2050–2054, 2008b.

Peter C. Austin. Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples. *Statistics in Medicine*, 28(25):3083–3107, 2009.

Elizabeth A. Stuart. Matching methods for causal inference: A review and a look forward. *Statistical Science*, 25(1):1–21, 2010.

Donald B. Rubin. Comment on 'Randomization analysis of experimental data: The Fisher randomization test' by D. Basu. *Journal of the American Statistical Association*, 75(371):591–593, 1980.

Donald B. Rubin. Which ifs have causal answers? (Comment on 'Statistics and causal inference' by Paul W. Holland). *Journal of the American Statistical Association*, 81(396):961–962, 1986.

David R. Cox. *Planning of Experiments*. Wiley, New York, NY, 1958.

Paul R. Rosenbaum and Donald B. Rubin. The bias due to incomplete matching. *Biometrics*, 41(1): 103–116, 1985.

Paul R. Rosenbaum. Optimal matching of an optimally chosen subset in observational studies. *Journal of Computational and Graphical Statistics*, 21(1):57–71, 2012.

Colin B. Fogarty, Mark E. Mikkelsen, David F. Gaieski, and Dylan S. Small. Discrete optimization for interpretable study populations and randomization inference in an observational study of severe sepsis mortality. *Journal of the American Statistical Association*, 111(514):447–458, 2016.

Mikhail Traskin and Dylan S. Small. Defining the study population for an observational study to ensure sufficient overlap: A tree approach. *Statistics in Biosciences*, 3(1):94–118, 2011.

Jason Wu and Peng Ding. Randomization tests for weak null hypotheses in randomized experiments. *Journal of the American Statistical Association*, 116(536):1898–1913, 2021.

Peter L. Cohen and Colin B. Fogarty. Gaussian prepivoting for finite population causal inference. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 84(2):295–320, 2022.

Eun Yi Chung and Joseph P. Romano. Exact and asymptotically robust permutation tests. *The Annals of Statistics*, 41(2):484–507, 2013.

Peng Ding. A paradox from randomization-based causal inference. *Statistical Science*, 32(3):331–345, 2017.

Colin B. Fogarty. Studentized sensitivity analysis for the sample average treatment effect in paired observational studies. *Journal of the American Statistical Association*, 115(531):1518–1530, 2020a.

Andrew Gelman. Causality and statistical learning. *American Journal of Sociology*, 117(3):955–966, 2011.

Devin Caughey, Allan Dafoe, Xinran Li, and Luke Miratrix. Randomisation inference beyond the sharp null: Bounded null hypotheses and quantiles of individual treatment effects. *Journal of the Royal Statistical Society Series B (Statistical Methodology)*, 85(5):1471–1491, 2023.

Nicole E. Pashley, Guillaume W. Basse, and Luke W. Miratrix. Conditional as-if analyses in randomized experiments. *Journal of Causal Inference*, 9(1):264–284, 2021.

Paul R. Rosenbaum. *Observation and Experiment: An Introduction to Causal Inference*. Harvard University Press, Cambridge, MA, 2017.

Samuel D. Pimentel and Yaxuan Huang. Covariate-adaptive randomization inference in matched designs. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 86(5):1312–1338, 2024.

Samuel D. Pimentel and Ruoqi Yu. Re-evaluating the impact of hormone replacement therapy on heart disease using match-adaptive randomization inference. *Journal of the American Statistical Association*, 2026. doi: 10.1080/01621459.2026.2691321. Published online.

Ronald A. Fisher. *The Design of Experiments*. Oliver and Boyd, Edinburgh, 1935.

Fan Li, Kari Lock Morgan, and Alan M. Zaslavsky. Balancing covariates via propensity score weighting. *Journal of the American Statistical Association*, 113(521):390–400, 2018.

Joseph L. Hodges, Jr. and Erich L. Lehmann. Estimates of location based on rank tests. *The Annals of Mathematical Statistics*, 34(2):598–611, 1963.

Paul R. Rosenbaum. Hodges–Lehmann point estimates of treatment effect in observational studies. *Journal of the American Statistical Association*, 88(424):1250–1253, 1993.

Jerzy Neyman. Sur les applications de la théorie des probabilités aux expériences agricoles: Essai des principes. *Roczniki Nauk Rolniczych*, 10:1–51, 1923.

James M. Robins. Confidence intervals for causal parameters. *Statistics in Medicine*, 7(7):773–785, 1988.

Peter M. Aronow, Donald P. Green, and Donald K.K. Lee. Sharp bounds on the variance in randomized experiments. *The Annals of Statistics*, 42(3):850–871, 2014.

Christopher Harshaw, Joel A. Middleton, and Fredrik Sävje. Optimized variance estimation under interference and complex experimental designs. *Journal of the American Statistical Association*, 2026. doi: 10.1080/01621459.2026.2627027. Published online.

Luke W. Miratrix, Jasjeet S. Sekhon, and Bin Yu. Adjusting treatment effect estimates by post-stratification in randomized experiments. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 75(2):369–396, 2013.

Paul R. Rosenbaum. Sensitivity analysis for certain permutation inferences in matched observational studies. *Biometrika*, 74(1):13–26, 1987a.

Paul R. Rosenbaum. Conditional permutation tests and the propensity score in observational studies. *Journal of the American Statistical Association*, 79(387):565–574, 1984.

Paul R. Rosenbaum. Quantiles in nonrandom samples and observational studies. *Journal of the American Statistical Association*, 90(432):1424–1431, 1995.

Colin B. Fogarty, Pixu Shi, Mark E. Mikkelsen, and Dylan S. Small. Randomization inference and sensitivity analysis for composite null hypotheses with binary outcomes in matched observational studies. *Journal of the American Statistical Association*, 112(517):321–331, 2017.

Colin B. Fogarty. Testing weak nulls in matched observational studies. arXiv Preprint, https://arxiv.org/abs/1908.07352v3, February 2020b.

Paul R. Rosenbaum. Reduced sensitivity to hidden bias at upper quantiles in observational studies with dilated treatment effects. *Biometrics*, 55(2):560–564, 1999.

Paul R. Rosenbaum. Effects attributable to treatment: Inference in experiments and observational studies with a discrete pivot. *Biometrika*, 88(1):219–231, 2001.

Paul R. Rosenbaum. Exact confidence intervals for nonconstant effects by inverting the signed rank test. *The American Statistician*, 57(2):132–138, 2003.

Ben B. Hansen and Jake Bowers. Attributing effects to a cluster-randomized get-out-the-vote campaign. *Journal of the American Statistical Association*, 104(487):873–885, 2009.

Jasjeet S. Sekhon and Yotam Shem-Tov. Inference on a new class of sample average treatment effects. *Journal of the American Statistical Association*, 116(534):798–804, 2021.

Winston Lin. Agnostic notes on regression adjustments to experimental data: Reexamining Freedman's critique. *The Annals of Applied Statistics*, 7(1):295–318, 2013.

Peter L. Cohen and Colin B. Fogarty. No-harm calibration for generalized Oaxaca-Blinder estimators. *Biometrika*, 111(1):331–338, 2024.

Kevin Guo and Guillaume W. Basse. The generalized Oaxaca-Blinder estimator. *Journal of the American Statistical Association*, 118(541):524–536, 2023.

Bo Lu, Elaine Zanutto, Robert Hornik, and Paul R. Rosenbaum. Matching with doses in an observational study of a media campaign against drug abuse. *Journal of the American Statistical Association*, 96 (456):1245–1253, 2001.

Bo Lu, Robert Greevy, Xinyi Xu, and Cole Beck. Optimal nonbipartite matching and its statistical applications. *The American Statistician*, 65(1):21–30, 2011.

Jeffrey H. Silber, Paul R. Rosenbaum, Richard N. Ross, Justin M. Ludwig, Wei Wang, Bijan A. Niknam, Nabanita Mukherjee, Philip A. Saynisch, Orit Even-Shoshan, Rachel R. Kelz, and Lee A. Fleisher. Template matching for auditing hospital cost and quality. *Health Services Research*, 49(5):1446–1474, 2014.

José R. Zubizarreta and Luke Keele. Optimal multilevel matching in clustered observational studies: A

case study of the effectiveness of private schools under a large-scale voucher system. *Journal of the American Statistical Association*, 112(518):547–560, 2017.

Samuel D. Pimentel, Lindsay C. Page, Matthew Lenard, and Luke Keele. Optimal multilevel matching using network flows: An application to a summer reading intervention. *The Annals of Applied Statistics*, 12(3):1479–1505, 2018.

Yunfei Paul Li, Kathleen J. Propert, and Paul R. Rosenbaum. Balanced risk set matching. *Journal of the American Statistical Association*, 96(455):870–882, 2001.

José R. Zubizarreta, Ricardo D. Paredes, and Paul R. Rosenbaum. Matching for balance, pairing for heterogeneity in an observational study of the effectiveness of for-profit and not-for-profit high schools in Chile. *The Annals of Applied Statistics*, 8(1):204–231, 2014.

Stefano M. Iacus, Gary King, and Giuseppe Porro. Causal inference without balance checking: Coarsened exact matching. *Political Analysis*, 20(1):1–24, 2012.

Paul R. Rosenbaum and Jeffrey H. Silber. Using the exterior match to compare two entwined matched control groups. *The American Statistician*, 67(2):67–75, 2013.

Fredrik Sävje, Michael J. Higgins, and Jasjeet S. Sekhon. Generalized full matching. *Political Analysis*, 29(4):423–447, 2021.

William G. Cochran. The planning of observational studies of human populations. *Journal of the Royal Statistical Society. Series A (General)*, 128(2):234–266, 1965.

Ruoqi Yu, Dylan S. Small, and Paul R. Rosenbaum. The information in covariate imbalance in studies of hormone replacement therapy. *The Annals of Applied Statistics*, 15(4):2023–2042, 2021.

Shoshana R. Daniel, Katrina Armstrong, Jeffrey H. Silber, and Paul R. Rosenbaum. An algorithm for optimal tapered matching, with application to disparities in survival. *Journal of Computational and Graphical Statistics*, 17(4):914–924, 2008.

Paul R. Rosenbaum, Richard N. Ross, and Jeffrey H. Silber. Minimum distance matched sampling with fine balance in an observational study of treatment for ovarian cancer. *Journal of the American Statistical Association*, 102(477):75–83, 2007.

Dan Yang, Dylan S. Small, Jeffrey H. Silber, and Paul R. Rosenbaum. Optimal matching with minimal deviation from fine balance in a study of obesity and surgical outcomes. *Biometrics*, 68(2):628–636, 2012.

Ruoqi Yu. How well can fine balance work for covariate balancing. *Biometrics*, 79(3):2346–2356, 2023.

Samuel D. Pimentel, Rachel R. Kelz, Jeffrey H. Silber, and Paul R. Rosenbaum. Large, sparse optimal matching with refined covariate balance in an observational study of the health outcomes produced by new surgeons. *Journal of the American Statistical Association*, 110(510):515–527, 2015a.

Samuel D. Pimentel, Frank Yoon, and Luke Keele. Variable-ratio matching with fine balance in a study of the Peer Health Exchange. *Statistics in Medicine*, 34(30):4070–4082, 2015b.

Paul R. Rosenbaum. Design sensitivity in observational studies. *Biometrika*, 91(1):153–164, 2004.

Ruth Heller, Paul R. Rosenbaum, and Dylan S. Small. Split samples and design sensitivity in observational studies. *Journal of the American Statistical Association*, 104(487):1090–1101, 2009.

Jesse Y. Hsu, Dylan S. Small, and Paul R. Rosenbaum. Effect modification and design sensitivity in observational studies. *Journal of the American Statistical Association*, 108(501):135–148, 2013.

Dylan S. Small, Jing Cheng, M. Elizabeth Halloran, and Paul R. Rosenbaum. Case definition and design sensitivity. *Journal of the American Statistical Association*, 108(504):1457–1468, 2013.

Johann Gagnon-Bartsch and Yotam Shem-Tov. The classification permutation test: A flexible approach to testing for covariate imbalance in observational studies. *The Annals of Applied Statistics*, 13(3): 1464–1483, 2019.

Zach Branson. Randomization tests to assess covariate balance when designing and analyzing matched datasets. *Observational Studies*, 7(2):1–36, 2021.

Paul R. Rosenbaum. Permutation tests for matched pairs with adjustments for covariates. *Journal of the Royal Statistical Society Series C: Applied Statistics*, 37(3):401–411, 1988.

Kan Chen, Siyu Heng, Qi Long, and Bo Zhang. Testing biased randomization assumptions and quantifying imperfect matching and residual confounding in matched observational studies. *Journal of Computational and Graphical Statistics*, 32(2):528–538, 2023.

Siyu Heng, Yanxin Shen, and Pengyun Wang. Reconciling overt bias and hidden bias in sensitivity analysis for matched observational studies. arXiv Preprint, https://arxiv.org/pdf/2311.11216v4, May 2025.

Jianan Zhu, Jeffrey Zhang, Zijian Guo, and Siyu Heng. Randomization-based inference for average treatment effects in inexactly matched observational studies. arXiv Preprint, https://arxiv.org/pdf/2308.02005v4, March 2025.

Richard K. Crump, V. Joseph Hotz, Guido W. Imbens, and Oscar A. Mitnik. Dealing with limited overlap in estimation of average treatment effects. *Biometrika*, 96(1):187–199, 2009.

Gary King and Langche Zeng. The dangers of extreme counterfactuals. *Political Analysis*, 14(2):131–159, 2006.

Donald T. Campbell. Factors relevant to the validity of experiments in social settings. *Psychological Bulletin*, 54(4):297–312, 1957.

Donald T. Campbell and Julian C. Stanley. *Experimental and Quasi-Experimental Designs for Research*. Houghton Mifflin Company, Boston, MA, 1963.

William R. Shadish, Thomas D. Cook, and Donald T. Campbell. *Experimental and Quasi-Experimental Designs for Generalized Causal Inference*. Houghton Mifflin Company, Boston, MA, 2002.

Paul R. Rosenbaum. *An Introduction to the Theory of Observational Studies*. Springer Texts in Statistics. Springer, Cham, Switzerland, 2025.

Paul R. Rosenbaum. The role of a second control group in an observational study. *Statistical Science*, 2 (3):292–306, 1987b.

Paul R. Rosenbaum. The role of known effects in observational studies. *Biometrics*, 45(2):557–569, 1989.

Paul R. Rosenbaum. *Replication and Evidence Factors in Observational Studies*. Number 167 in Monographs on Statistics and Applied Probability. CRC Press, Boca Raton, FL, 2021.

Paul R. Rosenbaum. A second evidence factor for a second control group. *Biometrics*, 79(4):3968–3980, 2023a.

Paul R. Rosenbaum. Sensitivity analyses informed by tests for bias in observational studies. *Biometrics*, 79(1):475–487, 2023b.