

# Variance of the Difference-in-Means Estimator for the SATE and Estimation of That Variance\*

Thomas Leavitt

This guide derives the variance of the Difference-in-Means estimator for the Sample Average Treatment Effect (SATE) under complete random assignment, and then develops a conservative estimator of that variance. This guide states the framework only as far as its derivations require and shares that framework with three companion guides: “Notation and Setup for Design-Based Causal Inference” develops the notation in full, “Unbiasedness of the Difference-in-Means Estimator for the SATE and the PATE” establishes that the estimator is unbiased, and “Variance of the Difference-in-Means Estimator for the PATE and Estimation of That Variance” extends the results below to a superpopulation, where the estimand is the Population Average Treatment Effect (PATE).

## 1 Variance of the Difference-in-Means estimator for the Sample Average Treatment Effect (SATE)

### 1.1 Setup

Let the index  $i \in \{1, \dots, n\}$  run over the  $n$  units of a finite sample,  $\mathcal{S}_n$ , of which  $n_T \geq 2$  receive treatment and  $n_C = n - n_T \geq 2$  receive control, so that  $n \geq 4$ ; these bounds ensure that the estimator of the variance developed below, which forms a separate sample variance within each arm, is well defined. The indicator  $Z_i \in \{0, 1\}$  records whether unit  $i$  receives treatment ( $Z_i = 1$ ) or control ( $Z_i = 0$ ), and the random assignment vector  $\mathbf{Z} = [Z_1, \dots, Z_n]^\top$  collects the indicators.

---

\*This is a live document that is subject to updating at any time.

Write  $n_T(\mathbf{z}) := \sum_{i=1}^n z_i$  for the number of treated units under an assignment  $\mathbf{z}$ , so that  $n_T(\mathbf{Z})$  is a random variable. Under *complete random assignment* the design fixes the count of treated units: The support of  $\mathbf{Z}$  is  $\Omega = \{\mathbf{z} \in \{0, 1\}^n : n_T(\mathbf{z}) = n_T\}$ , on which  $n_T(\mathbf{Z})$  equals the design constant  $n_T$  with probability one, and  $\mathbf{Z}$  is distributed uniformly on  $\Omega$ , whose cardinality<sup>1</sup> is  $|\Omega| = \binom{n}{n_T}$ . The unbiasedness guide shows how the same analysis covers simple random assignment after conditioning on the event  $\{n_T(\mathbf{Z}) = n_T\}$ .

Under the Stable Unit Treatment Value Assumption (SUTVA) (Cox, 1958; Rubin, 1980, 1986), unit  $i$ 's potential outcome depends on the assignment vector only through unit  $i$ 's own coordinate, so each unit has two *fixed* potential outcomes,  $y_i(1)$  and  $y_i(0)$ , with individual effect  $\tau_i := y_i(1) - y_i(0)$ . The observable outcome is  $Y_i := y_i(Z_i) = Z_i y_i(1) + (1 - Z_i) y_i(0)$ , random only through  $Z_i$ . The setup guide develops the potential outcomes schedule, states SUTVA precisely, and walks through the collapse from  $y_i(\mathbf{Z})$  to  $y_i(1)$  and  $y_i(0)$  in numbered steps.

Write  $\bar{y}(1) := n^{-1} \sum_{i=1}^n y_i(1)$  and  $\bar{y}(0) := n^{-1} \sum_{i=1}^n y_i(0)$  for the finite population means of the treatment and control potential outcomes; both are fixed quantities. The target of interest is the Sample Average Treatment Effect (SATE),

$$\tau_{\text{SATE}} := n^{-1} \sum_{i=1}^n \tau_i = \bar{y}(1) - \bar{y}(0),$$

which is likewise a fixed (though unknown) quantity. Define the Difference-in-Means estimator of  $\tau_{\text{SATE}}$  under complete random assignment as

$$(1) \quad \hat{\tau} := \frac{1}{n_T} \sum_{i=1}^n Z_i Y_i - \frac{1}{n_C} \sum_{i=1}^n (1 - Z_i) Y_i.$$

The estimator in Equation (1) is the same Difference-in-Means estimator whose unbiasedness the companion guide establishes. Under complete random assignment, where the design fixes the counts  $n_T(\mathbf{Z}) = n_T$  and  $n_C(\mathbf{Z}) = n_C$ , the estimator coincides with the random-denominator form  $(\sum_i Z_i Y_i) / (\sum_i Z_i) - (\sum_i (1 - Z_i) Y_i) / (\sum_i (1 - Z_i))$  that the setup guide gives. Unlike the estimand  $\tau_{\text{SATE}}$ , the estimator  $\hat{\tau}$  is a *random variable*, since  $\hat{\tau}$  is a function of the random assignment vector  $\mathbf{Z}$  and inherits all randomness from  $\mathbf{Z}$ . For the expectation and variance of  $\hat{\tau}$  in Equation (1), I write  $E_\Omega[\cdot]$  and  $\text{Var}_\Omega[\cdot]$  to emphasize that they run over only the randomness of the assignment process, i.e., over the distribution of  $\mathbf{Z}$  on  $\Omega$ .

---

<sup>1</sup>For an arbitrary set  $\mathcal{W}$ , let  $|\mathcal{W}|$  denote the cardinality of (i.e., the number of elements in) the set  $\mathcal{W}$ .

## 1.2 Derivation of variance of Difference-in-Means estimator for the SATE

**Proposition 1.** *The variance of  $\hat{\tau}$  for the  $\tau_{\text{SATE}}$  under complete random assignment is*

$$(2) \quad \text{Var}_{\Omega}[\hat{\tau}] = \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} - \frac{S_n^2(\boldsymbol{\tau})}{n},$$

where

$$(3) \quad S_n^2(\mathbf{y}(1)) = \left( \frac{1}{n-1} \right) \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right)^2$$

$$(4) \quad S_n^2(\mathbf{y}(0)) = \left( \frac{1}{n-1} \right) \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right)^2$$

$$(5) \quad S_n^2(\boldsymbol{\tau}) = \left( \frac{1}{n-1} \right) \sum_{i=1}^n \left( \tau_i - \frac{1}{n} \sum_{i=1}^n \tau_i \right)^2.$$

*Proof.* Building on [Imbens and Rubin \(2015, Chapter 6, Appendix A\)](#), I will break the proof into several steps:

**Step 1:** I will show that we can rewrite the Difference-in-Means estimator in (1) not in terms of the raw assignment indicators  $Z_i$ , but in terms of the centered treatment variable  $A_i := Z_i - \text{E}_{\Omega}[Z_i]$ . I center because the variance measures the fluctuations of  $\hat{\tau}$  around its mean,  $\tau_{\text{SATE}}$ . If we can algebraically separate  $\hat{\tau}$  into a fixed part (equal to  $\tau_{\text{SATE}}$ ) plus a random part with mean zero, then the variance of  $\hat{\tau}$  equals the variance of the random part alone. Centering the assignment variable produces this separation and, because  $\text{E}_{\Omega}[A_i] = 0$ , greatly simplifies the algebra that follows.

**Step 2:** The potential outcomes,  $\{y_i(1), y_i(0)\}_{i=1}^n$ , are fixed, and the observable outcomes inherit their randomness only from the treatment assignment. Once I write  $\hat{\tau}$  as a weighted sum of the  $A_i$ , the variance of  $\hat{\tau}$  depends only on the first and second moments of the  $A_i$ . I will therefore derive  $\text{E}_{\Omega}[A_i]$ ,  $\text{Var}_{\Omega}[A_i]$  and  $\text{Cov}_{\Omega}[A_i, A_j]$  for  $i \neq j$ . The covariance term is essential and nonzero. Under complete random assignment the design fixes the count of treated units at  $n_T$ , so if one unit is pulled into treatment another must be pulled out. This mechanical dependence among the assignment indicators is the source of the covariance term, and the covariance turns out to be negative.

**Step 3:** I will rely on the expressions for  $S_n^2(\mathbf{y}(1))$  and  $S_n^2(\mathbf{y}(0))$  above, as well as on the following algebraic equivalence that I will need to prove:

$$(6) \quad S_n^2(\boldsymbol{\tau}) = S_n^2(\mathbf{y}(1)) + S_n^2(\mathbf{y}(0)) - \frac{2}{n-1} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right).$$

**Step 4:** Once I have completed the three previous steps, I will rely only on the linearity of expectations, rules of variance and (often very messy) algebra.

### Step 1

First, we can rewrite the estimator as

$$\frac{1}{n_T} \sum_{i=1}^n Z_i Y_i - \frac{1}{n_C} \sum_{i=1}^n (1 - Z_i) Y_i = \frac{1}{n} \sum_{i=1}^n \left( \frac{n}{n_T} Z_i y_i(1) - \frac{n}{n_C} (1 - Z_i) y_i(0) \right)$$

Because  $n = n_C + n_T$ ,  $1 = \frac{n_C + n_T}{n}$ , so instead of  $(1 - Z_i)$  we can write  $(\frac{n_C + n_T}{n} - Z_i)$ :

$$\frac{1}{n} \sum_{i=1}^n \frac{n}{n_T} Z_i y_i(1) - \frac{n}{n_C} \left( \frac{n_C + n_T}{n} - Z_i \right) y_i(0),$$

which after a little bit of algebra yields

$$(7) \quad \frac{1}{n} \sum_{i=1}^n \frac{n}{n_T} \left( \underbrace{Z_i - \frac{n_T}{n}}_{=A_i} + \frac{n_T}{n} \right) y_i(1) - \frac{n}{n_C} \left[ \frac{n_C}{n} - \left( \underbrace{Z_i - \frac{n_T}{n}}_{=A_i} \right) \right] y_i(0),$$

where  $A_i$  is the centered treatment variable,  $A_i = Z_i - E_\Omega[Z_i]$ , because, under complete random assignment,  $E_\Omega[Z_i] = \frac{n_T}{n}$ .

Therefore, we can rewrite (7) as

$$\frac{1}{n} \sum_{i=1}^n \frac{n}{n_T} \left( A_i + \frac{n_T}{n} \right) y_i(1) - \frac{n}{n_C} \left( \frac{n_C}{n} - A_i \right) y_i(0),$$

which, with a bit more algebra, is equivalent to

$$\underbrace{\frac{1}{n} \sum_{i=1}^n (y_i(1) - y_i(0))}_{= \tau_{\text{SATE}}} + \frac{1}{n} \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) = \tau_{\text{SATE}} + \frac{1}{n} \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right)$$

## Step 2

Thus far, I have shown that the Difference-in-Means estimator in (1) is equivalent to

$$(8) \quad \tau_{\text{SATE}} + \frac{1}{n} \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right),$$

in which  $\tau_{\text{SATE}}$ ,  $n_C$ ,  $n_T$ ,  $n$ , and  $\{y_i(0), y_i(1)\}_{i=1}^n$  are all fixed quantities. The only random quantities in (8) are  $\{A_i\}_{i=1}^n$ .

Therefore, to derive the variance of (8), I now need to derive  $E_\Omega[A_i]$ ,  $\text{Var}_\Omega[A_i]$  and  $E_\Omega[A_i A_j]$  for  $i \neq j$ .

To do so, I write the two values that  $A_i$  takes:

$$(9) \quad A_i = Z_i - \frac{n_T}{n} = \begin{cases} \frac{n_C}{n} & \text{if } Z_i = 1 \\ -\frac{n_T}{n} & \text{if } Z_i = 0 \end{cases}$$

and recall that, under complete random assignment,  $\Pr_\Omega(Z_i = 1) = \frac{n_T}{n}$  and  $\Pr_\Omega(Z_i = 0) = 1 - \frac{n_T}{n} = \frac{n}{n} - \frac{n_T}{n} = \frac{n - n_T}{n} = \frac{n_C}{n}$ .

**Derivation of  $E_\Omega[A_i]$ :** Substituting the definition of  $A_i$  gives  $E_\Omega[A_i] = E_\Omega[Z_i - \frac{n_T}{n}] = E_\Omega[Z_i] - \frac{n_T}{n} = \frac{n_T}{n} - \frac{n_T}{n} = 0$  or, equivalently,

$$\begin{aligned} E_\Omega[A_i] &= \frac{n_C}{n} \Pr_\Omega(Z_i = 1) - \frac{n_T}{n} \Pr_\Omega(Z_i = 0) \\ &= \frac{n_C}{n} \frac{n_T}{n} - \frac{n_T}{n} \left( 1 - \frac{n_T}{n} \right) \\ &= \frac{n_C}{n} \frac{n_T}{n} - \frac{n_T}{n} \frac{n_C}{n} \\ &= 0. \end{aligned}$$

**Derivation of  $\text{Var}_\Omega[A_i]$ :** Since  $E_\Omega[A_i] = 0$ , the variance reduces to  $\text{Var}_\Omega[A_i] = E_\Omega[A_i^2] - E_\Omega[A_i]^2 = E_\Omega[A_i^2]$ . Thus,

$$\begin{aligned} \text{Var}_\Omega[A_i] &= E_\Omega[A_i^2] \\ &= \left( \frac{n_C}{n} \right)^2 \Pr_\Omega(Z_i = 1) + \left( -\frac{n_T}{n} \right)^2 \Pr_\Omega(Z_i = 0) \\ &= \frac{n_C^2}{n^2} \Pr_\Omega(Z_i = 1) + \frac{n_T^2}{n^2} \Pr_\Omega(Z_i = 0) \end{aligned}$$

$$\begin{aligned}
&= \frac{n_C^2}{n^2} \frac{n_T}{n} + \frac{n_T^2}{n^2} \frac{n_C}{n} \\
&= \frac{n_C^2 n_T}{n^3} + \frac{n_T^2 n_C}{n^3} \\
&= \frac{n_C^2 n_T + n_T^2 n_C}{n^3} \\
&= \frac{n_C n_T (n_C + n_T)}{n^3} \\
&= \frac{n_C n_T (n)}{n^3} \\
&= \frac{n_C n_T}{n^2}.
\end{aligned}$$

**Derivation of  $\text{Cov}_\Omega[A_i, A_j]$  for  $i \neq j$ :** The  $\{A_i\}_{i=1}^n$  all have the same expected value,  $E_\Omega[A_i] = 0$  (as I just showed), but they are *not* independent: Fixing the count  $n_T(\mathbf{Z})$  at  $n_T$  links the assignment indicators to one another. I therefore will need to derive  $\text{Cov}_\Omega[A_i, A_j]$  for  $i \neq j$ . To do so, first note that

$$\begin{aligned}
\text{Cov}_\Omega [A_i, A_j] &= E_\Omega \left[ (A_i - E_\Omega [A_i]) (A_j - E_\Omega [A_j]) \right] \\
&= E_\Omega \left[ (A_i - 0) (A_j - 0) \right] \\
&= E_\Omega [A_i A_j],
\end{aligned}$$

which is easier to work with.

The possible values that  $A_i A_j$  could take on are

$$A_i A_j = \begin{cases} \left( \frac{n_C}{n} \right) \left( \frac{n_C}{n} \right) = \frac{n_C^2}{n^2} & \text{if } Z_i = 1 \text{ and } Z_j = 1 \\ \left( -\frac{n_T}{n} \right) \left( \frac{n_C}{n} \right) = \frac{-n_T n_C}{n^2} & \text{if } Z_i = 0 \text{ and } Z_j = 1 \text{ or } Z_i = 1 \text{ and } Z_j = 0 \\ \left( -\frac{n_T}{n} \right) \left( -\frac{n_T}{n} \right) = \frac{n_T^2}{n^2} & \text{if } Z_i = 0 \text{ and } Z_j = 0. \end{cases}$$

Having derived the values that  $A_i A_j$  takes, I now need to derive the probabilities of the four assignment events that produce those values, namely,  $\Pr_\Omega(Z_i = 1, Z_j = 1)$ ,  $\Pr_\Omega(Z_i = 0, Z_j = 1)$ ,  $\Pr_\Omega(Z_i = 1, Z_j = 0)$  and  $\Pr_\Omega(Z_i = 0, Z_j = 0)$ .

Remember that  $Z_i$  and  $Z_j$  are *not* independent, which implies that, e.g.,  $\Pr_\Omega(Z_i = 1, Z_j = 1) \neq \Pr_\Omega(Z_i = 1) \Pr_\Omega(Z_j = 1)$ . Instead, I will appeal to the definition of joint probability in which

$$\Pr_{\Omega}(Z_i = z, Z_j = z') = \Pr_{\Omega}(Z_i = z) \Pr_{\Omega}(Z_j = z' \mid Z_i = z):$$

$$\Pr_{\Omega}(Z_i = z, Z_j = z') = \begin{cases} \left(\frac{n_T}{n}\right) \left(\frac{n_T - 1}{n - 1}\right) & \text{if } Z_i = 1 \text{ and } Z_j = 1 \\ \left(\frac{n_T}{n}\right) \left(\frac{n_C}{n - 1}\right) & \text{if } Z_i = 1 \text{ and } Z_j = 0 \\ \left(\frac{n_C}{n}\right) \left(\frac{n_T}{n - 1}\right) & \text{if } Z_i = 0 \text{ and } Z_j = 1 \\ \left(\frac{n_C}{n}\right) \left(\frac{n_C}{n - 1}\right) & \text{if } Z_i = 0 \text{ and } Z_j = 0. \end{cases}$$

We can therefore write the probability mass function of  $A_i A_j$  as

$$\Pr_{\Omega}(A_i A_j) = \begin{cases} \left(\frac{n_T}{n}\right) \left(\frac{n_T - 1}{n - 1}\right) = \frac{n_T (n_T - 1)}{n (n - 1)} & \text{if } A_i A_j = \frac{n_C^2}{n^2} \\ \left(\frac{n_T}{n}\right) \left(\frac{n_C}{n - 1}\right) + \left(\frac{n_C}{n}\right) \left(\frac{n_T}{n - 1}\right) = \frac{2 (n_T n_C)}{n (n - 1)} & \text{if } A_i A_j = \frac{-n_T n_C}{n^2} \\ \left(\frac{n_C}{n}\right) \left(\frac{n_C}{n - 1}\right) = \frac{n_C (n_C - 1)}{n (n - 1)} & \text{if } A_i A_j = \frac{n_T^2}{n^2}, \end{cases}$$

which implies that

$$\mathbb{E}_{\Omega}[A_i A_j] = \frac{n_C^2}{n^2} \left( \frac{n_T (n_T - 1)}{n (n - 1)} \right) + \frac{-n_T n_C}{n^2} \left( \frac{2 (n_T n_C)}{n (n - 1)} \right) + \frac{n_T^2}{n^2} \left( \frac{n_C (n_C - 1)}{n (n - 1)} \right)$$

or equivalently, after some algebra,

$$\frac{-n_T n_C}{n^2 (n - 1)}.$$

### Step 3

In this step, I will prove the algebraic equivalence between the expression for  $S_n^2(\boldsymbol{\tau})$  in Equation (5) above and

$$(10) \quad S_n^2(\boldsymbol{\tau}) = S_n^2(\mathbf{y}(1)) + S_n^2(\mathbf{y}(0)) - \frac{2}{n-1} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right),$$

which will be valuable for the derivation of  $\text{Var}_{\Omega}[\hat{\tau}]$  in **Step 4** to follow.

Recall from Equation (5) that

$$S_n^2(\boldsymbol{\tau}) = \frac{1}{n-1} \sum_{i=1}^n \left( \tau_i - \frac{1}{n} \sum_{i=1}^n \tau_i \right)^2.$$

Hence,

$$\begin{aligned}
S_n^2(\boldsymbol{\tau}) &= \frac{1}{n-1} \sum_{i=1}^n \left( \tau_i - \frac{1}{n} \sum_{i=1}^n \tau_i \right)^2 \\
&= \frac{1}{n-1} \sum_{i=1}^n \left( \underbrace{(y_i(1) - y_i(0))}_{=\tau_i} - \frac{1}{n} \sum_{i=1}^n \underbrace{(y_i(1) - y_i(0))}_{=\tau_i} \right)^2 \\
&= \frac{1}{n-1} \sum_{i=1}^n \left( \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) - \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \right)^2,
\end{aligned}$$

where, in the last line, I have used  $\frac{1}{n} \sum_{i=1}^n (y_i(1) - y_i(0)) = \frac{1}{n} \sum_{i=1}^n y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(0)$  to regroup each summand into a treated deviation,  $(y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1))$ , minus a control deviation,  $(y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0))$ . Expanding the square via  $(a - b)^2 = a^2 - 2ab + b^2$  and then distributing the sum yields

$$\begin{aligned}
S_n^2(\boldsymbol{\tau}) &= \underbrace{\frac{1}{n-1} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right)^2}_{=S_n^2(\mathbf{y}(1))} + \underbrace{\frac{1}{n-1} \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right)^2}_{=S_n^2(\mathbf{y}(0))} \\
&\quad - \frac{2}{n-1} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \\
&= S_n^2(\mathbf{y}(1)) + S_n^2(\mathbf{y}(0)) - \frac{2}{n-1} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right).
\end{aligned}$$

#### Step 4

To complete the derivation, I compute the variance of the Difference-in-Means estimator from the expression in (8):

$$\begin{aligned}
\text{Var}_\Omega [\hat{\tau}] &= \text{Var}_\Omega \left[ \tau_{\text{SATE}} + \frac{1}{n} \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) \right] \\
&= \frac{1}{n^2} \text{Var}_\Omega \left[ \sum_{i=1}^n A_i \underbrace{\left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right)}_{\text{constant}} \right],
\end{aligned}$$



which, since  $E_\Omega[A_i] = 0$ , is equivalent to

$$\begin{aligned} \text{Var}_\Omega[\hat{\tau}] &= \frac{1}{n^2} \left( E_\Omega \left[ \left( \sum_{i=1}^n A_i \underbrace{\left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right)}_{\text{constant}} \right)^2 \right] - E_\Omega \left[ \sum_{i=1}^n A_i \underbrace{\left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right)}_{\text{constant}} \right]^2 \right) \\ &= \frac{1}{n^2} E_\Omega \left[ \left( \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) \right)^2 \right]. \end{aligned}$$

Expanding the expression immediately above yields

$$\begin{aligned} \text{Var}_\Omega[\hat{\tau}] &= \frac{1}{n^2} E_\Omega \left[ \left( \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) \right)^2 \right] \\ &= \frac{1}{n^2} E_\Omega \left[ \left( \sum_{i=1}^n A_i \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) \right) \left( \sum_{j=1}^n A_j \left( \frac{n}{n_T} y_j(1) + \frac{n}{n_C} y_j(0) \right) \right) \right] \\ &= \frac{1}{n^2} E_\Omega \left[ \sum_{i=1}^n \sum_{j=1}^n A_i A_j \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) \left( \frac{n}{n_T} y_j(1) + \frac{n}{n_C} y_j(0) \right) \right]. \end{aligned}$$

In **Step 2** above, I showed that

$$\begin{cases} E_\Omega[A_i A_j] = \frac{-n_T n_C}{n^2(n-1)} & \text{when } i \neq j \\ E_\Omega[A_i A_j] = E_\Omega[A_i^2] = \text{Var}_\Omega[A_i] = \frac{n_C n_T}{n^2} & \text{when } i = j. \end{cases}$$

I now split the double sum  $\sum_{i=1}^n \sum_{j=1}^n$  into the  $n$  diagonal terms (where  $i = j$ , so that  $A_i A_j = A_i^2$ ) and the  $n(n-1)$  off-diagonal terms (where  $i \neq j$ ), because  $E_\Omega[A_i A_j]$  takes a different value in each case. The split yields

$$\begin{aligned} \text{Var}_\Omega[\hat{\tau}] &= \frac{1}{n^2} \left( \sum_{i=1}^n \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right)^2 \underbrace{E_\Omega[A_i^2]}_{=\frac{n_C n_T}{n^2}} \right. \\ &\quad \left. + \sum_{i=1}^n \sum_{j \neq i} \left( \frac{n}{n_T} y_i(1) + \frac{n}{n_C} y_i(0) \right) \left( \frac{n}{n_T} y_j(1) + \frac{n}{n_C} y_j(0) \right) \underbrace{E_\Omega[A_i A_j]}_{=\frac{-n_T n_C}{n^2(n-1)}} \right). \end{aligned}$$

I now substitute the expressions I derived for  $E_\Omega[A_i^2]$  and  $E_\Omega[A_i A_j]$ . To keep the bookkeeping

manageable, I abbreviate the constant weight multiplying  $A_i$  as  $w_i := \frac{n}{n_T}y_i(1) + \frac{n}{n_C}y_i(0)$ , so that the expression above becomes

$$\text{Var}_\Omega [\hat{\tau}] = \frac{1}{n^2} \left( \frac{n_C n_T}{n^2} \sum_{i=1}^n w_i^2 - \frac{n_T n_C}{n^2 (n-1)} \sum_{i=1}^n \sum_{j \neq i} w_i w_j \right).$$

Using the identity  $\sum_{i=1}^n \sum_{j \neq i} w_i w_j = (\sum_{i=1}^n w_i)^2 - \sum_{i=1}^n w_i^2$  and then collecting terms over the common denominator  $n-1$ ,

$$\text{Var}_\Omega [\hat{\tau}] = \frac{n_C n_T}{n^4 (n-1)} \left( n \sum_{i=1}^n w_i^2 - \left( \sum_{i=1}^n w_i \right)^2 \right) = \frac{n_C n_T}{n^3 (n-1)} \sum_{i=1}^n (w_i - \bar{w})^2,$$

where  $\bar{w} = \frac{1}{n} \sum_{i=1}^n w_i$ , and where the final equality uses the identity  $n \sum_i w_i^2 - (\sum_i w_i)^2 = n \sum_i (w_i - \bar{w})^2$ . Finally, I substitute back the centered weight,

$$w_i - \bar{w} = \frac{n}{n_T} \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) + \frac{n}{n_C} \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right),$$

and expanding the square yields

$$\begin{aligned} \text{Var}_\Omega [\hat{\tau}] &= \frac{n_C}{n n_T (n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right)^2 + \frac{n_T}{n n_C (n-1)} \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right)^2 \\ &+ \frac{2}{n(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right). \end{aligned} \quad (11)$$

Now recall that

$$\begin{aligned} S_n^2(\mathbf{y}(1)) &= \frac{1}{n-1} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right)^2 \quad \text{and} \\ S_n^2(\mathbf{y}(0)) &= \frac{1}{n-1} \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right)^2, \end{aligned}$$

which yields

$$\text{Var}_\Omega [\hat{\tau}] = \frac{n_C}{n n_T} S_n^2(\mathbf{y}(1)) + \frac{n_T}{n n_C} S_n^2(\mathbf{y}(0)) + \frac{2}{n(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right).$$

It then follows that

$$\begin{aligned}
\text{Var}_\Omega [\hat{\tau}] &= \frac{n_C}{nn_T} S_n^2(\mathbf{y}(1)) + \frac{n_T}{nn_C} S_n^2(\mathbf{y}(0)) + \frac{2}{n(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \\
&= \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} - \frac{S_n^2(\mathbf{y}(1))}{n} - \frac{S_n^2(\mathbf{y}(0))}{n} \\
&\quad + \frac{2}{n(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \\
&= \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} \\
&\quad - \underbrace{\frac{1}{n} \left( S_n^2(\mathbf{y}(1)) + S_n^2(\mathbf{y}(0)) - \frac{2}{(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \right)}_{=S_n^2(\boldsymbol{\tau})}.
\end{aligned}$$

As I showed in **Step 3**, the bracketed term in the expression immediately above is equal to  $S_n^2(\boldsymbol{\tau})$ , which leaves

$$(12) \quad \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} - \frac{S_n^2(\boldsymbol{\tau})}{n}.$$

□

Each of the three terms in this variance has a clear meaning. The estimator  $\hat{\tau}$  is the difference between the treated group mean and the control group mean, so its variability comes from two pieces. The first term,  $\frac{S_n^2(\mathbf{y}(1))}{n_T}$ , is the variability that the treated mean contributes. The first term grows with the spread of the treatment potential outcomes,  $S_n^2(\mathbf{y}(1))$ , and shrinks as the number of treated units,  $n_T$ , grows. The second term,  $\frac{S_n^2(\mathbf{y}(0))}{n_C}$ , is the analogous variability that the control mean contributes. If the treated and control means came from two *independently* drawn groups, these first two terms would account for the entire variance.

The two means are *not* independent, however, and the third term,  $-\frac{S_n^2(\boldsymbol{\tau})}{n}$ , corrects for the dependence that complete random assignment induces. The treated and control groups are complementary, since every unit the design sends to treatment is a unit it *withholds* from control, so the treated mean and the control mean are negatively coupled. The coupling runs through the individual treatment effects, which is why the correction involves  $S_n^2(\boldsymbol{\tau})$ , the finite population variance of the individual effects  $\tau_i = y_i(1) - y_i(0)$ . Two consequences follow. First, the correction enters with a minus sign and therefore *lowers* the variance; ignoring the correction would overstate the variance, which is what makes the conservative estimator of the variance (developed in the next section) conservative rather than anti-conservative. Second, when the treatment effect is constant across

units,  $\tau_i$  is the same for all  $i$ , so  $S_n^2(\boldsymbol{\tau}) = 0$ , the correction vanishes, and the spreads of the two sets of potential outcomes fully determine the variance.

The expression I just derived for  $\text{Var}_\Omega[\hat{\tau}]$  is the *true* variance of the Difference-in-Means estimator. That expression is mathematically equivalent to Equation 3.4 in Gerber and Green (2012, 57), which differs from the expression in Equation (12) above because the variances and covariance of treated and control potential outcomes in Gerber and Green (2012, Equation 3.4, 57) use a denominator of  $n$  as opposed to  $n - 1$  as in Equations (3) to (5) above. The corollary below establishes this equivalence.

**Corollary 1.** *An equivalent expression for the finite sample variance of the Difference-in-Means estimator under complete random assignment is*

$$(13) \quad \text{Var}_\Omega[\hat{\tau}] = \frac{1}{n-1} \left( \frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \frac{n_T \sigma_n^2(\mathbf{y}(0))}{n_C} + 2\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) \right),$$

where

$$(14) \quad \sigma_n^2(\mathbf{y}(1)) = \left( \frac{1}{n} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \right)^2$$

$$(15) \quad \sigma_n^2(\mathbf{y}(0)) = \left( \frac{1}{n} \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \right)^2$$

$$(16) \quad \sigma_n(\mathbf{y}(0), \mathbf{y}(1)) = \left( \frac{1}{n} \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right) \right) \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right).$$

*Proof.* Recall the expression for  $\text{Var}_\Omega[\hat{\tau}]$  in Equation (11):

$$\begin{aligned} \text{Var}_\Omega[\hat{\tau}] &= \frac{n_C}{nn_T(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right)^2 + \frac{n_T}{nn_C(n-1)} \sum_{i=1}^n \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right)^2 \\ &\quad + \frac{2}{n(n-1)} \sum_{i=1}^n \left( y_i(1) - \frac{1}{n} \sum_{i=1}^n y_i(1) \right) \left( y_i(0) - \frac{1}{n} \sum_{i=1}^n y_i(0) \right). \end{aligned}$$

Then the definitions of  $\sigma_n^2(\mathbf{y}(1))$ ,  $\sigma_n^2(\mathbf{y}(0))$  and  $\sigma_n(\mathbf{y}(0), \mathbf{y}(1))$  above yield

$$\text{Var}_\Omega[\hat{\tau}] = \frac{n_C}{n_T(n-1)} \sigma_n^2(\mathbf{y}(1)) + \frac{n_T}{n_C(n-1)} \sigma_n^2(\mathbf{y}(0)) + \frac{2}{(n-1)} \sigma_n(\mathbf{y}(0), \mathbf{y}(1))$$

$$= \frac{1}{n-1} \left( \frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \frac{n_T \sigma_n^2(\mathbf{y}(0))}{n_C} + 2\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) \right),$$

which completes the proof.  $\square$

## 2 Conservative estimation of the variance

Having derived the true variance  $\text{Var}_\Omega[\hat{\tau}]$ , I now turn to estimating it from the observed data. Absent additional assumptions, we *cannot* estimate  $\text{Var}_\Omega[\hat{\tau}]$  unbiasedly. The obstruction is the same in both equivalent expressions for the variance. Each expression depends on a quantity that couples a unit's two potential outcomes, namely the covariance  $\sigma_n(\mathbf{y}(0), \mathbf{y}(1))$  in the Gerber–Green form or the variance of the individual effects,  $S_n^2(\boldsymbol{\tau})$ , in the Imbens–Rubin form, and we never observe both potential outcomes for any single unit (the fundamental problem of causal inference). What we *can* do is construct an estimable quantity that is at least as large as the true variance and estimate that quantity instead. The resulting estimator is *conservative*, in the sense that on average it weakly overstates the variance, which yields valid confidence intervals (possibly too wide) rather than overconfident ones.

I establish the upper bound in two ways, starting from each of the two equivalent expressions for the variance. Both routes arrive at the *same* conservative estimator. I first record the two expressions for reference. The Imbens–Rubin form (Proposition 1) is

$$(17) \quad \text{Var}_\Omega[\hat{\tau}] = \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} - \frac{S_n^2(\boldsymbol{\tau})}{n},$$

and the Gerber–Green form (Corollary 1) is

$$(18) \quad \text{Var}_\Omega[\hat{\tau}] = \frac{1}{n-1} \left( \frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \frac{n_T \sigma_n^2(\mathbf{y}(0))}{n_C} + 2\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) \right).$$

### 2.1 Approach 1 (Gerber–Green expression): bounding the covariance

In Equation (18), we can estimate the variances  $\sigma_n^2(\mathbf{y}(1))$  and  $\sigma_n^2(\mathbf{y}(0))$  unbiasedly, but we cannot estimate the covariance  $\sigma_n(\mathbf{y}(0), \mathbf{y}(1))$ . The first approach replaces the unestimable covariance with an estimable upper bound. The Cauchy–Schwarz inequality implies that

$$\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) \leq \sqrt{\sigma_n^2(\mathbf{y}(0))\sigma_n^2(\mathbf{y}(1))},$$

and the arithmetic mean–geometric mean (AM–GM) inequality further implies that

$$\sqrt{\sigma_n^2(\mathbf{y}(0))\sigma_n^2(\mathbf{y}(1))} \leq \frac{\sigma_n^2(\mathbf{y}(0)) + \sigma_n^2(\mathbf{y}(1))}{2}.$$

Hence  $2\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) \leq \sigma_n^2(\mathbf{y}(0)) + \sigma_n^2(\mathbf{y}(1))$ . Substituting this bound for the covariance term in Equation (18), and then collecting the two terms in each potential outcome’s variance, yields a simple upper bound. The key simplification is that pairing the cross term with the matching squared term gives, for example,

$$\frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \sigma_n^2(\mathbf{y}(1)) = \frac{n_C + n_T}{n_T} \sigma_n^2(\mathbf{y}(1)) = \frac{n \sigma_n^2(\mathbf{y}(1))}{n_T},$$

and likewise for the control variance. Carrying out that substitution,

$$\begin{aligned} \text{Var}_\Omega [\hat{\tau}] &= \frac{1}{n-1} \left( \frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \frac{n_T \sigma_n^2(\mathbf{y}(0))}{n_C} + 2\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) \right) \\ &\leq \frac{1}{n-1} \left( \frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \frac{n_T \sigma_n^2(\mathbf{y}(0))}{n_C} + \sigma_n^2(\mathbf{y}(1)) + \sigma_n^2(\mathbf{y}(0)) \right) \\ &= \frac{1}{n-1} \left( \underbrace{\frac{n_C \sigma_n^2(\mathbf{y}(1))}{n_T} + \sigma_n^2(\mathbf{y}(1))}_{= n \sigma_n^2(\mathbf{y}(1))/n_T} + \underbrace{\frac{n_T \sigma_n^2(\mathbf{y}(0))}{n_C} + \sigma_n^2(\mathbf{y}(0))}_{= n \sigma_n^2(\mathbf{y}(0))/n_C} \right) \\ &= \frac{n}{n-1} \left( \frac{\sigma_n^2(\mathbf{y}(1))}{n_T} + \frac{\sigma_n^2(\mathbf{y}(0))}{n_C} \right). \end{aligned}$$

We can estimate the two remaining unknowns,  $\sigma_n^2(\mathbf{y}(1))$  and  $\sigma_n^2(\mathbf{y}(0))$ , unbiasedly. Following Cochran (1977, Theorem 2.4), unbiased estimators are

$$\begin{aligned} \hat{\sigma}_n^2(\mathbf{y}(1)) &= \left( \frac{n-1}{n(n_T-1)} \right) \sum_{i=1}^n Z_i \left( Y_i - \frac{1}{n_T} \sum_{i=1}^n Z_i Y_i \right)^2 \\ \hat{\sigma}_n^2(\mathbf{y}(0)) &= \left( \frac{n-1}{n(n_C-1)} \right) \sum_{i=1}^n (1-Z_i) \left( Y_i - \frac{1}{n_C} \sum_{i=1}^n (1-Z_i) Y_i \right)^2. \end{aligned}$$

These estimators depend only on the *observed* outcomes. The factor  $Z_i$  selects the treated units, for which  $Y_i = y_i(1)$ , and  $(1-Z_i)$  selects the control units, for which  $Y_i = y_i(0)$ . The leading factor  $\frac{n-1}{n}$  converts the ordinary sample variance, which is unbiased for  $S_n^2(\cdot)$ , into an unbiased estimator of  $\sigma_n^2(\cdot)$ , since  $\sigma_n^2(\cdot) = \frac{n-1}{n} S_n^2(\cdot)$ . Substituting these into the upper bound gives the conservative

estimator of the variance

$$(19) \quad \widehat{\text{Var}} [\hat{\tau}] = \frac{n}{n-1} \left( \frac{\hat{\sigma}_n^2(\mathbf{y}(1))}{n_T} + \frac{\hat{\sigma}_n^2(\mathbf{y}(0))}{n_C} \right).$$

## 2.2 Approach 2 (Imbens–Rubin expression): dropping the variance of the individual effects

The second approach starts from Equation (17) and is more direct. The only term we cannot estimate is  $S_n^2(\boldsymbol{\tau})$ , the finite population variance of the individual effects, since computing  $S_n^2(\boldsymbol{\tau})$  would require knowing both potential outcomes for each unit. Notice that  $S_n^2(\boldsymbol{\tau})$  enters with a *minus* sign and is always nonnegative. Therefore

$$\text{Var}_\Omega [\hat{\tau}] = \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} - \underbrace{\frac{S_n^2(\boldsymbol{\tau})}{n}}_{\geq 0} \leq \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C}.$$

That is, simply *dropping* the nonnegative term we cannot estimate produces an upper bound. The two remaining terms depend on one potential outcome per unit, and the corresponding sample variances within each group estimate them unbiasedly,

$$\begin{aligned} \hat{S}_n^2(\mathbf{y}(1)) &= \left( \frac{1}{n_T - 1} \right) \sum_{i=1}^n Z_i \left( Y_i - \frac{1}{n_T} \sum_{i=1}^n Z_i Y_i \right)^2 \\ \hat{S}_n^2(\mathbf{y}(0)) &= \left( \frac{1}{n_C - 1} \right) \sum_{i=1}^n (1 - Z_i) \left( Y_i - \frac{1}{n_C} \sum_{i=1}^n (1 - Z_i) Y_i \right)^2, \end{aligned}$$

which are well defined because  $n_T \geq 2$  and  $n_C \geq 2$ . Substituting these into the upper bound gives the conservative estimator of the variance

$$(20) \quad \widehat{\text{Var}} [\hat{\tau}] = \frac{1}{n_T} \hat{S}_n^2(\mathbf{y}(1)) + \frac{1}{n_C} \hat{S}_n^2(\mathbf{y}(0)).$$

## 2.3 The two approaches give the same estimator, and how conservative it is

The two routes produce the *same* estimator. To see the equivalence, recall that  $\hat{\sigma}_n^2(\mathbf{y}(1)) = \frac{n-1}{n} \hat{S}_n^2(\mathbf{y}(1))$  (compare the two pairs of estimators above), so that

$$\frac{n}{n-1} \frac{\hat{\sigma}_n^2(\mathbf{y}(1))}{n_T} = \frac{n}{n-1} \cdot \frac{n-1}{n} \cdot \frac{\hat{S}_n^2(\mathbf{y}(1))}{n_T} = \frac{\hat{S}_n^2(\mathbf{y}(1))}{n_T},$$

and likewise for the control term. Hence Equation (19) and Equation (20) are identical; bounding the covariance via the Cauchy–Schwarz/AM–GM route and dropping the nonnegative variance of the individual effects are two ways of describing the same bound.

Finally, the conservative estimator overstates the variance by an amount we can derive exactly. Because  $\widehat{S}_n^2(\mathbf{y}(1))$  and  $\widehat{S}_n^2(\mathbf{y}(0))$  are unbiased for  $S_n^2(\mathbf{y}(1))$  and  $S_n^2(\mathbf{y}(0))$ ,

$$\mathbb{E}_\Omega \left[ \widehat{\text{Var}} [\hat{\tau}] \right] = \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} = \text{Var}_\Omega [\hat{\tau}] + \frac{S_n^2(\boldsymbol{\tau})}{n},$$

where the last equality uses Equation (17). Therefore

$$\mathbb{E}_\Omega \left[ \widehat{\text{Var}} [\hat{\tau}] \right] - \text{Var}_\Omega [\hat{\tau}] = \frac{S_n^2(\boldsymbol{\tau})}{n} \geq 0,$$

so the estimator’s upward bias is exactly  $S_n^2(\boldsymbol{\tau})/n$ . The estimator is conservative in general, and it is *exactly* unbiased when the individual treatment effects are constant across units, i.e., when  $S_n^2(\boldsymbol{\tau}) = 0$  (equivalently, when  $2\sigma_n(\mathbf{y}(0), \mathbf{y}(1)) = \sigma_n^2(\mathbf{y}(0)) + \sigma_n^2(\mathbf{y}(1))$ , so that the Cauchy–Schwarz/AM–GM bound in Approach 1 holds with equality).

### 3 Extension: variance and its estimation under simple random assignment

So far the assignment mechanism has been complete random assignment, with the counts of treated and control units fixed at  $n_T$  and  $n_C$  by design. Under *simple random assignment*, each unit receives treatment independently with a common assignment probability, the two degenerate assignments (all treated and all control) are excluded, and the count  $n_T(\mathbf{Z})$  is a nondegenerate random variable with support  $\{1, \dots, n-1\}$ ; the setup and unbiasedness guides develop this mechanism in full. The estimator is the random-denominator form that the setup guide gives, with the counts of treated and control units,  $n_T(\mathbf{Z})$  and  $n_C(\mathbf{Z})$ , in the denominators.

The conditioning argument that the unbiasedness guide uses for the expectation extends to the variance, for one reason: Conditional on the event  $\{n_T(\mathbf{Z}) = n_T\}$ , the distribution of  $\mathbf{Z}$  is the complete random assignment distribution, as the unbiasedness guide proves (see also [Pashley et al., 2021](#)), so every result above holds conditionally, word for word. In particular,

$$\text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z}) = n_T] = \frac{S_n^2(\mathbf{y}(1))}{n_T} + \frac{S_n^2(\mathbf{y}(0))}{n_C} - \frac{S_n^2(\boldsymbol{\tau})}{n},$$

which is Equation (12) evaluated at the realized counts. Two facts then turn conditional statements



into unconditional ones. First, the conditional expectation of the estimator is the *same* constant at every count of treated units: The unbiasedness guide proves  $E_\Omega[\hat{\tau} \mid n_T(\mathbf{Z}) = k] = \tau_{\text{SATE}}$  for every  $k \in \{1, \dots, n-1\}$ . Second, the conservative estimator's conditional overshoot,  $S_n^2(\boldsymbol{\tau})/n$ , also does not depend on the count of treated units. Averaging over that count therefore costs nothing, as the following proposition records.

**Proposition 2.** *Under simple random assignment,*

$$\text{Var}_\Omega [\hat{\tau}] = S_n^2(\mathbf{y}(1)) E_\Omega \left[ \frac{1}{n_T(\mathbf{Z})} \right] + S_n^2(\mathbf{y}(0)) E_\Omega \left[ \frac{1}{n_C(\mathbf{Z})} \right] - \frac{S_n^2(\boldsymbol{\tau})}{n}.$$

*Proof.* I will complete this proof in three steps: I will (1) decompose the variance by conditioning on the count of treated units, (2) evaluate both pieces of the decomposition with results already in hand, and (3) average the conditional variance over the distribution of the count of treated units.

**Step 1:** By the law of total variance, conditioning on the count of treated units,

$$\text{Var}_\Omega [\hat{\tau}] = E_\Omega \left[ \text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \right] + \text{Var}_\Omega \left[ E_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \right].$$

**Step 2:** The second term is zero: The unbiasedness guide proves  $E_\Omega[\hat{\tau} \mid n_T(\mathbf{Z}) = k] = \tau_{\text{SATE}}$  for every  $k$ , and a constant has no variance. For the first term, conditional on  $n_T(\mathbf{Z}) = k$  the distribution of  $\mathbf{Z}$  is complete random assignment with  $k$  treated units — the reduction that the unbiasedness guide's conditional uniformity lemma proves — so Proposition 1 applies with  $n_T = k$  and  $n_C = n - k$ :

$$\text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z}) = k] = \frac{S_n^2(\mathbf{y}(1))}{k} + \frac{S_n^2(\mathbf{y}(0))}{n - k} - \frac{S_n^2(\boldsymbol{\tau})}{n}.$$

**Step 3:** The conditional variance is a random variable that depends on  $\mathbf{Z}$  only through its count, so the outer expectation averages the display above over the distribution of the count of treated units:

$$E_\Omega \left[ \text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \right] = \sum_{k=1}^{n-1} \Pr_\Omega (n_T(\mathbf{Z}) = k) \left( \frac{S_n^2(\mathbf{y}(1))}{k} + \frac{S_n^2(\mathbf{y}(0))}{n - k} - \frac{S_n^2(\boldsymbol{\tau})}{n} \right).$$

The three variances are fixed quantities, so the linearity of expectations splits the sum into three pieces and pulls each variance out front:

$$\begin{aligned} E_\Omega \left[ \text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \right] &= S_n^2(\mathbf{y}(1)) \sum_{k=1}^{n-1} \frac{\Pr_\Omega (n_T(\mathbf{Z}) = k)}{k} + S_n^2(\mathbf{y}(0)) \sum_{k=1}^{n-1} \frac{\Pr_\Omega (n_T(\mathbf{Z}) = k)}{n - k} \\ &\quad - \frac{S_n^2(\boldsymbol{\tau})}{n} \sum_{k=1}^{n-1} \Pr_\Omega (n_T(\mathbf{Z}) = k). \end{aligned}$$

The first sum is, by definition, the expected reciprocal  $E_\Omega[1/n_T(\mathbf{Z})]$ . The second sum is  $E_\Omega[1/n_C(\mathbf{Z})]$ ,

because  $n_C(\mathbf{Z}) = n - n_T(\mathbf{Z})$ , so the event  $\{n_T(\mathbf{Z}) = k\}$  is the event  $\{n_C(\mathbf{Z}) = n - k\}$ . The third sum equals one, because the count of treated units takes some value in  $\{1, \dots, n - 1\}$  with probability one. Substituting the three pieces yields the claim.  $\square$

Compared with Equation (12), the only change is that the fixed reciprocals  $1/n_T$  and  $1/n_C$  become the expected reciprocals  $E_\Omega[1/n_T(\mathbf{Z})]$  and  $E_\Omega[1/n_C(\mathbf{Z})]$ ; the correction term  $-S_n^2(\boldsymbol{\tau})/n$  is untouched. The expected reciprocals have no simple closed form, but the unbiasedness guide derives the distribution of the count of treated units,  $\Pr_\Omega(n_T(\mathbf{Z}) = k) = \binom{n}{k} p^k (1 - p)^{n-k} / c$  with  $c = 1 - p^n - (1 - p)^n$ , so direct summation computes them exactly:

$$E_\Omega \left[ \frac{1}{n_T(\mathbf{Z})} \right] = \sum_{k=1}^{n-1} \frac{1}{k} \Pr_\Omega (n_T(\mathbf{Z}) = k),$$

and likewise for  $E_\Omega[1/n_C(\mathbf{Z})]$ , with  $\frac{1}{n-k}$  in place of  $\frac{1}{k}$ . Simulation gives the same numbers without the formula: Draw many assignment vectors from the mechanism, discard any degenerate draws, and average the realized values of  $1/n_T(\mathbf{Z})$ ; by the law of large numbers the average converges to the expected reciprocal.

The two variance formulas share the correction term  $-S_n^2(\boldsymbol{\tau})/n$  and differ only in the reciprocals of the counts, so the comparison between the two mechanisms comes down to fixed versus random counts of treated and control units. Complete random assignment holds those counts at the balance the experimenter chooses, while simple random assignment lets chance push them away from that balance, and departures from balance only add to the variance. With  $p = \frac{1}{2}$ , even  $n$ , and the balanced design  $n_T = n_C = \frac{n}{2}$ , complete random assignment therefore never has larger variance than simple random assignment.

The conservative estimator extends in the same way. With random counts the estimator uses the realized counts,

$$\widehat{\text{Var}}[\hat{\tau}] = \frac{\widehat{S}_n^2(\mathbf{y}(1))}{n_T(\mathbf{Z})} + \frac{\widehat{S}_n^2(\mathbf{y}(0))}{n_C(\mathbf{Z})},$$

and each arm's sample variance requires at least two units, for the same reason the complete random assignment analysis assumes  $n_T, n_C \geq 2$ . I therefore condition on the event  $\mathcal{D} := \{2 \leq n_T(\mathbf{Z}) \leq n - 2\}$  that each arm contains at least two units.

**Proposition 3.** *Under simple random assignment,*

$$E_\Omega \left[ \widehat{\text{Var}}[\hat{\tau}] \mid \mathcal{D} \right] - \text{Var}_\Omega [\hat{\tau} \mid \mathcal{D}] = \frac{S_n^2(\boldsymbol{\tau})}{n}.$$

*Proof.* Every ingredient is already in hand, and the proof assembles those ingredients. First, each

count event  $\{n_T(\mathbf{Z}) = k\}$  with  $k \in \{2, \dots, n-2\}$  lies inside  $\mathcal{D}$ , so once the count of treated units is given, the additional conditioning on  $\mathcal{D}$  changes nothing. Second, conditional on any such count the distribution of  $\mathbf{Z}$  is complete random assignment — the unbiasedness guide’s conditional uniformity lemma again — so the conservative estimation results of the previous section hold at that count:

$$\mathbb{E}_\Omega \left[ \widehat{\text{Var}}[\hat{\tau}] \mid n_T(\mathbf{Z}) = k \right] = \text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z}) = k] + \frac{S_n^2(\boldsymbol{\tau})}{n}.$$

Third,  $\mathbb{E}_\Omega[\hat{\tau} \mid n_T(\mathbf{Z}) = k] = \tau_{\text{SATE}}$  for every  $k$ , so the conditional expectation of the estimator does not vary with the count of treated units.

Now average over the counts in  $\mathcal{D}$ . By the law of iterated expectations,

$$\mathbb{E}_\Omega \left[ \widehat{\text{Var}}[\hat{\tau}] \mid \mathcal{D} \right] = \mathbb{E}_\Omega \left[ \text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \mid \mathcal{D} \right] + \frac{S_n^2(\boldsymbol{\tau})}{n},$$

where the overshoot passes through the average untouched because the overshoot is the same constant at every count in  $\mathcal{D}$ . By the law of total variance, applied within the event  $\mathcal{D}$ ,

$$\text{Var}_\Omega [\hat{\tau} \mid \mathcal{D}] = \mathbb{E}_\Omega \left[ \text{Var}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \mid \mathcal{D} \right] + \text{Var}_\Omega \left[ \mathbb{E}_\Omega [\hat{\tau} \mid n_T(\mathbf{Z})] \mid \mathcal{D} \right],$$

and the second term is zero because the inner conditional expectation is the constant  $\tau_{\text{SATE}}$ . Subtracting the second display from the first leaves exactly  $S_n^2(\boldsymbol{\tau})/n$ .  $\square$

The estimator is exactly as conservative under simple random assignment as under complete random assignment.

## References

- Cochran, W. G. (1977). *Sampling Techniques* (3rd ed.). New York: John Wiley & Sons. [14](#)
- Cox, D. R. (1958). *Planning of Experiments*. New York: John Wiley & Sons. [2](#)
- Gerber, A. S. and D. P. Green (2012). *Field Experiments: Design, Analysis, and Interpretation*. New York: W. W. Norton. [12](#)
- Imbens, G. W. and D. B. Rubin (2015). *Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction*. New York: Cambridge University Press. [3](#)
- Pashley, N. E., G. W. Basse, and L. W. Miratrix (2021). Conditional as-if analyses in randomized experiments. *Journal of Causal Inference* 9(1), 264–284. [16](#)
- Rubin, D. B. (1980). Comment on “Randomization Analysis of Experimental Data: The Fisher

Randomization Test” by D. Basu. *Journal of the American Statistical Association* 75(371), 591–593. [2](#)

Rubin, D. B. (1986). Comment on “Statistics and Causal Inference” by P. W. Holland: Which ifs have causal answers. *Journal of the American Statistical Association* 81(396), 961–962. [2](#)