

# Notation and Setup for Design-Based Causal Inference\*

Thomas Leavitt

This guide collects, in one place, the notation and formal setup shared by the companion guides on (i) the unbiasedness of the Difference-in-Means estimator for both the Sample Average Treatment Effect (SATE) and the Population Average Treatment Effect (PATE), (ii) the variance of the estimator for the SATE and the estimation of that variance, and (iii) the variance of the estimator for the PATE and the estimation of that variance. Each of those guides gives just enough of the setup for its own proofs. This guide develops that setup in full. A reader who is already comfortable with the potential outcomes framework for randomized experiments can safely skim this guide.

## 1 Units, assignments, and the assignment mechanism

Let the index  $i \in \{1, \dots, n\}$  run over  $n$  units in a finite sample,  $\mathcal{S}_n$ , where  $n \geq 4$ . Of these  $n$  units, the design assigns  $n_T \geq 2$  to the treatment condition and  $n_C \geq 2$  to the control condition, where  $n_T + n_C = n$ . The bounds  $n_T \geq 2$  and  $n_C \geq 2$  (and hence  $n \geq 4$ ) play no role in the unbiasedness of the estimator or in the variance formula itself; they ensure that the conservative (Neyman) estimator of the variance (developed in a companion guide) is well defined. That estimator computes a separate sample variance within each treatment arm, dividing by  $n_T - 1$  for the treated units and by  $n_C - 1$  for the control units, and a sample variance requires at least two observations; the estimator therefore needs at least two units in each arm, i.e.,  $n_T \geq 2$  and  $n_C \geq 2$ , so that  $n = n_T + n_C \geq 4$ .

Let the binary indicator variable  $Z_i \in \{0, 1\}$  record whether unit  $i$  receives treatment ( $Z_i = 1$ ) or control ( $Z_i = 0$ ), and collect these indicators in the random assignment vector  $\mathbf{Z} = [Z_1, \dots, Z_n]^\top$ . The set of all logically possible assignment vectors is  $\{0, 1\}^n$ .

---

\*This is a live document that is subject to updating at any time.

Every assignment vector determines how many units it sends to treatment: Write  $n_T(\mathbf{z}) := \sum_{i=1}^n z_i$  for this count, and  $n_C(\mathbf{z}) := n - n_T(\mathbf{z})$ . Evaluated at the random assignment vector, this counting function yields  $n_T(\mathbf{Z})$ , itself a random variable and a function of  $\mathbf{Z}$  in exactly the way the potential outcomes below are functions of  $\mathbf{z}$ . The assignment mechanism places positive probability only on a subset of the  $2^n$  logically possible vectors. Under *complete random assignment* the design fixes the count, restricting the support to

$$\Omega = \{\mathbf{z} \in \{0, 1\}^n : n_T(\mathbf{z}) = n_T\},$$

where  $n_T$  is a constant that the experimenter chooses.

On  $\Omega$  the random count  $n_T(\mathbf{Z})$  equals the constant  $n_T$  with probability one, so I write the plain constant, just as  $z_i$  denotes the realized value of  $Z_i$ . The set  $\Omega$  is the support of  $\mathbf{Z}$ . Under complete random assignment,  $\mathbf{Z}$  is distributed uniformly on  $\Omega$ , and the number of elements in  $\Omega$ , denoted  $|\Omega|$ ,<sup>1</sup> is  $\binom{n}{n_T}$ . By contrast,  $n$  independent Bernoulli assignments would place positive probability on all  $2^n$  vectors. The definition of complete random assignment thus has two separable parts, and the guides use both: The fixed count determines the support, and uniformity is a further assumption about the distribution the mechanism places on that support.

I distinguish three objects that are easily conflated. The *assignment space*  $\{0, 1\}^n$  is the set of all conceivable assignments. The *support*  $\Omega \subseteq \{0, 1\}^n$  is the (typically smaller) set of assignments the mechanism can actually produce. And the *randomization distribution* is the probability distribution the mechanism places on  $\Omega$  (uniform, under complete random assignment). All of the randomness in what follows comes from this distribution. One convention holds throughout the guides:  $\Omega$  always denotes the support of  $\mathbf{Z}$  under the assignment mechanism in force, and when an argument moves between mechanisms I condition explicitly rather than change the symbol. The same way of building a support, holding a count fixed and collecting every vector that achieves it, extends to stratified designs: A blocked or matched study fixes a treated count within each stratum, and the study's support collects the assignment vectors that hit all of those counts at once.

The same framework accommodates *simple random assignment*, under which the mechanism assigns each of the  $n$  units independently, with a common assignment probability  $p \in (0, 1)$  for every unit. Two features change. First, no design constraint pins down the count, so  $n_T(\mathbf{Z})$  is a nondegenerate random variable. Second, I exclude the two degenerate assignment vectors (all treated and all control), under which one group is empty and the Difference-in-Means estimator is undefined; the support is therefore  $\Omega = \{\mathbf{z} \in \{0, 1\}^n : 0 < n_T(\mathbf{z}) < n\}$ , which has  $2^n - 2$  elements. The exclusion changes the distribution, so I state the distribution explicitly:  $\mathbf{Z}$  follows the distribution of  $n$  independent Bernoulli draws with success probability  $p$ , *conditioned on* the event  $\{0 < n_T(\mathbf{Z}) < n\}$ ,

---

<sup>1</sup>For an arbitrary set  $W$ , let  $|W|$  denote the cardinality of (i.e., the number of elements in) the set  $W$ .

and the conditioning breaks the independence of the coordinates.

Note where the probabilities live: Simple random assignment defines them at the level of individual units, and the distribution of the whole vector follows as a consequence — an assignment  $\mathbf{z}$  in the support has probability proportional to  $p^{n_T(\mathbf{z})}(1-p)^{n-n_T(\mathbf{z})}$ , which depends on  $\mathbf{z}$  only through its count and is uniform on the support only when  $p = \frac{1}{2}$ . Complete random assignment runs in the opposite direction: The distribution is defined at the level of the whole vector, uniform on  $\Omega$ , and the unit-level probabilities are consequences, which the unbiasedness guide derives.

Even though the design does not fix  $n_T(\mathbf{Z})$ , we can fix the count by conditioning on its realized value: The randomization distribution conditional on the event  $\{n_T(\mathbf{Z}) = n_T\}$  is exactly the complete random assignment distribution above, whatever the value of  $p$  (Pashley et al., 2021); the unbiasedness guide proves the reduction. Hence results that I derive under complete random assignment extend to simple random assignment after conditioning on the realized number of treated units.

## 2 The potential outcomes schedule and the Stable Unit Treatment Value Assumption

Adopting the terminology of Freedman (2009) and later Gerber and Green (2012), define a potential outcomes schedule as a vector-valued function  $\mathbf{y} : \{0, 1\}^n \rightarrow \mathbb{R}^n$  that maps each possible assignment vector  $\mathbf{z} \in \{0, 1\}^n$  to an  $n$ -dimensional vector of real-valued outcomes. The schedule lists how each of the  $n$  study participants would respond to every assignment  $\mathbf{z}$  that the experiment could in principle produce. The schedule is a *fixed* feature of the units; randomness enters only through which assignment the mechanism could select. I move in numbered steps from this general object to the familiar pair of potential outcomes: for each unit, one fixed outcome under treatment and another under control. I will use the same format, general object followed by assumption followed by licensed simplification, whenever an assumption changes what the notation must express.

**Step 1 (general object).** Unit  $i$ 's outcome under assignment  $\mathbf{z}$  is the  $i$ th coordinate of the schedule's value, written  $y_i(\mathbf{z})$ . Composed with the random assignment,  $y_i(\mathbf{Z})$  is a random variable that could in principle take as many as  $2^n$  distinct values, one for each assignment vector.

**Step 2 (assumption invoked).** Under the Stable Unit Treatment Value Assumption (SUTVA)<sup>2</sup> (Cox, 1958; Rubin, 1980, 1986), unit  $i$ 's outcome depends on the assignment vector only through

---

<sup>2</sup>SUTVA implies that (1) units in the experiment respond to only the treatment condition to which each unit is individually assigned and (2) the treatment condition is actually the same treatment for all units assigned to treatment and the control condition is the same for all units assigned to control.

unit  $i$ 's own coordinate: For any  $\mathbf{z}, \mathbf{z}'$  with  $z_i = z'_i$ ,  $y_i(\mathbf{z}) = y_i(\mathbf{z}')$ .

**Step 3 (licensed simplification).** By Step 2, the function  $\mathbf{z} \mapsto y_i(\mathbf{z})$  depends on  $\mathbf{z}$  only through  $z_i$ , so  $y_i(1)$  is well defined as the common value of  $y_i(\mathbf{z})$  across all  $\mathbf{z}$  with  $z_i = 1$ , and  $y_i(0)$  as the common value across all  $\mathbf{z}$  with  $z_i = 0$ . SUTVA thus collapses  $2^n$  numbers per unit to two fixed numbers per unit. The individual causal effect for unit  $i$  on the additive scale is  $\tau_i := y_i(1) - y_i(0)$ . The vectors  $\mathbf{y}(1)$  and  $\mathbf{y}(0)$  collect the treatment and control potential outcomes for all  $n$  units, and  $\boldsymbol{\tau}$  collects the  $n$  individual effects; all of these are fixed, finite population quantities.

**Step 4 (realization).** The observable outcome for unit  $i \in \{1, \dots, n\}$  is

$$Y_i := y_i(Z_i) = Z_i y_i(1) + (1 - Z_i) y_i(0),$$

which is random only through  $Z_i$ ; under the realized assignment  $\mathbf{Z} = \mathbf{z}$ , the observed outcome is the number  $y_i(z_i)$ . Potential outcomes are fixed, the assignment is random, and observable outcomes are random only because a random assignment selects which fixed potential outcome we see.

### 3 Estimands and the Difference-in-Means estimator

Write  $\bar{y}(1) := n^{-1} \sum_{i=1}^n y_i(1)$  and  $\bar{y}(0) := n^{-1} \sum_{i=1}^n y_i(0)$  for the finite population means of the treatment and control potential outcomes; both are fixed quantities. The target of interest is the Sample Average Treatment Effect (SATE),

$$\tau_{\text{SATE}} := n^{-1} \sum_{i=1}^n \tau_i = n^{-1} \sum_{i=1}^n (y_i(1) - y_i(0)) = \bar{y}(1) - \bar{y}(0),$$

which is likewise a fixed (though unknown) quantity. The Difference-in-Means estimator of  $\tau_{\text{SATE}}$  is

$$\hat{\tau} := \frac{\sum_{i=1}^n Z_i Y_i}{\sum_{i=1}^n Z_i} - \frac{\sum_{i=1}^n (1 - Z_i) Y_i}{\sum_{i=1}^n (1 - Z_i)}.$$

Unlike the estimand  $\tau_{\text{SATE}}$ , the estimator  $\hat{\tau}$  is a *random variable*, since  $\hat{\tau}$  is a function of the random assignment vector  $\mathbf{Z}$  and inherits all randomness from  $\mathbf{Z}$ . The denominators of the ratio form are the counts  $n_T(\mathbf{Z})$  and  $n_C(\mathbf{Z})$ ; under complete random assignment the design fixes them at  $n_T$  and  $n_C$ , so we can equivalently write  $\hat{\tau}$  as

$$\hat{\tau} = \frac{1}{n_T} \sum_{i=1}^n Z_i Y_i - \frac{1}{n_C} \sum_{i=1}^n (1 - Z_i) Y_i.$$

I use the ratio form when the number of treated units may be random (e.g., under simple random assignment) and the form with fixed denominators when  $n_T$  and  $n_C$  are fixed.

Because the only randomness is the assignment, I write  $E_\Omega[\cdot]$  and  $\text{Var}_\Omega[\cdot]$  for expectations and variances over the distribution of  $\mathbf{Z}$  on  $\Omega$ , to emphasize that they pertain to the randomization distribution alone. The subscript names the distribution in force on  $\Omega$ , not the set itself. In these guides that distribution is the uniform one just described. However, other distributions are possible on the same support, as when a sensitivity analysis considers departures from uniform assignment.

## 4 Finite population variances of the potential outcomes

Two equivalent families of finite population variances appear in the variance results. The first uses the divisor  $n - 1$ :

$$\begin{aligned} S_n^2(\mathbf{y}(1)) &= \left(\frac{1}{n-1}\right) \sum_{i=1}^n (y_i(1) - \bar{y}(1))^2 \\ S_n^2(\mathbf{y}(0)) &= \left(\frac{1}{n-1}\right) \sum_{i=1}^n (y_i(0) - \bar{y}(0))^2 \\ S_n^2(\boldsymbol{\tau}) &= \left(\frac{1}{n-1}\right) \sum_{i=1}^n (\tau_i - \tau_{\text{SATE}})^2. \end{aligned}$$

The second uses the divisor  $n$  and is the form that [Gerber and Green \(2012, Equation 3.4\)](#) uses:

$$\begin{aligned} \sigma_n^2(\mathbf{y}(1)) &= \left(\frac{1}{n}\right) \sum_{i=1}^n (y_i(1) - \bar{y}(1))^2 \\ \sigma_n^2(\mathbf{y}(0)) &= \left(\frac{1}{n}\right) \sum_{i=1}^n (y_i(0) - \bar{y}(0))^2 \\ \sigma_n(\mathbf{y}(0), \mathbf{y}(1)) &= \left(\frac{1}{n}\right) \sum_{i=1}^n (y_i(0) - \bar{y}(0)) (y_i(1) - \bar{y}(1)). \end{aligned}$$

The two families are related by  $\sigma_n^2(\cdot) = \frac{n-1}{n} S_n^2(\cdot)$ , and likewise for the covariance term. All of these are fixed, finite population quantities, and they will be the building blocks of the variance of  $\hat{\tau}$ . The naming rule deserves one explicit statement: The letter  $S$  marks the divisor one less than the size, the letter  $\sigma$  marks the divisor equal to the size itself, the subscript names the index set ( $n$  for the sample  $\mathcal{S}_n$ ,  $N$  for the superpopulation  $\mathcal{P}_N$ ), and the absence of a subscript marks the infinite superpopulation limit introduced at the end of the next section.

## 5 Extension to a superpopulation: the Population Average Treatment Effect

For results about the Population Average Treatment Effect (PATE), I embed the finite sample in a superpopulation. Consider a superpopulation  $\mathcal{P}_N$  of size  $N \geq n \geq 4$ , with index  $i \in \{1, \dots, N\}$ , and let simple random sampling draw  $n$  units from  $\mathcal{P}_N$  into the experimental sample  $\mathcal{S}_n$ , leaving the remaining  $N - n$  units unsampled. Among the  $n$  sampled units, complete random assignment sends  $n_T \geq 2$  to treatment and  $n_C \geq 2$  to control, exactly as above. The companion guides develop the superpopulation results under complete random assignment at this second stage; the variance guides close with extensions to simple random assignment.

The binary indicator  $R_i \in \{0, 1\}$  records whether unit  $i$  enters the sample ( $R_i = 1$ ) or stays out of it ( $R_i = 0$ ), and I collect these indicators in  $\mathbf{R} = [R_1, \dots, R_N]^\top$ . Write  $n(\mathbf{r}) := \sum_{i=1}^N r_i$  for the number of units a sampling vector selects; the support of  $\mathbf{R}$  is then  $\Pi = \{\mathbf{r} \in \{0, 1\}^N : n(\mathbf{r}) = n\}$ , which holds a count fixed exactly as  $\Omega$  does, now at the sampling stage. *Simple random sampling*, in the form the guides use, has a sample size fixed by design: The mechanism selects one of the  $\binom{N}{n}$  subsets of exactly  $n$  of the  $N$  units uniformly at random — sampling *without replacement* — which is the same as distributing  $\mathbf{R}$  uniformly on  $\Pi$ . The sample size is therefore not random. Every draw produces exactly  $n$  sampled units, so  $n(\mathbf{R}) = n$  with probability one, and I write the plain constant  $n$ , exactly as I write  $n_T$  under complete random assignment. The qualifier matters because a different mechanism, *Bernoulli sampling*, would flip a coin for each of the  $N$  units separately and so make the sample size  $n(\mathbf{R})$  a nondegenerate random variable, exactly as simple random assignment makes  $n_T(\mathbf{Z})$  one. Throughout the guides the sample size stays fixed at  $n$ , even in the infinite superpopulation model of the PATE variance guide, which draws a fixed number  $n$  of units independently.

One symbol  $\Omega$  serves every sample, and the license for that deserves its own statement. The assignment stage operates on whichever units the sampling stage selects, so in full generality its support could depend on the realized sample: Write  $\Omega_{\mathbf{r}}$  for the set of ways to assign the units that  $\mathbf{r}$  selects. Two features of the design make the single symbol sufficient. First, simple random sampling fixes the sample size, so every realized sample presents the same assignment problem —  $n$  units, of which the design sends exactly  $n_T$  to treatment. Second, the assignment mechanism is the same for every realized sample; the design does not, say, change  $n_T$  according to which units it happens to draw. Under these two conditions, listing each sample's units in a fixed order (by superpopulation index, say) identifies every  $\Omega_{\mathbf{r}}$  with the same set of binary  $n$ -vectors, and  $\Omega$  denotes that common object. After the identification, the assignment vector is independent of the sampling vector — every sample faces the same lottery — although which *units of the superpopulation* receive

treatment of course depends on both stages. When one formula mixes the stages, as in the products  $R_i Z_i$  of the companion guides, assignments wear superpopulation labels; for an unsampled unit the product is zero whatever value  $Z_i$  might take, so no definition of  $Z_i$  off the sample is ever needed. Under Bernoulli sampling the identification would fail at the first step: The length of the assignment vector itself would be random.

The potential outcomes deserve the same two-stage scrutiny, in the same numbered format as before.

**Step 1 (general object).** With both stages in view, unit  $i$ 's outcome could in principle depend on the entire configuration — on which sample the sampling stage draws and on which assignment the mechanism then selects — written  $y_i(\mathbf{r}, \mathbf{z})$ .

**Step 2 (assumption invoked).** I extend the stability assumption to both stages: Unit  $i$ 's outcome depends on the configuration only through unit  $i$ 's own treatment status. Being sampled does not itself change any outcome — sampling determines only whether an outcome becomes observable — and no unit's outcome depends on which other units the design samples or on how it assigns them.

**Step 3 (licensed simplification).** Every unit of  $\mathcal{P}_N$  then has just two fixed potential outcomes,  $y_i(1)$  and  $y_i(0)$ , defined whether or not the sampling stage ever selects the unit; the schedule of the earlier sections is the within-sample view of the same collapse. The estimand below is well defined precisely because unsampled units have potential outcomes too.

**Step 4 (observability).** For a sampled unit the observable outcome is  $Y_i = y_i(Z_i)$ , exactly as before; for an unsampled unit both potential outcomes exist, but neither is observable. Sampling and assignment never create or change potential outcomes; the two stages only determine which of the fixed values we see.

The estimand is the PATE,

$$\tau_{\text{PATE}} := N^{-1} \sum_{i=1}^N \tau_i,$$

and the corresponding population variances, which follow the naming rule of the previous section (letter  $S$ , divisor  $N - 1$ , subscript  $N$  for  $\mathcal{P}_N$ ), are

$$\begin{aligned} S_N^2(\mathbf{y}(1)) &= \left( \frac{1}{N-1} \right) \sum_{i=1}^N \left( y_i(1) - \frac{1}{N} \sum_{i=1}^N y_i(1) \right)^2 \\ S_N^2(\mathbf{y}(0)) &= \left( \frac{1}{N-1} \right) \sum_{i=1}^N \left( y_i(0) - \frac{1}{N} \sum_{i=1}^N y_i(0) \right)^2 \end{aligned}$$

$$S_N^2(\boldsymbol{\tau}) = \left( \frac{1}{N-1} \right) \sum_{i=1}^N (\tau_i - \tau_{\text{PATE}})^2.$$

The divisor  $N - 1$  pays off in the PATE variance guide: With this convention, the sample variance  $S_n^2(\cdot)$  is *exactly* unbiased for  $S_N^2(\cdot)$  under simple random sampling, with no correction factor.

In this setting the Difference-in-Means estimator has *two* sources of randomness, sampling  $\{R_i\}_{i=1}^N$  and assignment  $\{Z_i\}_{i=1}^n$ . I write  $E_{\Pi}[\cdot]$  and  $\text{Var}_{\Pi}[\cdot]$  for expectations and variances over the sampling distribution,  $E_{\Omega}[\cdot]$  and  $\text{Var}_{\Omega}[\cdot]$  for those over the assignment distribution (conditional on the sample), and the unsubscripted  $E[\cdot]$  and  $\text{Var}[\cdot]$  for those over both. The law of total variance links the overall variance to the stage-specific moments,

$$\text{Var}[\hat{\tau}] = E_{\Pi}[\text{Var}_{\Omega}[\hat{\tau}]] + \text{Var}_{\Pi}[E_{\Omega}[\hat{\tau}]],$$

which is the organizing identity for the PATE variance derivation. Finally, as  $N \rightarrow \infty$  with the population variances bounded, the quantities  $S_N^2(\cdot)$  converge to limits that I write  $\sigma^2(\cdot)$ , with no subscript; these limiting superpopulation variances govern the infinite superpopulation results in the PATE variance guide.

## References

- Cox, D. R. (1958). *Planning of Experiments*. New York: John Wiley & Sons. [3](#)
- Freedman, D. A. (2009). *Statistical Models: Theory and Practice* (2nd ed.). New York: Cambridge University Press. [3](#)
- Gerber, A. S. and D. P. Green (2012). *Field Experiments: Design, Analysis, and Interpretation*. New York: W. W. Norton. [3](#), [5](#)
- Pashley, N. E., G. W. Basse, and L. W. Miratrix (2021). Conditional as-if analyses in randomized experiments. *Journal of Causal Inference* 9(1), 264–284. [3](#)
- Rubin, D. B. (1980). Comment on “Randomization Analysis of Experimental Data: The Fisher Randomization Test” by D. Basu. *Journal of the American Statistical Association* 75(371), 591–593. [3](#)
- Rubin, D. B. (1986). Comment on “Statistics and Causal Inference” by P. W. Holland: Which ifs have causal answers. *Journal of the American Statistical Association* 81(396), 961–962. [3](#)