

# CS727 HW5

Zhili Feng

November 21, 2018

1. (5.4.19)

- (a) Let  $P = \{x : Dx \geq d; Fx = f; x \geq 0\}$ . We have  $P$  is closed. Note that  $P$  contains no line since any  $x$  with negative entry is not in  $P$ . Then by theorem 4.3.4,  $P = \text{conv}(X) + \text{pos}(Y)$ , where  $X$  is the set of extreme points of  $P$ , and  $Y$  is a set of points of  $\mathbb{R}^n$  such that  $\text{pos}(Y)$  is the set of extreme rays of  $P$ .

Now by Minkowski-Weyl, since  $P$  is polyhedral, for any  $x \in P$ , we can write  $x = \sum_{i=1}^p \lambda_i x_i + \sum_{i=1}^q \mu_i y_i$  such that  $\lambda_i \geq 0, \sum_i \lambda_i = 1, \mu_i \geq 0$ , and  $x_i \in X, y_i \in Y$ .

Note that we must have  $\langle c^*, y_i \rangle \geq 0$  for all  $i = 1, \dots, q$ . If there exists  $\langle c^*, y_i \rangle < 0$ , then we can make the corresponding  $\mu_i$  approaching  $\infty$  and  $\inf\{\langle c^*, x \rangle, x \in P\}$  will be  $-\infty$ . Hence:

$$\forall x \in P : \langle c^*, x \rangle \geq \left\langle c^*, \sum_{i=1}^p \lambda_i x_i \right\rangle$$

note  $\sum_{i=1}^p \lambda_i x_i \in P$  since  $P$  is convex.

Since  $X$  is finite, we can find  $x^* \in X$  with the smallest  $\langle c^*, x^* \rangle$ , i.e.  $\langle c^*, x^* \rangle \leq \langle c^*, x_i \rangle \forall x_i \in X$ . Then we have

$$\langle c^*, x^* \rangle \leq \left\langle c^*, \sum_{i=1}^p \lambda_i x_i \right\rangle, \text{ for all suitable } \lambda_i$$

Hence this  $x^*$  satisfies the condition that  $\forall x \in P, \langle c^*, x^* \rangle \leq \langle c^*, x \rangle$ . By definition,  $x^*$  attains the infimum that we want.

- (b) First let us rewrite the constraints in  $P$  in terms of  $Ax \leq a$ :

$$Dx \geq d; Fx = f; x \geq 0 \iff \underbrace{\begin{bmatrix} -D \\ F \\ -F \\ -I \end{bmatrix}}_A x \leq \underbrace{\begin{bmatrix} -d \\ f \\ -f \\ 0 \end{bmatrix}}_a$$

where  $I \in \mathbb{R}^{n \times n}$  is the identity matrix.

Let  $x_0 \in P$  be the feasible solution that satisfies base demand  $d_0$ . Use the Hoffman's theorem, there exists  $\gamma_A > 0$  such that given  $x_0$ , we can find  $x \in P$  (equivalently,  $Ax \leq a$ ) such that

$$\|x - x_0\| \leq \gamma_A \|(Ax_0 - a)_+\|$$

Note that for  $x_0$ :

$$Ax_0 = \begin{bmatrix} -D \\ F \\ -F \\ -I \end{bmatrix} x \leq \begin{bmatrix} -d_0 \\ f \\ -f \\ 0 \end{bmatrix}$$

therefore:

$$(Ax_0 - a)_+ \leq \left( \begin{bmatrix} -d_0 \\ f \\ -f \\ 0 \end{bmatrix} - \begin{bmatrix} -d \\ f \\ -f \\ 0 \end{bmatrix} \right)_+ = \left( \begin{bmatrix} d - d_0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right)_+$$

Let the solution we get from (a) be  $x^*$ , we have:

$$\begin{aligned} & \langle c^*, x^* \rangle \\ & \leq \langle c^*, x \rangle \\ & = \langle c^*, x_0 \rangle + \langle c^*, x - x_0 \rangle \\ & \leq c_0 + \|c^*\| \|x - x_0\| \\ & \leq c_0 + \|c^*\| \gamma_A \|(Ax_0 - a)_+\| \\ & \leq c_0 + \|c^*\| \gamma_A \|(d - d_0)_+\| \end{aligned}$$

Let  $\alpha = \gamma_A \|c^*\|$ , and we finish the proof.