



The globular cluster system of the Auriga simulations

Journal:	<i>Monthly Notices of the Royal Astronomical Society</i>
Manuscript ID	MN-19-3257-MJ.R1
Manuscript type:	Main Journal
Date Submitted by the Author:	n/a
Complete List of Authors:	Halbesma, Timo; Max-Planck-Institut für Astrophysik Grand, Robert; Max-Planck-Institut für Astrophysik, Gomez, Facundo; Universidad de La Serena Marinacci, Federico; University of Bologna, Physics and Astronomy Pakmor, Ruediger; Max-Planck-Institut für Astrophysik Trick, Wilma; Max-Planck-Institut für Astrophysik, Busch, Philipp; Max-Planck-Institut für Astrophysik, White, Simon; Max-Planck-Institut für Astrophysik, Cosmology
Keywords:	galaxies: formation < Galaxies, galaxies: star clusters: general < Galaxies

The globular cluster system of the Auriga simulations

Timo L. R. Halbesma¹*, Robert J. J. Grand¹, Facundo A. Gómez^{2,3}, Federico Marinacci⁴,
Rüdiger Pakmor¹, Wilma H. Trick¹, Philipp Busch¹, Simon D. M. White¹

¹ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85741 Garching, Germany

² Instituto de Investigación Multidisciplinar en Ciencia y Tecnología, Universidad de La Serena, Raúl Bitrán 1305, La Serena, Chile

³ Departamento de Física y Astronomía, Universidad de La Serena, Av. Juan Cisternas 1200 N, La Serena, Chile

⁴ Department of Physics & Astronomy, University of Bologna, via Gobetti 93/2, 40129 Bologna, Italy

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We investigate whether the galaxy and star formation model used for the Auriga simulations can produce a realistic globular cluster (GC) population. We compare statistics of GC candidate star particles in the Auriga haloes with catalogues of the Milky Way (MW) and Andromeda (M31) GC populations. We find that the Auriga simulations do produce sufficient stellar mass for GC candidates at radii and metallicities that are typical for the MW GC system (GCS). We also find varying mass-ratios of the simulated GC candidates relative to the observed mass in the MW and M31 GC systems for different bins of galactocentric radius-metallicity ($r_{\text{gal}} - [\text{Fe}/\text{H}]$). Overall, the Auriga simulations produce GC candidates with higher metallicities than the MW and M31 GCS and they are found at larger radii than observed. The Auriga simulations would require bound cluster formation efficiencies higher than ten percent for the metal-poor GC candidates, and those within the Solar radius should experience negligible destruction rates to be consistent with observations. GC candidates in the outer halo, on the other hand, should either have low formation efficiencies, or experience high mass loss for the Auriga simulations to produce a GCS that is consistent with that of the MW or M31. Finally, the scatter in the metallicity as well as in the radial distribution between different Auriga runs is considerably smaller than the differences between that of the MW and M31 GCSs. The Auriga model is unlikely to give rise to a GCS that can be consistent with both galaxies.

Key words: galaxies: formation – galaxies: star clusters: general.

1 INTRODUCTION

Globular star clusters (GCs) are old, bright, and ubiquitous. Various properties of GC systems (GCS) show correlations with their host galaxies. GCSs are believed to retain information about the galactic (gas) conditions at their times of formation. Thus they could offer a unique insight into the (chemodynamical) evolution of their parent galaxies, if the formation and evolution of GCs- and GC systems themselves is adequately understood. However, despite decades of research (e.g. Harris 1991; Harris 2001; West et al. 2004; Brodie & Strader 2006; Gratton et al. 2012; Kruijssen 2014; Forbes et al. 2018), consensus on an exhaustive picture of the formation of GCs is yet to be reached. Most galaxies host a GCS with a bimodal colour distribution that may reflect a bimodal metallicity distribution which is argued to arise from two distinct formation channels (Brodie & Strader 2006). GCs are typically referred to as ‘blue’ or ‘red’ (e.g. Zinn 1985; Gebhardt & Kissler-Patig 1999; Larsen et al. 2001) where the former class is more metal-poor, more radially extended,

and showing little to no rotation whereas the latter is more metal-rich, radially concentrated, and rotating with the galaxy.

The literature offers a wealth of formation scenarios, many of which focus on the distinct characteristics of the blue or red subpopulation. Peebles & Dicke (1968) and Peebles (1984) argue that GCs form as the earliest bound structures in the Universe (i.e. prior to formation of the main galaxy), noting that the Jeans length and mass shortly after recombination is consistent with typical GC masses and sizes. Diemand et al. (2005) and Boley et al. (2009) argue that the blue, metal-poor GCs ($[\text{Fe}/\text{H}] < -1$) form in radially biased dark matter (DM) halos at high redshift. Other recent hypotheses of GC formation prior to collapse of the proto-galaxy include formation in colliding supershells (Recchi et al. 2017), in supersonically induced gaseous objects (Chiou et al. 2019), or in high-speed collisions of dark matter subhaloes (Madau et al. 2019). Boylan-Kolchin (2017) suggests that (the blue subpopulation of) GCs could form in high density regions along the cosmic filament before or during collapse.

Other models also date GCs formation during formation of the proto-galaxy itself, for example as a result of thermal instabilities in hot gas-rich haloes (Fall & Rees 1985, also see the discussions in

* E-mail: Halbesma@MPA-Garching.MPG.DE

Kang et al. 1990). Alternative formation triggers are also explored, such as (other causes of) shock compression, or cloud-cloud collisions (e.g. Gunn 1980; Murray & Lin 1992; Harris & Pudritz 1994; Vietri & Pesce 1995; Larson 1996; Cen 2001).

Yet another hypothesis is that star cluster formation is triggered by major gas-rich (spiral) galaxy mergers (Schweizer 1987; Ashman & Zepf 1992), which is naturally expected within the framework of hierarchical assembly. One testable prediction of this scenario is the formation of young clusters in interacting and merging galaxies, which has been observed and are found to show remarkable similarities with GCs in the Milky Way (MW) (e.g. Whitmore & Schweizer 1995; Holtzman et al. 1996; Zepf et al. 1999; Whitmore et al. 1999). Moreover, modelling efforts of this framework yield GC (sub)populations consistent with various observables (e.g. Muratov & Gnedin 2010; Choksi et al. 2018), and the recent numerical simulation of an isolated dwarf-dwarf merger executed at very high resolution (baryonic mass $m_b \sim 4M_\odot$; softening $\epsilon = 0.1$ pc) produces star clusters that could be GC progenitors (Lahén et al. 2019).

As for the formation timeline, the scenarios (and flavours thereof) of Fall & Rees (1985) and Ashman & Zepf (1992) are intertwined because accretion and mergers continually occur during the hierarchical build-up of galaxies. Various (other) hierarchical formation channels thus combine different aspects of the aforementioned paradigms, such as GC formation in (small) galactic disks before they are accreted onto an assembling galaxy (e.g. Côté et al. 2000, 2002; Beasley et al. 2002; Gnedin 2003). We refer to Gnedin et al. (2001) for a scoreboard of GC formation models compared to observations of the MW GCSs.

An idea that has recently been studied extensively is the hypothesis that the formation mechanisms of young massive clusters (YMCs, see Portegies Zwart et al. 2010, for a review) and GCs are the same. The proximity of YMCs in the local Universe allows to study these processes with a level of detail that cannot be achieved for GCs at high redshifts. In this framework, differences between the two classes of objects are caused by nearly a Hubble time of (dynamical) evolution (e.g. Spitzer 1987). This picture is based on observed similarities between YMCs and GCs (e.g. Holtzman et al. 1992; Whitmore et al. 1999) and strengthened by observations of gravitationally lensed objects at high redshifts ($z = 2 - 6$). These sources have properties reminiscent of (local) YMCs and may be GC progenitors at times of formation (Vanzella et al. 2017; Johnson et al. 2017). The modelling work by Kruijssen et al. (2011, 2012); Kruijssen (2015) is now incorporated into cosmological zoom simulations and shows promising results (Pfeffer et al. 2018; Kruijssen et al. 2019).

The (mass) resolution of cosmological zoom simulations has reached the mass range populated by GCs, and the gravitational force softening can be as low as several parsec. A number of groups can thus incorporate formation of (globular) star clusters into their high-resolution hydrodynamical simulations. Ricotti et al. (2016) ran parsec scale simulations of the high-redshift universe prior to reionization (the simulations stop at $z = 9$), Li et al. (2017) implement a new subgrid model for star (cluster) formation and run simulations that reach $z = 3.3$, and the run of Renaud et al. (2017) reaches $z = 0.5$. Kim et al. (2018) find that mergers can push gas to high density that quickly forms clustered stars that end up tightly bound by the end of the simulation. A somewhat different approach couples semi-analytical models to DM-only simulations (Muratov & Gnedin 2010; Li & Gnedin 2014; Choksi et al. 2018; Choksi & Gnedin 2019b,a), also done in the work by El-Badry et al. (2019).

In this work we use state of the art simulations that produce realistic spiral galaxies (MW analogues) at redshift zero for which

several global properties are consistent with the observations. The question thus naturally arises whether the star formation histories of the simulations could give rise to a GCS similar to the MW and/or Andromeda (M31) GCSs. In particular, we use the Auriga simulations (Grand et al. 2017, hereafter G17), further described in Section 2. We investigate whether the star formation model implemented produces metallicity, radial, and metallicity-radial distributions that are consistent with the MW and/or M31 GCSs, and whether the model produces enough stellar mass with the right properties. Moreover, the simulations provide us with the history of the star particles, which allows us to investigate the differences between these properties for sub-populations that have formed *in situ* versus those that have been accreted. We focus on the metallicity and radial distributions because observations of the red and blue GCs show distinct differences between these properties.

The plan of the paper is as follows. We summarise the relevant characteristics of the Auriga simulations in Section 2, followed by a summary of the observations of the MW and M31 GCSs in Section 3 that we compare our simulations to in Section 4. We discuss our findings in Section 5 to come to our conclusions in Section 6.

2 THE AURIGA SIMULATIONS

We use the Auriga simulations (G17), a suite of high-resolution cosmological zoom simulations of MW analogues. The MW-mass selected initial conditions in the range $1 - 2 \times 10^{12} M_\odot$ were selected from the EAGLE DM-only box, see G17 for details. The simulations are performed with the AREPO code (Springel 2010; Pakmor et al. 2016) that solves the magnetohydrodynamical equations on a moving mesh.

The galaxy formation model includes primordial and metal-line cooling with self-shielding corrections. Reionization is completed at redshift 6 by a time-varying spatially uniform UV background (Faucher-Giguère et al. 2009; Vogelsberger et al. 2013). The interstellar medium (ISM) is described by an equation of state for a two-phase medium in pressure equilibrium (Springel & Hernquist 2003); cold clouds embedded in a hot ambient medium, whose mass and energy densities are described by processes important for star formation, such as the condensation of cold clouds and the photoevaporation of clouds. Stars form stochastically in thermally unstable gas with a density threshold of $n = 0.13 \text{ cm}^{-3}$, and consecutive stellar evolution is accounted for. The channel for energetic stellar feedback is from non-local type II supernovae (SN), which is modelled with wind particles (Marinacci et al. 2014; G17). Galactic winds enrich the ISM with metals from SNIa, SNII, and asymptotic giant branch stars (Vogelsberger et al. 2013). Supermassive black holes grow via Bondi–Hoyle–Lyttleton accretion (Bondi & Hoyle 1944; Bondi 1952), and feedback from the active galactic nuclei has a low-accretion and high-accretion mode (radio and quasar) (Springel et al. 2005; Marinacci et al. 2014; G17). Finally, the simulations follow the evolution of a magnetic field of 10^{-14} (comoving) G seeded at $z = 127$ (Pakmor & Springel 2013; Pakmor et al. 2014).

This model was tailored to the AREPO code and calibrated to reproduce key observables of galaxies, such as the history of the cosmic star formation rate density, the stellar mass to halo mass relation, and galaxy luminosity functions. The galaxy formation model produces realistic spiral galaxies at redshift zero that match observations such as sizes, morphologies (G17), HI disc properties (Marinacci et al. 2017), and the diversity of accreted stellar halo properties (Monachesi et al. 2019). Furthermore, the simula-

tions are able to reproduce major Galactic components, such as the chemical thin/thick disc dichotomy (Grand et al. 2018) and the recently discovered *Gaia* Sausage (Fattahi et al. 2019). They are thus a large set of detailed simulations similar enough to the Galaxy to make predictions for the GC population properties of the MW and M31.

The Auriga suite has a fiducial baryonic mass resolution $m_b = 5 \times 10^4 M_\odot$, with gravitational softening of collisionless particles $\epsilon = 369$ pc. Simulations at this resolution are referred to as level 4 (L4). Selected initial condition runs are accompanied by the lower (higher) resolution level L5 (L3). The resolution for these levels is $m_b = 4 \times 10^5 M_\odot$ with $\epsilon = 738$ pc, and $m_b = 6 \times 10^3 M_\odot$ with $\epsilon = 184$ pc, respectively. The mass resolution of the Auriga simulations is thus close $10^5 M_\odot$, to the characteristic peak mass of the lognormal GC mass distribution (Harris 1991), although the gravitational softening is two orders of magnitudes larger than typical GC radii of several parsec. High-density gaseous regions are thus not expected to produce surviving stellar clumps with masses and radii consistent with GCs because such objects would numerically disperse, even in the highest-resolution runs. On the other hand, we can investigate (statistical) properties of age-selected GC candidates because each star particle represents a single stellar population with a total mass that could be consistent with one GC. This means that their formation sites could be consistent with those of real-world GCs.

3 OBSERVATIONAL DATA

We describe observations of the MW GCS in Section 3.1 for these clusters, and of the M31 GCS in Section 3.2. We discuss age-estimates in Section 3.3, their distribution of galactocentric radius in Section 3.4, and their total mass in 2D bins of metallicity-galactocentric radius in Section 3.5. We focus on both metallicity and galactocentric radius because these data are available in the observations and in the Auriga simulations.

3.1 Milky Way

Harris (1996, 2010 edition; hereafter H96e10) provides a catalogue¹ of the MW GCS that contains properties of 157 GCs. The authors initially estimated the number of GCs in the MW GCS to be 180 ± 10 , thus, their catalogue to be $\sim 85\%$ complete. However, an additional 59 GC candidates have been discovered more recently by various authors. The total number of GCs in the MW might thus be 216 with recent estimates now anticipating an additional 30 GCs yet to be discovered (e.g. Ryu & Lee 2018, and references therein). We still use data from the Harris catalogue, but caution that it may (only) be 50-70% complete. Specifically, the relevant data fields that we use from H96e10 are the metallicity [Fe/H], the Galactic distance components X , Y , and Z (in kpc)², and absolute magnitude in the V band, M_V . We use the V band magnitude to calculate mass-estimates by assuming $M_{V,\odot} = 4.83$ and a mass to light ratio $M/L_V = 1.7 M_\odot/L_\odot$, the mean for MW clusters (McLaughlin & van der Marel 2005). We supplement the catalogue with age-estimates from isochrone fits to stars near the

main-sequence turnoff in 55 GCs (VandenBerg et al. 2013, hereafter V13).

3.2 Andromeda

Caldwell et al. (2011, hereafter C11) and Caldwell & Romanowsky (2016, hereafter CR16) present a uniform set of spectroscopic observations of the inner 1.6° (~ 21) kpc of M31 that is believed to be 94% complete. In addition, the outer stellar halo of M31 up to $r_{\text{proj}} \sim 150$ kpc is observed in the Pan-Andromeda Archaeological Survey (PAndAS, Huxor et al. 2014; Veljanoski et al. 2014; Mackey et al. 2019, hereafter H14). In fact, the work of H14 is incorporated in the latest public release³ of the C11 and CR16 data sets. Therefore we use the latest data set of CR16 because it is the most recent aggregated dataset of M31's GCS that contains GCs in the inner region and in the outer halo. The relevant fields in the CR16 dataset that we use are the age, metallicity, and the mass-estimate⁴. We calculate the galactocentric radii from the observed positions RA and DEC, as further discussed in Section 3.4.

3.3 Age-estimates

The top panel of Figure 1 shows a histogram of the age-estimates of the 55 MW GCs in V13 and 88 GCs in M31 for which age-estimates are available in CR16. The mean age of the MW GCs in this data set is 11.9 Gyr and the standard deviation is 0.8 Gyr. Furthermore, only 1 of the 55 GC age-estimates is below 10 Gyr. The M31 GCS has a mean age of 11.0 Gyr with a standard deviation of 2.2 Gyr, and 24 GCs have age-estimates below 10 Gyr with a minimum age of 4.8 Gyr. Based on these data, we find that the age distributions of the MW and M31 GCSs are not statistically consistent. The MW appears to host a GCS that is somewhat older than that of M31. This is somewhat surprising given that M31 is generally considered to be earlier type than the MW so that an older stellar population would naively be expected. However, we do caution that both data sets are incomplete and the age measurements have large uncertainties (of 1 – 2 Gyr). On the other hand, the magnitude of the uncertainty is insufficient to explain the low-age tail in M31.

3.4 Radial distribution

The bottom panel of Figure 1 shows the radial distribution of the MW and M31 GCS. We divide the radii of M31 by its virial radius⁵ $r_{\text{vir},\text{M31}} = 299$ kpc and multiply by $r_{\text{vir},\text{MW}} = 261$ kpc to account for the different intrinsic sizes of the two galaxies, adopting the values and cosmology from Patel et al. (2017). Values for r_{gal} are readily available in H96e10 (assuming $r_\odot = 8.0$ kpc), but the galactocentric radius of GCs in M31 is not available in CR16. Therefore, we follow Wang et al. (2019, section 4.1) to calculate the projected radius r_{proj} from the observed positions, adopting M31's central position from the NASA Extragalactic Database⁶ (α_0, δ_0) = ($0^{\text{h}}42^{\text{m}}44.35^{\text{s}}, +41^\circ 16'08.63''$) and the distance $D_{\text{M31}} = 780$ kpc

³ Last revised 23 Sep 2015, see https://www.cfa.harvard.edu/oir/eg/m31clusters/M31_Hectospec.html

⁴ The authors assumed $M/L_V = 2$ independent of [Fe/H]

⁵ r_{vir} is the radius in a spherical ‘top-hat’ perturbation model at which the average density reaches an overdensity of $\Delta_{\text{vir}} = 357$ times the background density

⁶ <https://ned.ipac.caltech.edu/>

¹ See https://www.physics.mcmaster.ca/Fac_Harris/mwgc.dat
² In a Sun-centered coordinate system: X points toward Galactic center, Y in direction of Galactic rotation, and Z toward the North Galactic Pole. We calculate the galactocentric radius $r_{\text{gal}} = \sqrt{(X - r_\odot)^2 + Y^2 + Z^2}$, assuming the solar radius $r_\odot = 8$ kpc.




Figure 1. *Top:* Age distribution of 55 GCs in the MW (data from [VandenBerg et al. 2013](#)) and 88 GCs in M31 (data from [Caldwell & Romanowsky 2016](#)). *Bottom:* Distribution of galactocentric radius in the MW and M31. The dotted lines show the subset of data that also have age measurements (i.e. the same sample as in the top panel). We divide the radii of M31 by $r_{\text{vir},\text{M31}} = 299$ kpc and multiply by $r_{\text{vir},\text{MW}} = 261$ kpc ([Patel et al. 2017](#)) to compensate for the different intrinsic sizes of both galaxies.

([McConnachie et al. 2005](#); [Conn et al. 2012](#)). We estimate r_{gal} as ‘average deprojected distance’ $r_{\text{gal}} = r_{\text{proj}} \times (4/\pi)$.

The solid lines show the distributions using all available data (because the sky coordinates are known for each GC), while the subset for which age-estimates are available is indicated using dotted lines. The latter shows a narrower range of radii than the full data set: it appears that few age-estimates are available for the innermost (< 1 kpc) GCs, and none for those beyond roughly 20 kpc (the halo GCs). We compare the two distributions of the full data set (solid lines) and find that the MW has more GCs in the range 1 – 4 kpc than M31 when accounting for the larger number of total GCs in M31. Interestingly, the two distributions show a similar trend for $r_{\text{gal}} > 4$ kpc and host a subpopulation of halo GCs. Our naive expectation would be that the spiral galaxies in the Local Group host similar GC systems. However, based on a two-sample KS test we find that the radial distributions of the GCS of the MW and M31 are not statistically consistent with being drawn from the same underlying distribution due to substantial differences at intermediate radii.

3.5 Total GC mass in metallicity-radial space

We show the two-dimensional mass-weighted metallicity-radius distribution of the MW and the M31 GCS in the top and bottom panel of Figure 2, respectively. Note that the number of GCs is different than in Figure 1 because here we plot $[\text{Fe}/\text{H}]$ between -2.5 and 0 and r_{gal} between 1 and 250 kpc whereas Figure 1 shows the full range of metallicities and radii.

The observations indicate very few GCs with high metallicities at large radii (the three bins in the upper right corner, both for MW and for M31), and relatively few GCs at large radii in general ($r_{\text{gal}} > 30$ kpc; right column: 11 GCs or 7.3% in the MW and 17 or 4.6% in M31). Moreover, M31 hosts more metal-rich ($[\text{Fe}/\text{H}] > -1$) GCs in each radial bin in comparison to the MW after accounting for the fact that M31 hosts a larger GCS. Finally, given that the marginalized (i.e. the metallicity and radial) distributions are not statistically consistent between the two galaxies, we find that the two-dimensional distributions are also not consistent. More generally, the GCS of the MW differs significantly from that of M31, not only in the number of clusters but also in their distribution over radius and metallicity. We compare these observations to the Auriga simulations in Section 4.3.

4 RESULTS

We define GC candidates in the Auriga simulations as all star particles older than 10 Gyr based on the age distribution of the MW GCS (top panel of Figure 1), adopting the definition of a GC from [Portegies Zwart et al. \(2010\)](#) and following the method of [Renaud et al. \(2017\)](#). However, we caution that his definition, although statistically consistent with the estimated ages of all GCs in the MW, is not consistent with the tail of younger clusters as stated in Section 3.3, and further discussed in Section 5.3.

Throughout our analysis we compare distributions of age-selected GC candidates (age > 10 Gyr) with those of all star particles (i.e. with no age cut). The mean number of star particles in the Auriga L4 simulations that we select as GC candidates is $3.1 \cdot 10^5$ (with a mean total mass of $1.15 \cdot 10^{10} M_{\odot}$), and the mean number of all star particles is $1.93 \cdot 10^6$ ($7.3 \cdot 10^{10} M_{\odot}$). The subset of GC candidates is thus around 15% of all the stars in the galaxy when averaged over all Auriga L4 haloes. We further split the GC candidates into an *in situ* subset (defined as GC candidates that are bound to the most-massive halo/subhalo in the first snapshot that the particle was recorded), and an accreted subpopulation (those that have formed *ex situ* but are bound to the most-massive halo/subhalo at $z = 0$). A GC candidate either formed *in situ* or it was accreted. We consider the metallicity distribution in Section 4.1, the distribution of galactocentric radii in Section 4.2, and the combination of the two in Section 4.3.

4.1 Metallicity distribution

We investigate whether the star formation model implemented in Auriga produces metallicity distributions consistent with the MW and M31 GCSs, and whether the simulations generate sufficient total mass at metallicities typical for the MW and M31 GCSs. To visually inspect the former we show the normalized metallicity distribution of three specific Auriga galaxies in Figure 3 in comparison to the MW and M31 $[\text{Fe}/\text{H}]$ distributions.




Figure 2. *Top:* Mass-weighted r_{gal} -[Fe/H] distribution of 151 GCs in the MW (data from [Harris 1996](#), 2010 ed.), colour-coded by the **median** (values also shown in each bin). The numbers in parenthesis show how many star particles fall within the bin. *Bottom:* Same for M31, showing the 366 GCs in CR16 (data from [Caldwell & Romanowsky 2016](#)). Radii of M31 are again divided by $r_{\text{vir},\text{M31}}$ and multiplied by $r_{\text{vir},\text{MW}}$. Note that the range of the colourmap differs in both figures,

We select Au4-10⁷, Au4-21, and Au4-4 to show the distributions of three individual simulation runs that we consider representative for specific behaviour, and to highlight that different runs give rise to different distributions. We show the age-selected GC candidates as well as the accreted and *in situ* subsets. The top panel of Figure 3 shows the MW and M31 GCSs, where we overplot a double Gaussian for the MW GCS by adopting literature values of the mean μ and standard deviation σ of the metal-rich and metal-poor subpopulations ([Harris 2001](#)).

We find that the age cut, on average, lowers the mean metallicity $\langle[\text{Fe}/\text{H}]\rangle$ from 0.0 to -0.6 . This reflects the enrichment history of the galaxy because the age cut removes the more metal-rich stellar population which shifts the mean metallicity towards lower values. Furthermore, accreted GC candidates generally have lower mean




Figure 3. *Top:* Metallicity distribution of the MW and M31 GCS. The solid purple (magenta) line shows the GCSs of the MW (M31), and we indicate the mean value with an arrow at the bottom of the x-axis. We overplot a double Gaussian for the MW GCS using purple dotted lines, taking the mean and dispersion from ([Harris 2001](#)). *Other panels:* three selected Auriga haloes as indicated in the upper right corner. In each of these three panels we show the GC candidates in beige, and use a beige dotted line for all star particles. We split the GC candidates up into two subpopulations, those that have formed *in situ* (brown), and those that have been accreted (teal). The numbers in parenthesis show how many star particles are included in the subset.

metallicities than the *in situ* subset with differences of roughly ~ 0.3 dex for the majority of the simulation runs. This behaviour can be seen in Au4-21, while a slightly larger difference of ~ 0.5 dex is seen for Au4-10 (as plotted), Au4-16, Au4-17, Au4-18 and Au4-22. However, this trend is reversed for Au4-1 and Au4-4 for which the *in situ* subset has a lower mean metallicity instead. Au4-1 is undergoing a major merger at redshift zero and we find a mean metallicity in this simulation $\mu = -1.51$ (-0.74) for the *in situ* (accreted) GC candidates, although the former consists of only 1019 particles (1.3% of all GC candidates, and with a total mass of $5 \times 10^7 M_{\odot}$). For Au4-4, only 10.8% of the GC candidates is classified as *in situ* (compared to *in situ* fractions of 40 – 80% for other haloes). We note that Au4-4 has undergone a major merger around 2 Gyr lookback time, which is probably why the accreted population is more metal-rich. After inspection of the same figure for every one of the 30 Auriga L4 haloes we find that the simulations produce (sub)populations of GC candidates that are more metal-rich

⁷ The nomenclature is ‘Au’ for Auriga, followed by the resolution level (4) and halo number (10, indicating which set of initial conditions was used).


6 *T. L. R. Halbesma et al.*

Figure 4. First vs. second central moment of the Auriga L4 metallicity distributions. Each cross (for a given colour) represents one simulation. The beige (brown) [teal] crosses show the values calculated using all (only *in situ*) [only accreted] GC candidates. Beige triangles indicate that all stars were used. The purple (magenta) cross denotes our calculation using all MW (M31) observations (which would be appropriate for a unimodal distribution). The purple solid (open) dots indicate the literature values of a bimodal Gaussian fit to the data (values from [Ashman & Zepf 1998](#), page 38), showing the metal-rich (metal-poor) component of the MW (separated at $[\text{Fe}/\text{H}] = -1$ as indicated by the purple dotted vertical line).

than the MW and M31 GCSs. Moreover, none of the simulations produces GC candidates with a bimodal metallicity distribution.

We show the mean metallicity and standard deviation of all 30 Auriga L4 haloes in Figure 4 to illustrate the individual behaviour for the full set of Auriga galaxies. The beige crosses are to be compared to the purple and magenta crosses, which show the mean value of all MW and M31 GCs, respectively. In addition, we show the metal-rich population of the MW GCS using a solid dot, and the metal-poor population with an open dot. We caution that these literature values result from individual Gaussian fits to subsets of the observational data cut at $[\text{Fe}/\text{H}] = -1$. We do not include corresponding data points calculated using such an artificial cut for the simulated star particles. It is interesting that the mean metallicities of the *in situ* GC candidates appear roughly consistent with that of the metal-rich population of the MW despite the lack of such a metallicity cut, although the simulations show larger dispersions. The latter could simply be caused by the hard separation of the MW data into two groups, which means the range is smaller and the resulting dispersion lower. With regard to the M31 GCS, so far no definitive consensus has been reached in the literature concerning uni- bi- or trimodality in the $[\text{Fe}/\text{H}]$ distribution, but CR16 argues that the data, after removal of younger objects due to improved age classification, hints at three populations separated at $[\text{Fe}/\text{H}] = -0.4$ and $[\text{Fe}/\text{H}] = 1.5$. None of the Auriga subsets has a mean value that offers much hope of reconciling the simulated distribution with the lowest metallicity group in the M31 GCS. This is also true for the metal-poor population of the MW with a mean value of -1.6 . The main result of this figure is that all Auriga L4 galaxies have metallicity distributions with (much) larger mean values than observed for the MW and M31 GCS, and that we systematically find lower mean metallicities for the accreted GC candidates than for those that have formed *in situ*.

We now turn to our second goal, which is to see whether suffi-




Figure 5. Mass-weighted metallicity distribution of star particles in the Auriga L4 simulations. We show the median value of all Auriga haloes for all stars (beige dotted) and GC candidates (i.e. stars with age > 10 Gyr; beige solid). The latter subset is further split up between stars that formed *in situ* (brown solid), and those that were accreted (teal solid). Shaded regions indicate the 25-75 percentile range. The MW (M31) GCS is shown in purple (magenta). The middle (bottom) panel shows the ratio of the simulated mass to the mass in the MW (M31) GCS.

cient total mass in GC candidates is produced by the Auriga simulations. To answer this question we plot a mass-weighted metallicity distribution in Figure 5. We show the median over all the 30 Auriga L4 haloes and the 25-75 percentile range as shaded regions, which shows the scatter between runs that have different initial conditions, thus, have varying merger histories. We chose to aggregate the data to indicate general trends that we find when the GC candidates are split up into the accreted and *in situ* subsets, rather than selecting typical examples of individual (simulated) galaxies. Once again we notice that the peak metallicity shifts down from $[\text{Fe}/\text{H}] = 0$ to $[\text{Fe}/\text{H}] = -0.6$ for GC candidates compared to all stars, and we find that the mass at the peak lowers by roughly one dex. The mass budget of the GC candidates is dominated by the *in situ* subpopulation above $[\text{Fe}/\text{H}] = -1$, and by the accreted subset below this value. We show the MW GCS in purple and that of M31 in magenta, and notice that the difference between the MW and M31 distributions is substantially larger than the scatter between different Auriga galaxies, particularly around $[\text{Fe}/\text{H}] = -1$. In addition, M31 does host (a handful of) GCs with $[\text{Fe}/\text{H}] < -2.5$ as well as GCs with $[\text{Fe}/\text{H}] > 0$ while the MW does not.

We show the ratio of the simulated to the observed profiles in




Figure 6. Median and standard deviation of the logarithm of radial distribution of star particles in each of the 30 Auriga L4 haloes compared to the MW (M31) GCS shown in purple (magenta). The Auriga (M31) radii are divided by $r_{\text{vir,Au}}$ ($r_{\text{vir,M31}}$) and multiplied by $r_{\text{vir,MW}}$ to compensate for different intrinsic sizes of the galaxies.

the middle and bottom panels. This mass excess can be thought of the ‘mass budget’ that the Auriga GC candidates can ‘afford to lose’ (due to a combination of smaller than unity bound cluster formation efficiencies combined with a Hubble time of dynamical evolution), while still producing sufficient mass at the right metallicities. In particular, the combined efficiency would have to increase with decreasing metallicity for Auriga GC candidates to produce a population of GC candidates that is consistent with the MW. For the GC candidates in M31 we find a constant mass ratio up to $[\text{Fe}/\text{H}] = -0.9$, above which the simulations produce a higher mass ratio with increasing metallicity. If dynamical evolution is not expected to more efficiently disrupt GCs of higher metallicity, then we would find that the efficiency to form bound star clusters would have to decrease with increasing metallicity. This trend could be consistent with the cluster formation efficiency model of Kruijssen (2012) as lower metallicity star particles formed at earlier times when the bound cluster formation efficiencies are higher due to higher pressure birth environments.

4.2 Radial distribution

We repeat the analysis of the previous subsection for the distribution of galactocentric radius instead of metallicity. We divide the Auriga radii by $r_{\text{vir,Au}}$ and multiply by $r_{\text{vir,MW}}$ to compensate for the different intrinsic sizes of the Auriga galaxies. We look for general trends present in all Auriga L4 galaxies. Figure 6 shows the median and standard deviation of $\log_{10}(r_{\text{gal}})$ of star particles in all Auriga L4 simulations, respectively. The purple and magenta cross show the median of the MW and M31 GCs, which are to be compared to the beige crosses. The solid dot now shows our calculation of the median value of the ‘red’ metal-rich population, and the open dot that of the ‘blue’ metal-poor population. We split the radii up into the metal-rich/metal-poor groups by taking a metallicity cut at $[\text{Fe}/\text{H}] = -1$ as is done in the literature.

We notice that the *in situ* GC candidates are more centrally distributed, whereas the accreted subsets have a larger radial extent. This is not surprising because the classification of *in situ* requires star particles to have formed within the virial radius, thus they




Figure 7. Mass-weighted radial distribution of star particles in the Auriga L4 simulations, where the Auriga (M31) radii are again divided by $r_{\text{vir,Au}}$ ($r_{\text{vir,M31}}$) and multiplied by $r_{\text{vir,MW}}$. We show the median value of all Auriga haloes for all stars (beige dotted) and GC candidates (i.e. stars with age > 10 Gyr; beige solid). The latter subset is further split up between stars that formed *in situ* (brown solid), and those that were accreted (teal solid). Shaded regions indicate the 25-75 percentile range. The MW (M31) GCS is shown in purple (magenta). The middle (bottom) panel shows the ratio of the simulated mass to the mass in the MW (M31) GCS.

could naturally be expected to end up at small galactocentric radii. Accreted star particles, on the other hand, have formed in another (sub)halo beyond the virial radius, thus, they would first have to migrate inwards in order to populate the innermost radii. Moreover, we find that the simulations have a wider dispersion in $\log(r_{\text{gal}})$ than the MW GCS, while the dispersion of M31 seems to lie within the range of dispersions found in the Auriga galaxies. Furthermore, the median of the MW GCS lies roughly within the range of values produced by the Auriga simulations, whereas the median of M31 is somewhat larger and slightly closer to typical median values of the accreted GC candidates in the Auriga simulations. The larger radial extent of the M31 GCS is generally believed to hint at a more accretion-dominated origin of the GCS, and may reflect a richer accretion history of M31 in comparison to the MW.

Figure 7 shows the mass-weighted radial distribution of the Auriga L4 haloes. We notice a subtle peak around 10 kpc for all star particles that is not present for the GC candidates, indicating that the stellar disc is no longer present when applying the latter selection criterion. Furthermore, we find that the dominant contribution to the total mass in GC candidates changes from those formed *in situ* to the accreted population around 10 kpc. Again we show the mass


8 *T. L. R. Halbesma et al.*

Figure 8. Mass-weighted $[\text{Fe}/\text{H}]$ - r_{gal} distribution of all 30 Auriga L4 haloes. Here we consider only the GC candidates and colour-code by the median mass (values also shown in each bin). The numbers in parenthesis show how many star particles fall within the bin. Note that the range of the colourmap differs from both panels in Figure 2 (for improved contrast within the plot).

ratio of the simulations compared to the MW and M31 GCS. We find a decreasing mass ratio with increasing radius in the range 0.2 to ~ 5 kpc, followed by an increase attributed to the accreted sub-population. For the MW we notice that significantly fewer GCs are found beyond 40 kpc than for M31, and that accreted GC candidates contribute mostly to the Auriga GCS at these radii. We further investigate a breakdown of the total mass in Auriga GC candidates into bins of both metallicity and radius in the following section.

4.3 Total mass in metallicity-radial space

We investigate whether the Auriga simulations still produce sufficient mass when the GC candidates are two-dimensionally binned in $[\text{Fe}/\text{H}]$ and r_{gal} . First we sum the total mass of GC candidates in each bin (for an individual Auriga simulation), then we calculate the median total mass in each bin over all 30 Auriga L4 haloes. The result is shown in Figure 8, which can be compared to the observed total GC mass shown in Figure 2. We notice that the simulations produce relatively little mass in metal-poor GC candidates at small radii and in metal-rich GC candidates at large radii. However, there is still few times $10^7 M_{\odot}$ produced by the simulations in the uppermost 3 bins where no observed GCs are found.

To ease this comparison between the simulations and observations we take the ratio of the total mass of GC candidates in Auriga to the total GC mass in the observations. We use a diverging colour map centered around the overall median of the mass ratio (i.e. we subtract the median of these 25 bins from the value in each bin). We show this in the top and bottom panel of Figure 9 for the MW and M31, respectively. Note that the 3 bins in the top right are empty (white) because the mass ratio cannot be calculated due to zero mass in the observations. If the product of the bound cluster formation efficiency and consecutive GC disruption over a Hubble time would be constant, then the value in each bin would be zero. The red and blue bins indicate deviations from such a scenario and indicate a higher and lower than average mass excess in the simulations. The median value used to shift the mass ratio is 2.68 and 1.69 dex for the MW and M31, respectively. This means that the median ‘mass

budget’ that the simulations ‘can afford to lose’ is roughly a factor 480 and 50 for the MW and M31, respectively. The lowest mass surplus for both galaxies is the bin with metallicities in the range $[\text{Fe}/\text{H}] \in [-2.5, -1.5]$ and galactocentric radius $r_{\text{gal}} \in [3, 8]$ kpc, for which the simulations produce a factor $10^{2.68}/10^{1.33} \approx 22$ and $10^{1.69}/10^{0.75} \approx 9$ more mass in GC candidates than is observed in the MW and M31 respectively. The highest factor for the MW is 2.1×10^5 , and 1.1×10^3 for M31.

We now focus on four quadrants of this diagram, split at $[\text{Fe}/\text{H}] = -1$ (metal-poor or metal-rich) and $r_{\text{gal}} = 8$ kpc (inside or outside the Solar radius). The majority of the mass in Auriga GC candidates is produced in metal-rich star particles inside the Solar radius. The total mass is dominated by the *in situ* component, for $[\text{Fe}/\text{H}]$ as well as for r_{gal} as can be seen in Figure 5 and Figure 7. The total mass excess in this region increases with decreasing metallicity for the MW and M31, and there is a weak trend of decreasing over-production with increasing radius for the *in situ* subsets but no trend with radius for the accreted GC candidates. As for the metal-rich GC candidates outside the Solar radius, we find that the mass excess with respect to the median is largest in this corner for the MW and M31. The main reason for this trend are the lower GC counts in the observations, whereas the Auriga simulations efficiently form metal-rich GC candidates and have no problem to produce them at large radii. The opposite is true for the metal-poor GC candidates inside the Solar radius, where the mass surplus with respect to the median is smallest for both galaxies. However, the disruption rates are expected to be particularly high for GCs in the inner regions of galaxies which must have orbits with pericentres close to the galactic bulge or bar. In addition, disk crossings in this region will cause spikes in the tidal tensor due to steep density gradients that could be highly disruptive to the GCs. The bins with the lowest ‘mass budget’ are in this region (a factor 9 for the MW and 22 for M31). The bound cluster formation efficiency has to exceed 10%, and none of the GC candidates in this region can experience significant mass loss in order for Auriga to produce sufficient mass in this quadrant. Finally, the metal-poor GC candidates outside the Solar radius have lower than average mass ratios, which is not surprising given that the Auriga simulations overall underproduce low-metallicity stars particles.

5 DISCUSSION

5.1 Metallicity distribution

GCs are ~ 0.5 dex more metal-poor than spheroid stars observed at the same radius for almost all galaxies (Harris 1991). Our selection function (age cut) does lower the mean metallicity of GC candidates by 0.5 dex with respect to all star particles, but we still find that the GC candidates in the Auriga simulations are more metal-rich than the MW and M31 GCSs. Although the (old) star particles represent single stellar populations with a mass consistent with that of GCs, they are in fact statistical tracers of the stellar population of the galaxy as a whole. Therefore only a (small) fraction of the star particles may represent plausible formation sites of GCs, whereas the majority represents (halo) field stars - the disk component effectively falls outside our selection of star particles due to the age cut. It is implausible that all halo stars come from disrupted GCs because the total mass in GCSs at redshift zero is generally a few orders of magnitude lower than the total mass of stellar haloes of galaxies. If stellar haloes solely consist of disrupted GCs, then the GC disruption rate would have to be considerably higher than expected from




Figure 9. The top (bottom) panel shows the logarithm of the ratio of simulated mass to mass in the MW (M31) GCS, i.e. the logarithm of the mass ratio, minus the median value of these 25 bins. The values are 2.68 and 1.69 dex for the MW and M31, respectively. Red (blue) indicates a higher (lower) overall mass excess. The number in each bin shows the colour-coded value that is plotted, and the (median) star particle count is given in parenthesis. Note that the three bins in the upper right corner are white because the observations have zero mass there.

theoretical models. Moreover, disrupted dwarf galaxies are more likely to contribute to the build up of the stellar halo than GCs (e.g. Deason et al. 2015; Conroy et al. 2019). Overall, our set of GC candidates most likely contains an overabundance of regular stars. It could be that the resulting metallicity distribution is then biased towards higher values than observed for real GCs. An alternative explanation could be overmixing of metals at early times.

In general, the colour (metallicity) distribution of most GCSs appears to be bimodal with typical separating value $[\text{Fe}/\text{H}] \sim -1$ (Zinn 1985; Gebhardt & Kissler-Patig 1999; Larsen et al. 2001; Peng et al. 2006). Indeed, Ashman & Zepf (1998) and Harris (2001) find that the MW GCS has a bimodal $[\text{Fe}/\text{H}]$ distribution. Observations of GCs in M31, however, may be best split into three distinct metallicity groups (Caldwell & Romanowsky 2016). The numerical simulation of Renaud et al. (2017) does yield a bimodal metallic-

ity distribution where the metal-poor population is dominated by accreted star particles and the metal-rich population by *in situ* stars.

However, we find that none of the Auriga simulations produces a bimodal metallicity distribution for age-selected GC candidates. Interestingly, the cross-over point above (below) which the mass-weighted metallicity distribution of GC candidates is dominated by those that have formed *in situ* (were accreted) does coincide with the separation between the metal-rich and metal-poor populations of GCSs at $[\text{Fe}/\text{H}] = -1$, as for the MW. Moreover, we do find that the mean metallicity shifts when we split the GC candidates according to birth location (accreted or *in situ*). In particular, the mean values of the *in situ* GC candidates are roughly consistent with the metal-rich MW GCs. On the other hand, we also notice that most of the simulations have a similar mean value for all GC candidates, and the simulated *in situ* GC candidates have larger dispersions than is claimed for metal-rich GCs in the MW. Moreover, the offset between the means of the metal-rich and metal-poor populations in the MW is 1 dex, a factor 2 – 3 larger than the offset that we find between the *in situ* and accreted populations. However, we caution that Kruijssen et al. (2019) suggests to reserve the ‘*ex situ*’ classification for accretion after $z = 2$, when the central galaxy has formed and accretion unambiguously contributes to the radially extended halo GC population. In fact, our classification might not hold for Au4-1.

Finally, we note that Brodie & Strader (2006) compares the number of metal-poor GCs to the stellar halo mass and find⁸ $T_{\text{blue}}^n \sim 100$, while the number of metal-rich GCs compared to the bulge mass yields $T_{\text{red}}^n \sim 5$, and therefore conclude that the formation efficiency of metal-poor GCs is 20 times higher than the metal-rich GCs with respect to field stars. We find that a gradual increase in formation efficiency of GCs with respect to field stars would be required with decreasing metallicity for the GC candidates in the Auriga simulations to yield sufficient total mass to be consistent with the MW or M31 GCS. In addition, the metal-poor GC candidates would have to experience low mass loss rates, especially those within the Solar radius.

5.2 Radial distribution

The M31 GCS has a factor 2-3 more GCs at large radii compared to the MW GCS, which may indicate that M31 has a richer merger history than the MW (Caldwell & Romanowsky 2016). In the Auriga simulations we find that the GC candidates at radii larger than ~ 20 kpc are indeed dominated by accreted star particles. However, we select star particles that are bound to the main halo and main subhalo, which means that we include particles up to the virial radius r_{200} . The Auriga simulations have no problem populating the stellar halo up to the virial radius, even with our additional age cut. On the other hand, the MW and M31 have fewer GCs at large radii. We expect tidal disruption to be less efficient at larger radii, thus, the formation efficiencies of the accreted GC candidates would have to be lower. Alternatively, the fraction of old star particles in our subset of GC candidates may contain considerably more star particles that represent stars in the old stellar halo that have been accreted during hierarchical buildup of the galaxy. Our classification of GC candidates could be improved for the accreted subset in particular. For example, we could select only a subset of the accreted star particles as GC candidates where the number of GCs

⁸ The specific frequency T is the number of GCs per $10^9 M_{\odot}$ of galaxy stellar mass.

10 *T. L. R. Halbesma et al.*

would be determined by the satellite halo mass, given the $M_{\text{vir}} - N_{\text{gal}}$ correlation found by [Burkert & Forbes \(2019\)](#). However, we did not include such an analysis in this study.

5.3 Lower age cut in the GC selection function

Our method is sensitive to the definition that we adopt to classify GC candidates (e.g. [Forbes et al. 2018](#)). Therefore we explore how our results change if we include all star particles older than 8 Gyr, and if we further lower the age cut to 6 Gyr. The average fraction of GC candidates that we select from the all the star particles increases to 35% and 55% for an age cut of 8 and 6 Gyr, respectively.

We find that including more recently formed star particles in our subset of GC candidates shifts the mean metallicity to higher values. Moreover, the total mass in metal-rich ($[\text{Fe}/\text{H}] > -1$) GC candidates increases for a lower age threshold and moves closer to the dotted line that shows the distribution of all star particles in Figure 5. However, we notice that the Auriga simulations, on average, do not form new stars with a metallicity below $[\text{Fe}/\text{H}] = -1$ more recent than 6 Gyr ago: the behaviour of Figure 5 does not change for metallicities below $[\text{Fe}/\text{H}] = -1$ when we change the age cut to 8 or 6 Gyr, and the slope of the mass ratio for metallicities above $[\text{Fe}/\text{H}] = -1$ is slightly steeper. In addition, we still find that the accreted GC candidates are more metal-poor than those that have formed *in situ*.

The qualitative behaviour of the median distribution of galactocentric radii (Figure 7) is unaffected by lowering the age cut other than including more star particles (more mass) which leads to a higher normalisation and a higher mass ratio with respect to the observed GC systems of the MW and M31. The qualitative behaviour presented in Figure 9 does not change when the age cut is lowered to 8 or 6 Gyr. GC candidates is not affected definition because our selection function mostly includes star particles that have formed before roughly $z = 2$. Overall, reducing the age cut worsens the agreement between the observed GCSs and the candidate GCs in the Auriga galaxies.

6 SUMMARY AND CONCLUSIONS

We investigate statistics of age-selected (> 10 Gyr) GC candidates in the Auriga simulations and compare the simulations to the MW and M31 GCSs to test whether the star formation model implemented for the Auriga simulations could give rise to metallicity and radial distributions that are consistent with these two observed GC systems. Based on our analysis we draw the following conclusions.

- The star formation model implemented in the Auriga simulations produces metallicity distributions that are more metal-rich than the MW and M31 GCSs. For most of the Auriga simulations we find that the subset of accreted GC candidates has a lower mean metallicity than the *in situ* subpopulation. However, the accreted subsets are still more metal-rich than the MW and M31 GCSs. Moreover, none of the simulated subsets has a mean metallicity as low as the mean of the ‘blue’ metal-poor component of the MW GCS.

- We find that the difference between the metallicity distributions of the MW and M31 GCSs is big compared to the scatter in the Auriga GC candidates of different Auriga runs. The radial distributions of the MW and M31 GCSs are somewhat more similar, but they are not consistent with being drawn from the same distribution. The scatter between the radial distributions of different

Auriga simulations is smaller than the difference between the MW and M31 GCSs.

- GC candidates in the Auriga simulations may be found out to r_{200} , given that our selection function of star particles selects all stars bound to the main subhalo of the halo. The total mass of GC candidates is dominated by accreted star particles at radii beyond 20 kpc. The GCs in the MW and M31, on the other hand, have a much smaller radial extent.

- Tidal forces are larger in the inner regions of galaxies than in the outer halo. Therefore, we expect dynamical evolution to more strongly affect GC candidates at small radii than at large radii. However, we find that the Auriga simulations produce more mass in GC candidates with high metallicities at large radii than with low metallicities at small radii. The cluster formation efficiency would have to increase with decreasing metallicity for GC candidates in the Auriga simulations to be consistent with the MW GCS. This trend of over-production of old star particles that are metal-rich and at large radii compared to observed GCs is less clear for the M31 GCS.

- Overall, we conclude that the metallicity and radial distribution of age-selected star particles in the Auriga simulations are not consistent with globular cluster system of the Milky Way or with that of the Andromeda galaxy.

ACKNOWLEDGEMENTS

We thank the anonymous referee for a thoughtful and elaborate report. TLRH acknowledges support from the International Max-Planck Research School (IMPRS) on Astrophysics. TLRH thanks the Max Planck Computing and Data Facility for maintaining the Freya compute cluster, where the simulations were post-processed. FM is supported by the program “Rita Levi Montalcini” of the Italian MIUR. FAG acknowledges financial support from CONICYT through the project FONDECYT Regular Nr. 1181264, and funding from the Max Planck Society through a Partner Group grant. This work has been supported by the European Research Council under ERC-StG grant EXAGAL- 308037. Part of the simulations of this paper used the SuperMUC system at the Leibniz Computing Centre, Garching, under the project PR85JE of the Gauss Centre for Supercomputing. This work used the DiRAC Data Centric system at Durham University, operated by the Institute for Computational Cosmology on behalf of the STFC DiRAC HPC Facility ‘www.dirac.ac.uk’. This equipment was funded by BIS National E-infrastructure capital grant ST/K00042X/1, STFC capital grant ST/H008519/1 and STFC DiRAC Operations grant ST/K003267/1 and Durham University. DiRAC is part of the UK National E-Infrastructure.

The analysis in this work was performed using the Python ([van Rossum & de Boer 1991](#)) programming language, the IPython ([Perez & Granger 2007](#)) environment, the NumPy ([van der Walt et al. 2011](#)), SciPy ([Jones et al. 2001](#)), and Astropy ([Astropy Collaboration et al. 2013](#)) packages. Plots were created with Matplotlib ([Hunter 2007](#)), using the perceptually uniform colour maps from [Kovesi \(2015\)](#) for two-dimensional plots. This research has made use of NASA’s Astrophysics Data System Bibliographic Services.

REFERENCES

- Ashman K. M., Zepf S. E., 1992, [ApJ](#), **384**, 50
 Ashman K. M., Zepf S. E., 1998, *Globular Cluster Systems*
 Astropy Collaboration et al., 2013, [A&A](#), **558**, A33

- Beasley M. A., Baugh C. M., Forbes D. A., Sharples R. M., Frenk C. S., 2002, *MNRAS*, **333**, 383
- Boley A. C., Lake G., Read J., Teyssier R., 2009, *ApJ*, **706**, L192
- Bondi H., 1952, *MNRAS*, **112**, 195
- Bondi H., Hoyle F., 1944, *MNRAS*, **104**, 273
- Boylan-Kolchin M., 2017, *MNRAS*, **472**, 3120
- Brodie J. P., Strader J., 2006, *ARA&A*, **44**, 193
- Burkert A., Forbes D., 2019, arXiv e-prints,
- Caldwell N., Romanowsky A. J., 2016, *ApJ*, **824**, 42
- Caldwell N., Schiavon R., Morrison H., Rose J. A., Harding P., 2011, *AJ*, **141**, 61
- Cen R., 2001, *ApJ*, **560**, 592
- Chiou Y. S., Naoz S., Burkhardt B., Marinacci F., Vogelsberger M., 2019, arXiv e-prints,
- Choksi N., Gnedin O. Y., 2019a, arXiv e-prints,
- Choksi N., Gnedin O. Y., 2019b, *MNRAS*, **486**, 331
- Choksi N., Gnedin O. Y., Li H., 2018, *MNRAS*, **480**, 2343
- Conn A. R., et al., 2012, *ApJ*, **758**, 11
- Conroy C., Naidu R. P., Zaritsky D., Bonaca A., Cargile P., Johnson B. D., Caldwell N., 2019, *ApJ*, **887**, 237
- Côté P., Marzke R. O., West M. J., Minniti D., 2000, *ApJ*, **533**, 869
- Côté P., West M. J., Marzke R. O., 2002, *ApJ*, **567**, 853
- Deason A. J., Belokurov V., Weisz D. R., 2015, *MNRAS*, **448**, L77
- Diemand J., Madau P., Moore B., 2005, *MNRAS*, **364**, 367
- El-Badry K., Quataert E., Weisz D. R., Choksi N., Boylan-Kolchin M., 2019, *MNRAS*, **482**, 4528
- Fall S. M., Rees M. J., 1985, *ApJ*, **298**, 18
- Fattahi A., et al., 2019, *MNRAS*, **484**, 4471
- Faucher-Giguère C.-A., Lidz A., Zaldarriaga M., Hernquist L., 2009, *ApJ*, **703**, 1416
- Forbes D. A., et al., 2018, *Proceedings of the Royal Society of London Series A*, **474**, 20170616
- Gebhardt K., Kissler-Patig M., 1999, *AJ*, **118**, 1526
- Gnedin O. Y., 2003, in Kissler-Patig M., ed., *Extragalactic Globular Cluster Systems*. p. 224 (arXiv:astro-ph/0210556), doi:10.1007/10857603_35
- Gnedin O. Y., Lahav O., Rees M. J., 2001, arXiv Astrophysics e-prints,
- Grand R. J. J., et al., 2017, *MNRAS*, **467**, 179
- Grand R. J. J., et al., 2018, *MNRAS*, **474**, 3629
- Gratton R. G., Carretta E., Bragaglia A., 2012, *A&ARv*, **20**, 50
- Gunn J. E., 1980, in Hanes D., Madore B., eds, *Globular Clusters*. p. 301
- Harris W. E., 1991, *ARA&A*, **29**, 543
- Harris W. E., 1996, *AJ*, **112**, 1487
- Harris W. E., 2001, *Globular Cluster Systems*. Springer Berlin Heidelberg, p. 223–408, doi:10.1007/3-540-31634-5_2, https://doi.org/10.1007/3-540-31634-5_2
- Harris W. E., Pudritz R. E., 1994, *ApJ*, **429**, 177
- Holtzman J. A., et al., 1992, *AJ*, **103**, 691
- Holtzman J. A., et al., 1996, *AJ*, **112**, 416
- Hunter J. D., 2007, *Computing in Science and Engineering*, **9**, 90
- Huxor A. P., et al., 2014, *MNRAS*, **442**, 2165
- Johnson T. L., et al., 2017, *ApJ*, **843**, L21
- Jones E., Oliphant T., Peterson P., et al., 2001, *SciPy: Open source scientific tools for Python*, http://www.scipy.org/
- Kang H., Shapiro P. R., Fall S. M., Rees M. J., 1990, *ApJ*, **363**, 488
- Kim J.-h., et al., 2018, *MNRAS*, **474**, 4232
- Kovesi P., 2015, arXiv e-prints, p. arXiv:1509.03700
- Kruijssen J. M. D., 2012, *MNRAS*, **426**, 3008
- Kruijssen J. M. D., 2014, *Classical and Quantum Gravity*, **31**, 244006
- Kruijssen J. M. D., 2015, *MNRAS*, **454**, 1658
- Kruijssen J. M. D., Pelupessy F. I., Lamers H. J. G. L. M., Portegies Zwart S. F., Icke V., 2011, *MNRAS*, **414**, 1339
- Kruijssen J. M. D., Pelupessy F. I., Lamers H. J. G. L. M., Portegies Zwart S. F., Bastian N., Icke V., 2012, *MNRAS*, **421**, 1927
- Kruijssen J. M. D., Pfeffer J. L., Crain R. A., Bastian N., 2019, *MNRAS*, **486**, 3134
- Lahén N., Naab T., Johansson P. H., Elmegreen B., Hu C.-Y., Walch S., 2019, arXiv e-prints,
- Larsen S. S., Brodie J. P., Huchra J. P., Forbes D. A., Grillmair C. J., 2001, *AJ*, **121**, 2974
- Larson R. B., 1996, in Morrison H. L., Sarajedini A., eds, *Astronomical Society of the Pacific Conference Series Vol. 92, Formation of the Galactic Halo...Inside and Out*. p. 241
- Li H., Gnedin O. Y., 2014, *ApJ*, **796**, 10
- Li H., Gnedin O. Y., Gnedin N. Y., Meng X., Semenov V. A., Kravtsov A. V., 2017, *ApJ*, **834**, 69
- Mackey A. D., et al., 2019, *MNRAS*, **484**, 1756
- Madau P., Lupi A., Diemand J., Burkert A., Lin D. N. C., 2019, arXiv e-prints,
- Marinacci F., Pakmor R., Springel V., 2014, *MNRAS*, **437**, 1750
- Marinacci F., Grand R. J. J., Pakmor R., Springel V., Gómez F. A., Frenk C. S., White S. D. M., 2017, *MNRAS*, **466**, 3859
- McConnachie A. W., Irwin M. J., Ferguson A. M. N., Ibata R. A., Lewis G. F., Tanvir N., 2005, *MNRAS*, **356**, 979
- McLaughlin D. E., van der Marel R. P., 2005, *ApJS*, **161**, 304
- Monachesi A., et al., 2019, *MNRAS*, **485**, 2589
- Muratov A. L., Gnedin O. Y., 2010, *ApJ*, **718**, 1266
- Murray S. D., Lin D. N. C., 1992, *ApJ*, **400**, 265
- Pakmor R., Springel V., 2013, *MNRAS*, **432**, 176
- Pakmor R., Marinacci F., Springel V., 2014, *ApJ*, **783**, L20
- Pakmor R., Springel V., Bauer A., Mocz P., Munoz D. J., Ohlmann S. T., Schaaf K., Zhu C., 2016, *MNRAS*, **455**, 1134
- Patel E., Besla G., Sohn S. T., 2017, *MNRAS*, **464**, 3825
- Peebles P. J. E., 1984, *ApJ*, **277**, 470
- Peebles P. J. E., Dicke R. H., 1968, *ApJ*, **154**, 891
- Peng E. W., et al., 2006, *ApJ*, **639**, 95
- Perez F., Granger B. E., 2007, *Computing in Science and Engineering*, **9**, 21
- Pfeffer J., Kruijssen J. M. D., Crain R. A., Bastian N., 2018, *MNRAS*, **475**, 4309
- Portegies Zwart S. F., McMillan S. L. W., Gieles M., 2010, *ARA&A*, **48**, 431
- Recchi S., Wünsch R., Palouš J., Dinnbier F., 2017, *Ap&SS*, **362**, 183
- Renaud F., Agertz O., Gieles M., 2017, *MNRAS*, **465**, 3622
- Ricotti M., Parry O. H., Gnedin N. Y., 2016, *ApJ*, **831**, 204
- Ryu J., Lee M. G., 2018, *ApJ*, **863**, L38
- Schweizer F., 1987, in Faber S. M., ed., *Nearly Normal Galaxies. From the Planck Time to the Present*. pp 18–25
- Spitzer L., 1987, *Dynamical evolution of globular clusters*
- Springel V., 2010, *MNRAS*, **401**, 791
- Springel V., Hernquist L., 2003, *MNRAS*, **339**, 289
- Springel V., Di Matteo T., Hernquist L., 2005, *MNRAS*, **361**, 776
- VandenBerg D. A., Brogaard K., Leaman R., Casagrande L., 2013, *ApJ*, **775**, 134
- Vanzella E., et al., 2017, *MNRAS*, **467**, 4304
- Veljanoski J., et al., 2014, *MNRAS*, **442**, 2929
- Vietri M., Pesce E., 1995, *ApJ*, **442**, 618
- Vogelsberger M., Genel S., Sijacki D., Torrey P., Springel V., Hernquist L., 2013, *MNRAS*, **436**, 3031
- Wang S., Ma J., Liu J., 2019, arXiv e-prints
- West M. J., Côté P., Marzke R. O., Jordán A., 2004, *Nature*, **427**, 31
- Whitmore B. C., Schweizer F., 1995, *AJ*, **109**, 960
- Whitmore B. C., Zhang Q., Leitherer C., Fall S. M., Schweizer F., Miller B. W., 1999, *AJ*, **118**, 1551
- Zepf S. E., Ashman K. M., English J., Freeman K. C., Sharples R. M., 1999, *AJ*, **118**, 752
- Zinn R., 1985, *ApJ*, **293**, 424
- van Rossum G., de Boer J., 1991, *CWI Quarterly*, **4**, 283
- van der Walt S., Colbert S. C., Varoquaux G., 2011, *Computing in Science and Engineering*, **13**, 22

This paper has been typeset from a \LaTeX file prepared by the author.

Introduction

We thank the referee for a thoughtful and elaborate report. We address the comments in sections with headings that match those of the referee report, starting with the following listing of our response to the first 2 paragraphs in the introduction.

1. We updated the abstract and conclusion to clarify that we meant to describe what would be needed for the environmental dependence of the formation and disruption of bound clusters for the Auriga simulations to be consistent with the observations, rather than to argue that the Auriga simulations reflect these trends.
2. We feel that our conclusion would be too strongly phrased if we would generalise our results obtained using the Auriga simulations to cosmological simulations in general. We find that age-selected stellar particles in the Auriga simulations do not reproduce key properties of observed GC systems in the Milky Way and M31. However, our results might reflect characteristics that are specific to Auriga, e.g. that the galaxy formation model overmixes metals at early times. Therefore we did add a final bulletpoint to our conclusion to clarify the above, but we limit the scope to the Auriga simulations. Thanks for pointing this out!

Furthermore, we clarified in the manuscript that we investigate the metallicity and galactocentric radius because these properties are available for the observed GC systems and in the simulations.

General comments about style

- Numerical quantities are now described in arabic numbers instead of words.
- We now first use the name of the physical quantities in the given example, followed by the symbolic representation.
- The text has been updated to reduce the usage of parenthesis.
- Colloquial expressions have been rephrased.
- We replaced ‘Sec.’ by ‘Section’ to refer to sections in the present work. Moreover, we changed ‘Sec.’ to ‘section’ to refer to a section in one of the references.
- All acronyms are introduced the first time they appear in the text (MW, M31, GC, GCS, YMC, ISM, SN, DM, and Au). The subset thereof that appears in the abstract is also introduced the first time it appears.
- The colour palette has been updated to improve colourblind-friendliness.
- We no longer use blue and red to prevent confusion because they are typically used for metal-poor and metal-rich populations.

Major comments and concerns

- We updated the introduction to make it more clear why we consider the metallicity and galactocentric radius in light of *in situ* formation or accretion of GC candidates.
- We added section 5.3 to discuss how our results change if we adopt an age cut of 8 or 6 Gyr.

Additional major comments

- We added a section 5.3 to discuss how our results change if we adopt an age cut of 8 or 6 Gyr instead of 10 Gyr.
- We clarified that the number of observed GCs differs between Figure 1 and Figure 2 because the latter shows metallicities between -2.5 and 0, galactocentric radii between 1 and 250 kpc. Figure 1, on the other hand, shows the full range of radii, and the top panel of Figure 3 shows the full range of metallicities.
- We normalise the Auriga simulations by the virial radius of the dark matter halo to compensate for scatter between different simulation runs. We deliberately chose for r_{vir} because there are several stellar length scales that could be used (e.g. the effective radius of the bulge, the radial scale-length or vertical scaleheight of the disk, the optical radius, the extent of the stellar halo, the half-mass stellar radius, etc). The simulations show a wide variety between properties of the Auriga galaxies with little correlation between these different stellar length scales. Therefore we could introduce unknown biases by normalising to any one of these options which is why we use the virial radius instead.

Comments for each section

Introduction

- We clarified that the blue and red GCs are observed to have different colour distributions that may reflect a different metallicity distribution which is argued to arise from two distinct formation channels. We briefly summarise that differences between their properties include the metallicity and radial distributions, which is why we investigate these properties in the Auriga simulations.
- We removed the quoted sentence.

Sect. 2 -

- We added the halo mass range of the Auriga simulations.
- The last paragraph should now make more sense after updating the introduction.

Sect. 3 -

- We added references to section 3.3, 3.4, and 3.5 in the first paragraph of section 3.
- Section 3.2 has been restructured.
- We removed the standard error on the mean and replaced ‘dispersion’ by ‘standard deviation’ in section 3.3.
- We added a sentence to clarify the last statement in section 3.4.

Sect. 4 -

- We added the mean number of GC candidates compared to the mean number of star particles in the Auriga L4 galaxies. In addition, Figure 3 now shows the number of star particles included in each subset.
- We clarified at various places whether we mean the mean or median among the simulations, or if we refer to the absolute value of one of the galaxies.
- We added the suggestion in the 3rd paragraph of section 4.1.
- Bimodality in GC systems is now discussed in the introduction.
- Figure 4 - now has a vertical line to indicate the metallicity cut between metal-poor and metal-rich GC subpopulations in the Milky Way.
- We added a sentence about the cluster formation efficiency model of Kruijssen (2012) in the last paragraph of section 4.1.
- We now show the median instead of the mean for the galactocentric radius in section 4.2 and Figure 6.
- Figure 6: no changes were made for rescaling the galaxies (i.e. we still use the virial radius)
- Figure 5 and 7: shaded regions now show the 25-75th percentiles.
- We added a brief discussion of Figure 8.
- We clarified that we are referring to Figure 5 and Figure 7 for the claims about the *in situ* and accreted component in the third paragraph of section 4.3.

- Figure 9: The values are removed from the masked upper right corner bins.
- Figure 2,8,9: The values of the upper and right axes have been removed from Figure 2 and 9 (observations) for clarity (rather than the suggestion to add axis labels and units). However, we did not mask these bins for Figure 8 (Auriga) because it illustrates our finding that the simulations produce a considerable amount of mass in GC candidates with low metallicities at large radii.

Sect. 5 -

- We added references to Deason et al. (2015) and Conroy et al. (2019) in the first paragraph of section 5.1.
- We included a discussion of bimodality in metallicities in the introduction. In addition to that, the results of Usher et al. (2012) “imply that most early-type galaxies in the SLUGGS survey have bimodal metallicity as well as colour distributions”, and “Combined with previous studies, nine of 12 galaxies show evidence for bimodality while in the remaining three galaxies bimodality cannot be ruled out.”. Our interpretation of this study seems to differ from that of the referee.
- We still find that the accreted star particles are more metal-poor than those that have formed *in situ* if we lower the age cut to 8 or 6 Gyr. This has been added in section 5.3.
- We removed the quoted sentence.
- Our definition of ‘accreted GC candidate’ considers when the GC candidate was accreted onto the main object, not when it was formed. The age cut does not affect the time of accretion, but the time of accretion could have been before redshift $z = 2$ (which would be inconsistent with the definition of Kruijssen et al. 2019).
- We clarified which which galaxy Brodie & Strader provide the quoted number.
- We clarified footnote 8. of Kruijssen (2019) for *ex situ* GC formation.

Sect. 6 -

- We repeat the goal of the study in section 6.
- We clarified the 4th point.

Minor comments

- We clarified that ‘blue’ GC subpopulation means metal-poor with $[\text{Fe}/\text{H}] < -1$
- The units of mass-to-light ratio have been corrected.
- The indicated typos have been corrected.
- ‘missing error bars and references for the virial radii’ → The reference was given in a footnote, but is now moved to the main text.
- 4.1: ‘which model do the authors refer to?’ → added ‘star formation’.
- 4.1: ‘The top half of the left figure’ → ‘The top panel of Figure 3’.

The globular cluster system of the Auriga simulations

Timo L. R. Halbesma¹*, Robert J. J. Grand¹, Facundo A. Gómez^{2,3}, Federico Marinacci⁴,
Rüdiger Pakmor¹, Wilma H. Trick¹, Philipp Busch¹, Simon D. M. White¹

¹ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85741 Garching, Germany

² Instituto de Investigación Multidisciplinar en Ciencia y Tecnología, Universidad de La Serena, Raúl Bitrán 1305, La Serena, Chile

³ Departamento de Física y Astronomía, Universidad de La Serena, Av. Juan Cisternas 1200 N, La Serena, Chile

⁴ Department of Physics & Astronomy, University of Bologna, via Gobetti 93/2, 40129 Bologna, Italy

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We investigate whether the galaxy and star formation model used for the Auriga simulations can produce a realistic **GC-globular cluster (GC)** population. We compare statistics of GC candidate star particles in the Auriga haloes with catalogues of the Milky Way (MW) and Andromeda (M31) GC populations. We find that the Auriga simulations do produce sufficient stellar mass for GC candidates at radii and metallicities that are typical for the MW GC system (GCS). We ~~do observe also find~~ varying mass-ratios of the simulated GC candidates relative to the observed mass in the MW and M31 GC systems for different bins of galactocentric radius-metallicity ($r_{\text{gal}}-[\text{Fe}/\text{H}]$). ~~This may imply that different values for the combined product of the bound cluster formation efficiency and the GC disruption rate are needed.~~ Overall, the Auriga simulations produce GC candidates with higher metallicities than the MW and M31 GCS and they are found at larger radii than observed. ~~Bound~~ The Auriga simulations would require bound cluster formation efficiencies higher than ten percent ~~are needed~~ for the metal-poor GC candidates, and those within the Solar radius should experience negligible destruction rates to be consistent with observations. GC candidates in the outer halo, on the other hand, should either have low formation efficiencies, or experience high mass loss for the Auriga simulations to produce a GCS that is consistent with that of the MW or M31. Finally, the scatter in the metallicity as well as in the radial distribution between different Auriga runs is considerably smaller than the differences between that of the MW and M31 GCSs. The Auriga model is unlikely to give rise to a GCS that can be consistent with both galaxies.

Key words: galaxies: formation – galaxies: star clusters: general.

1 INTRODUCTION

Globular star clusters (GC)s are old, bright, and ubiquitous. Various properties of GC systems (GCS) show correlations with their host galaxies. GCSs are believed to retain information about the galactic (gas) conditions at their times of formation. Thus they could offer a unique insight into the (chemodynamical) evolution of their parent galaxies, if the formation and evolution of GCs and GC systems themselves is adequately understood. However, despite decades of research ~~(the extensive literature on GCs is summarised in several books and review articles, e.g. Harris 1991; Harris 2001; West et al. 2004; Brodie & Strader 2006; Gratton et al. 2012; Kruijssen 2014; Forbes et al. 2018)~~, consensus on an exhaustive picture of the formation of GCs is yet to be reached. Most galaxies host a GCS with a bimodal colour distribution that may reflect a bimodal metallicity distribution which is argued to

arise from two distinct formation channels (Brodie & Strader 2006). GCs are typically referred to as ‘blue’ or ‘red’ (e.g. Zinn 1985; Gebhardt & Kissler-Patig 1999; Larsen et al. 2001) where the former class is more metal-poor, more radially extended, and showing little to no rotation whereas the later is more metal-rich, radially concentrated, and rotating with the galaxy.

The literature offers a wealth of formation scenarios, many of which focus on the distinct characteristics of the blue or red subpopulation. Peebles & Dicke (1968) and Peebles (1984) argue that GCs form as the earliest bound structures in the Universe (i.e. prior to formation of the main galaxy), noting that the Jeans length and mass shortly after recombination is consistent with typical GC masses and sizes. Diemand et al. (2005) and Boley et al. (2009) argue that ~~(the bluesubpopulations of) GCs the blue, metal-poor GCs~~ ($[\text{Fe}/\text{H}] < -1$) form in radially biased dark matter (DM) halos at high redshift. Other recent hypotheses of GC formation prior to collapse of the proto-galaxy include formation in colliding supershells (Recchi et al. 2017), in supersonically induced gaseous objects

* E-mail: Halbesma@MPA-Garching.MPG.DE

2 T. L. R. Halbesma et al.

(Chiou et al. 2019), or in high-speed collisions of dark matter sub-haloes (Madau et al. 2019). Boylan-Kolchin (2017) suggests that (the blue subpopulation of) GCs could form in high density regions along the cosmic filament before or during collapse.

Other models also date GCs formation during formation of the proto-galaxy itself, for example as a result of thermal instabilities in hot gas-rich haloes (Fall & Rees 1985, also see the discussions in Kang et al. 1990). Alternative formation triggers are also explored, such as (other causes of) shock compression, or cloud-cloud collisions (e.g. Gunn 1980; Murray & Lin 1992; Harris & Pudritz 1994; Vietri & Pesce 1995; Larson 1996; Cen 2001).

Yet another hypothesis is that star cluster formation is triggered by major gas-rich (spiral) galaxy mergers (Schweizer 1987; Ashman & Zepf 1992), which is naturally expected within the framework of hierarchical assembly. One testable prediction of this scenario is the formation of young clusters in interacting and merging galaxies, which has been observed and are found to show remarkable similarities with GCs in the MW–Milky Way (MW) (e.g. Whitmore & Schweizer 1995; Holtzman et al. 1996; Zepf et al. 1999; Whitmore et al. 1999). Moreover, modelling efforts of this framework yield GC (sub)populations consistent with various observables (e.g. Muratov & Gnedin 2010; Choksi et al. 2018), and the recent numerical simulation of an isolated dwarf-dwarf merger executed at very high resolution (baryonic mass $m_b \sim 4M_\odot$; softening $\epsilon = 0.1$ pc) produces star clusters that could be GC progenitors (Lahén et al. 2019).

As for the formation timeline, the scenarios (and flavours thereof) of Fall & Rees (1985) and Ashman & Zepf (1992) are intertwined because accretion and mergers ~~continuously occurs~~ continually occur during the hierarchical build-up of galaxies. Various (other) hierarchical formation channels thus combine different aspects of the aforementioned paradigms, such as GC formation in (small) galactic disks before they are accreted onto an assembling galaxy (e.g. Côté et al. 2000, 2002; Beasley et al. 2002; Gnedin 2003). We refer to Gnedin et al. (2001) for a scoreboard of GC formation models compared to observations of the MW GCSs.

An idea that has recently been studied extensively is the hypothesis that the formation mechanisms of young massive clusters (YMCs, see Portegies Zwart et al. 2010, for a review) and GCs are the same. The proximity of YMCs in the local Universe allows to study these processes with a level of detail that cannot be achieved for GCs at high redshifts. In this framework, differences between ~~both the two~~ classes of objects are caused by nearly a Hubble time of (dynamical) evolution (e.g. Spitzer 1987). This picture is based on observed similarities between YMCs and GCs (e.g. Holtzman et al. 1992; Whitmore et al. 1999) and strengthened by observations of gravitationally lensed objects at high redshifts ($z = 2 - 6$). These sources have properties reminiscent of (local) YMCs and may be GC progenitors at times of formation (Vanzella et al. 2017; Johnson et al. 2017). The modelling work by Kruijssen et al. (2011, 2012); Kruijssen (2015) is now incorporated into cosmological zoom simulations and shows promising results (Pfeffer et al. 2018; Kruijssen et al. 2019).

The (mass) resolution of cosmological zoom simulations has reached the mass range populated by GCs, and the gravitational force softening can be as low several parsec. A number of groups can thus incorporate formation of (globular) star clusters into their high-resolution hydrodynamical simulations. Ricotti et al. (2016) ran parsec scale simulations of the high-redshift universe prior to reionization (the simulations stop at $z = 9$), Li et al. (2017) implement a new subgrid model for star (cluster) formation and run simulations that reach $z = 3.3$, and the run of Renaud et al. (2017)

reaches $z = 0.5$. Kim et al. (2018) find that mergers can push gas to high density that quickly forms clustered stars that end up tightly bound by the end of the simulation. A somewhat different approach couples semi-analytical models to DM-only simulations (Muratov & Gnedin 2010; Li & Gnedin 2014; Choksi et al. 2018; Choksi & Gnedin 2019b,a), ~~as-is-also~~ done in the work by El-Badry et al. (2019).

In this work we use state of the art simulations that produce realistic spiral galaxies (MW analogues) at redshift zero for which several global properties are consistent with the observations. The question thus naturally arises whether the star formation histories of the simulations could give rise to a GCS similar to the MW and/or Andromeda (M31) GCSs. In particular, we use the Auriga simulations (Grand et al. 2017, hereafter G17), further described in See Section 2. We investigate whether the star formation model implemented produces metallicity, radial, and metallicity-radial distributions that are consistent with the MW and/or M31 GCSs, and whether the model produces enough stellar mass with the right properties ~~to allow for formation efficiencies lower than unity and the expected (dynamical) mass loss over the cluster lifetime (not modelled explicitly in this study)~~. Moreover, the simulations provide us with the history of the star particles, which allows us to investigate the differences between these properties for sub-populations that have formed in situ versus those that have been accreted. We focus on the metallicity and radial distributions because observations of the red and blue GCs show distinct differences between these properties.

The plan of the paper is as follows. We summarise the relevant characteristics of the Auriga simulations in section Section 2, followed by a summary of the observations of the MW and M31 GCSs in section Section 3 that we ~~use to~~ compare our simulations to in section Section 4. We discuss our findings in section Section 5 to come to our conclusions in section Section 6.

2 THE AURIGA SIMULATIONS

We use the Auriga simulations (G17), a suite of high-resolution cosmological zoom simulations of MW analogues. The MW-mass selected initial conditions in the range $1 - 2 \times 10^{12} M_\odot$ were selected from the EAGLE ~~DM-only~~ DM-only box, see G17 for details. The simulations are performed with the AREPO code (Springel 2010; Pakmor et al. 2016) that solves the magnetohydrodynamical equations on a moving mesh.

The galaxy formation model includes primordial and metal-line cooling with self-shielding corrections. Reionization is completed at redshift ~~six-6~~ by a time-varying spatially uniform UV background (Faucher-Giguère et al. 2009; Vogelsberger et al. 2013). The interstellar medium (ISM) is described by an equation of state for a two-phase medium in pressure equilibrium (Springel & Hernquist 2003): cold clouds embedded in a hot ambient medium, whose mass and energy densities are described by processes important for star formation, such as the condensation of cold clouds and the photoevaporation of clouds. Stars form ~~stoeastically~~ stochastically in thermally unstable gas with a density threshold of $n = 0.13 \text{ cm}^{-3}$, and consecutive stellar evolution is accounted for. The channel for energetic stellar feedback is from non-local SNH type II supernovae (SN), which is modelled with wind particles (Marinacci et al. 2014; G17). Galactic winds enrich the ISM with metals from SNIa, SNII, and AGB-asymptotic giant branch stars (Vogelsberger et al. 2013). Supermassive black holes grow via Bondi–Hoyle–Lyttleton accretion (Bondi & Hoyle 1944; Bondi 1952), and feedback from the active galactic nuclei has a low-accretion and high-accretion mode

(radio and quasar) (Springel et al. 2005; Marinacci et al. 2014; G17). Finally, the simulations follow the evolution of a magnetic field of 10^{-14} (comoving) G seeded at $z = 127$ (Pakmor & Springel 2013; Pakmor et al. 2014).

This model was tailored to the AREPO code and calibrated to reproduce key observables of galaxies, such as the history of the cosmic star formation rate density, the stellar mass to halo mass relation, and galaxy luminosity functions. The galaxy formation model produces realistic spiral galaxies at redshift zero that match observations such as sizes, morphologies (G17), HI disc properties (Marinacci et al. 2017), and the diversity of accreted stellar halo properties (Monachesi et al. 2019). Furthermore, the simulations are able to reproduce major Galactic components, such as the chemical thin/thick disc dichotomy (Grand et al. 2018) and the recently discovered *Gaia* Sausage (Fattahi et al. 2019). They are thus a large set of detailed simulations similar enough to the Galaxy to make predictions for the GC population properties of the MW and M31.

The Auriga suite has a fiducial baryonic mass resolution $m_b = 5 \times 10^4 M_\odot$, with gravitational softening of collisionless particles $\epsilon = 369$ pc. Simulations at this resolution are referred to as level 4 (L4). Selected initial condition runs are accompanied by the lower (higher) resolution level L5 (L3). The resolution for these levels is $m_b = 4 \times 10^5 M_\odot$ with $\epsilon = 738$ pc, and $m_b = 6 \times 10^3 M_\odot$ with $\epsilon = 184$ pc, respectively. The mass resolution of the Auriga simulations is thus close $10^5 M_\odot$ to the characteristic peak mass of the lognormal GC mass distribution of $10^5 M_\odot$ (Harris 1991), although the gravitational softening is two orders of magnitudes larger than typical GC radii of several parsec. High-density gaseous regions are thus not expected to produce surviving stellar clumps with masses and radii consistent with GCs because such objects would numerically disperse, even in the highest-resolution runs. On the other hand, we can investigate (statistical) properties of age-selected GC candidates because each star particle represents a single stellar population with a total mass that could be consistent with one GC. This means that their formation sites and that of a could be consistent with those of real-world GC may be consistent GCs.

3 OBSERVATIONAL DATA

We describe the observations of the MW GCS in See. 3.1 Section 3.1 for these clusters, and of the M31 GCS in See. 3.2 Section 3.2. We discuss age-estimates in Section 3.3, their distribution of galactocentric radius in Section 3.4, and their total mass in 2D bins of metallicity-galactocentric radius in Section 3.5. We focus on both metallicity and galactocentric radius because these data are available in the observations and in the Auriga simulations.

3.1 Milky Way

Harris (1996, 2010 edition; hereafter H96e10) provides a catalogue¹ of the MW GCS that contains properties of 157 GCs. The authors initially estimated the number of GCs in the MW GCS to be 180 ± 10 , thus, their catalogue to be $\sim 85\%$ complete. However, an additional 59 GCs are claimed to GC candidates have been discovered more recently by various authors. The total number of GCs in the MW might add up to thus be 216 with recent estimates now

anticipating an additional thirty-30 GCs yet to be discovered (e.g. Ryu & Lee 2018, and references therein). We still use data from the Harris catalogue, but caution that it may (only) be 50-70% complete. Specifically, the relevant data fields that we use from H96e10 are the metallicity [Fe/H], the Galactic distance components X , Y , and Z (in kpc)², and absolute magnitude in the V band, M_V . We use the V band magnitude to calculate mass-estimates by assuming $M_{V,\odot} = 4.83$ and a mass to light ratio $M/L_V = 1.7 M_\odot/L_\odot$, the mean for MW clusters (McLaughlin & van der Marel 2005). We supplement the catalogue with age-estimates from isochrone fits to stars near the main-sequence turnoff in 55 GCs (VandenBerg et al. 2013, hereafter V13).

3.2 Andromeda

The fifth revision of the revised Bologna catalogue (RBC 5, last updated August, 2012) is the latest edition of three decades of systematically collecting integrated properties of the GCS of M31 (Galletti et al. 2004, and references therein). One contribution to RBC 5 is the work by Caldwell et al. (2011, hereafter C11), subsequently updated by Caldwell & Romanowsky (2016, hereafter CR16).

C11 and CR16 Caldwell et al. (2011, hereafter C11) and Caldwell & Romanowsky (2016, hereafter CR16) present a uniform set of spectroscopic observations calibrated on the MW GCS of the inner 1.6° (~ 21) kpc of M31 that is believed to be 94% complete. GCs in In addition, the outer stellar halo of M31 up to $r_{\text{proj}} \sim 150$ kpc, are is observed in the Pan-Andromeda Archaeological Survey (PAndAS, Huxor et al. 2014, hereafter H14), but see also Veljanoski et al. (2014) and Mackey et al. (2019). H14 presents the discovery of 59 new GCs and publishes updates to RBC 5. The (PAndAS, Huxor et al. 2014; Veljanoski et al. 2014; Mackey et al. 2019, hereafter). In fact, the work of H14 is incorporated in the latest public release³ of the C11 dataset, further revised by and CR16. We use data sets. Therefore we use the latest data set of CR16 because it is the most recent aggregated dataset of M31's GCS that contains properties of interest for our study and it includes GCs in the inner region and in the outer halo. The relevant fields in the CR16 dataset that we use are the age, metallicity, and the mass-estimate⁴. Radii are calculated from We calculate the galactocentric radii from the observed positions RA and DEC, as further discussed in See. Section 3.4.

3.3 Age-estimates

The top panel of Figure 1 shows a histogram of the age-estimates of the 55 MW GCs in V13 and 88 GCs in M31 for which age-estimates are available in CR16. The mean age of the MW GCs in this data set is $11.9 \pm 0.11.9$ Gyr and the dispersion standard deviation is 0.8 Gyr. Furthermore, only one-1 of the 55 GC age-estimates is below 10 Gyr. The M31 GCS has a mean age of $11.0 \pm 0.211.0$ Gyr with a dispersion standard deviation of 2.2 Gyr, and 24 GCs have

² In a Sun-centered coordinate system: X points toward Galactic center, Y in direction of Galactic rotation, and Z toward the North Galactic Pole. We calculate the galactocentric radius $r_{\text{gal}} = \sqrt{(X - r_\odot)^2 + Y^2 + Z^2}$, assuming the solar radius $r_\odot = 8$ kpc.

³ Last revised 23 Sep 2015, see https://www.cfa.harvard.edu/oir/eg/m31clusters/M31_Hectospec.html

⁴ The authors assumed $M/L_V = 2$ independent of [Fe/H]

¹ See https://www.physics.mcmaster.ca/Fac_Harris/mwgc.dat

4 T. L. R. Halbesma et al.

age-estimates below 10 Gyr with a minimum age of 4.8 Gyr. Based on these data, we find that the age distributions of the MW and M31 GCSs are not statistically consistent. Moreover, the MW appears to host a GCS that is somewhat older than that of M31. This is somewhat surprising given that M31 is generally considered to be earlier type than the MW meaning so that an older stellar population would be naively expected. However, we do caution that both data sets are incomplete and the age measurements have large uncertainties (of 1 – 2 Gyr). On the other hand, the magnitude of the uncertainty is insufficient to explain the low-age tail in M31.

3.4 Radial distribution

The bottom panel of Figure 1 shows the radial distribution of the MW and M31 GCS. We divide the radii of M31 by its virial radius⁵ $r_{\text{vir},\text{M31}} = 299$ kpc and multiply by $r_{\text{vir},\text{MW}} = 261$ kpc to account for the different intrinsic sizes of both galaxies. Values for $R_{\text{gal}} - r_{\text{gal}}$ are readily available in H96e10 (assuming $r_{\odot} = 8.0$ kpc), but the galactocentric radius of GCs in M31 is not available in CR16. Therefore, we follow Wang et al. (2019, Sec. 4.1) Wang et al. (2019, section 4.1) to calculate the projected radius r_{proj} from the observed positions, adopting M31's central position from the NASA Extragalactic Database⁶ (α_0, δ_0) = (0^h42^m44.35^s, +41°16′08.63″) and the distance $D_{\text{M31}} = 780$ kpc (McConnachie et al. 2005; Conn et al. 2012). We calculate estimate r_{gal} as ‘average deprojected distance’ $r_{\text{gal}} = r_{\text{proj}} \times (4/\pi)$.

The solid lines show the distributions using all available data (because the sky coordinates are known for each GC), while the subset for which age-estimates are available is indicated using dotted lines. The latter shows a narrower range of radii than the full data set: it appears that few age-estimates are available for the innermost (< 1 kpc) GCs, and none for those beyond roughly twenty–20 kpc (the halo GCs). We compare both the two distributions of the full data set (solid lines) and find that the MW has more GCs in the range 1 – 4 kpc than M31 (when accounting for the larger number of total GCs in M31). Interestingly, both the two distributions show a similar trend for $r_{\text{gal}} > 4$ kpc and host a subpopulation of halo GCs. However, the two radial distributions. Our naive expectation would be that the spiral galaxies in the Local Group host similar GC systems. However, based on a two-sample KS test we find that the radial distributions of the GCS of the MW and M31 are not statistically consistent with being drawn from the same underlying distribution due to substantial differences at intermediate radii.

3.5 Total GC mass in metallicity-radial space

We show the two-dimensional mass-weighted metallicity-radial metallicity-radius distribution of the MW (and the M31) GCS in the top (bottom) and bottom panel of Figure 2, respectively. Note that the number of GCs is different than in Figure 1 because here we plot [Fe/H] between -2.5 and 0 and r_{gal} between 1 and 250 kpc whereas Figure 1 shows the full range of metallicities and radii.

The observations indicate that no very few GCs with high




Figure 1. Top: Age distribution of 55 GCs in the MW (data from VandenBerg et al. 2013) and 88 GCs in M31 (data from Caldwell & Romanowsky 2016). Bottom: Distribution of galactocentric radius in the MW and M31. The dotted lines show the subset of data that also have age measurements (i.e. the same sample as in the top panel). We divide the radii of M31 by $r_{\text{vir},\text{M31}} = 299$ kpc and multiply by $r_{\text{vir},\text{MW}} = 261$ kpc (Patel et al. 2017) to compensate for the different intrinsic sizes of both galaxies.

metallicities are to be expected at large radii (the three bins in the upper right corner, both for MW and for M31), and relatively few GCs at large radii in general ($r_{\text{gal}} > 30$ kpc; right column: 11 GCs or 7.3% in the MW and 17 or 4.6% in M31). Moreover, the M31 hosts more metal-rich ([Fe/H] > -1) GCs in each radial bin in comparison to the MW GCSs than what would be expected when after accounting for the fact that M31 hosts a larger GC than the MW. Finally, given that both the marginalized (i.e. the metallicity and radial) distributions are not statistically consistent between the two galaxies, we find that the two-dimensional distributions are also not consistent. More generally, the GCS of the MW differs significantly from that of M31, not only in the number of clusters but also in their distribution over radius and metallicity. We compare these observations to the Auriga simulations later on in Sec. in Section 4.3.

⁵ r_{vir} is the radius in a spherical ‘top-hat’ perturbation model at which the average density reaches an overdensity of $\Delta_{\text{vir}} = 357$ times the background density, adopting the values and cosmology from Patel et al. (2017)

⁶ <https://ned.ipac.caltech.edu/>




Figure 2. *Top:* Mass-weighted r_{gal} - $[\text{Fe}/\text{H}]$ distribution of 151 GCs in the MW (data from Harris 1996, 2010 ed.), color-coded by the median (values also shown in each bin). The numbers in parenthesis show how many star particles fall within the bin. *Bottom:* Same for M31, showing the 366 GCs in CR16 (data from Caldwell & Romanowsky 2016). Radii of M31 are again divided by $r_{\text{vir},\text{M31}}$ and multiplied by $r_{\text{vir},\text{MW}}$. Note that the range of the colourmap differs in both figures.

4 RESULTS

We define GC candidates in the Auriga simulations as all star particles older than 10 Gyr based on the age distribution of the MW GCS (top panel of Figure 1), and following Renaud et al. (2017), adopting the definition of a GC from Portegies Zwart et al. (2010) and following the method of Renaud et al. (2017). However, we caution that his definition, although statistically consistent with the estimated ages of all GCs in the MW, is not consistent with the tail of younger clusters as stated in Section 3.3, and further discussed in Section 5.3.

Throughout our analysis we compare distributions of age-selected GC candidates (age > 10 Gyr) with those of all star particles (i.e. when no additional selection criterion is applied) with no age cut. The mean number of star particles in the Auriga L4 simulations that we select as GC candidates is $3.1 \cdot 10^5$ (with a mean total mass of $1.15 \cdot 10^{10} M_{\odot}$), and the mean number of all star particles is $1.93 \cdot 10^6$ ($7.3 \cdot 10^{10} M_{\odot}$). The subset of GC candidates is thus

around 15% of all the stars in the galaxy when averaged over all Auriga L4 haloes. We further split the GC candidates into an *in situ* subset (defined as GC candidates that are bound to the most-massive halo/subhalo in the first snapshot that the particle was recorded), and an accreted subpopulation (those that have formed *ex situ* and but are bound to the most-massive halo/subhalo at $z = 0$). A GC candidate either formed *in situ* or it was accreted. We consider the metallicity distribution in See:Section 4.1, the distribution of galactocentric radii in See:Section 4.2, and the combination of both in See:the two in Section 4.3.

4.1 Metallicity distribution

We investigate whether the star formation model implemented in Auriga produces metallicity distributions consistent with the MW and M31 GCSs, and whether the simulations generate sufficient total mass at metallicities typical for the MW and M31 GCSs. To visually inspect the former we show the normalized metallicity distribution of three specific Auriga galaxies in Figure 3 in comparison to the MW and M31 $[\text{Fe}/\text{H}]$ distributions.

We select Au4-10⁷, Au4-21, and Au4-4 to show the distributions of three individual simulation runs that we consider representative for specific behaviour, and to highlight that different runs could give rise to different distributions. We show the age-selected GC candidates as well as the accreted and *in situ* subsets. The top half of the left figure panel of Figure 3 shows the MW and M31 GCSs, where we overplot a double Gaussian for the MW GCS by adopting literature values of the mean μ and standard deviation σ of the metal-rich and metal-poor subpopulations (Harris 2001).

We find that the age cut, on average, lowers the mean metallicity $\langle [\text{Fe}/\text{H}] \rangle$ from 0.0 to -0.6 . This reflects the enrichment history of the galaxy because the age cut removes the more metal-rich stellar population which shifts the mean metallicity towards lower values. Furthermore, accreted GC candidates generally have lower mean metallicities than the *in situ* subset with differences of roughly ~ 0.3 dex for the majority of the simulation runs. This behaviour can be seen in Au4-21, while a slightly larger difference of ~ 0.5 dex is seen for Au4-10 (as plotted), Au4-16, Au4-17, Au4-18 and Au4-22. However, this trend is reversed for Au4-1 and Au4-4 for which the *in situ* subset has a lower mean metallicity instead. Au4-1 is undergoing a major merger at redshift zero and we find a mean metallicity in this simulation $\mu = -1.51$ (-0.74) for the *in situ* (accreted) GC candidates, although the former consists of only 1019 particles (1.3% of all GC candidates, and with a total mass of $5 \times 10^7 M_{\odot}$). For Au4-4, only 10.8% of the GC candidates is classified as *in situ* (compared to *in situ* fractions of 40 – 80% for other haloes). We note that Au4-4 has undergone a major merger around 2 Gyr lookback time, which is probably why the accreted population is more metal-rich. After inspection of the same figure for every one of the thirty-30 Auriga L4 haloes we find that the simulations produce (sub)populations of GC candidates that are more metal-rich than the MW and M31 GCSs. Moreover, none of the simulations produces GC candidates with a bimodal metallicity distribution.

We show the mean metallicity and standard deviation of all thirty-30 Auriga L4 haloes in Figure 4 to illustrate the individual behaviour for the full set of Auriga galaxies. The green-beige crosses are to be compared to the purple (magenta) cross, which shows

⁷ The nomenclature is ‘Au’ for Auriga, followed by the resolution level (4) and halo number (10, indicating which set of initial conditions was used).


6 *T. L. R. Halbesma et al.*

Figure 3. *Top:* Metallicity distribution of the MW and M31 GCS. The solid purple (magenta) line shows the GCSs of the MW (M31), and we indicate the mean value with an arrow at the bottom of the x-axis. We overplot a double Gaussian for the MW GCS using purple dotted lines, taking the mean and dispersion from (Harris 2001). *Other panels:* three selected Auriga haloes as indicated in the upper right corner. In each of these three panels we show the GC candidates in green (blue) dotted line for all star particles. We split the GC candidates up into two subpopulations, those that have formed *in situ* (blue), and those that have been accreted (red). The numbers in parenthesis show how many star particles are included in the subset.

and magenta crosses, which show the mean value of all MW (and M31) GCs, respectively. In addition, we show the metal-rich (metal-poor) population of the MW GCS using a solid (open) dot, and the metal-poor population with an open dot. We caution that these literature values result from individual Gaussian fits to subsets of the observational data cut at $[\text{Fe}/\text{H}] = -1$. We do not include corresponding data points calculated using such an artificial cut for the simulated star particles. Having said that, we do find it interesting that the mean metallicities of the *in situ* GC candidates appear roughly consistent with that of the metal-rich population of the MW despite the lack of such a metallicity cut, although the simulations show larger dispersions. The latter could simply be caused by the hard separation of the MW data into two groups, which means the range is smaller and the resulting dispersion lower. With regard to the M31 GCS, so far no definitive consensus has been reached in the literature concerning uni- bi- or trimodality in the $[\text{Fe}/\text{H}]$ distribution, but CR16 argues that the data, after removal of




Figure 4. First vs. second central moment of the Auriga L4 metallicity distributions. Each cross (for a given colour) represents one simulation. The green (blue) crosses show the values calculated using all (only *in situ*) [only accreted] GC candidates. Green (blue) triangles indicate that all stars were used. The purple (magenta) cross denotes our calculation using all MW (M31) observations (which would be appropriate for a unimodal distribution). The black (purple) solid (open) dots indicate the literature values of a bimodal Gaussian fit to the data (values from Ashman & Zepf 1998, p. 38) (values from Ashman & Zepf 1998, page 38), showing the metal-rich (metal-poor) component of the MW (separated at $[\text{Fe}/\text{H}] \approx -1$ as indicated by the purple dotted vertical line).

younger objects due to improved age classification, hints at three populations separated at $[\text{Fe}/\text{H}] = -0.4$ and $[\text{Fe}/\text{H}] = 1.5$. None of the Auriga sub-sets has a mean value that offers much hope that of reconciling the simulated distribution is consistent with the lowest metallicity group in the M31 GCS. This is also true for the metal-poor population of the MW with a mean value of -1.6 . The main take-away from this plot, again, result of this figure is that all Auriga L4 galaxies have metallicity distributions with (much) larger mean values than what is observed for the MW and M31 GCS, and that we systematically find lower mean metallicities for the accreted GC candidates than for those that have formed *in situ*.

We now turn to our second goal, which is to see whether sufficient total mass in GC candidates is produced by the Auriga simulations. To answer this question we plot a mass-weighted metallicity distribution in Figure 5. We show the median for all thirty over all the 30 Auriga L4 haloes with the 1σ interval around it (and the 25-75 percentile range as shaded regions, which shows the scatter between runs that have different initial conditions, thus, have unique merger histories) varying merger histories. We chose to aggregate the data to indicate general trends that we find when the GC candidates are split up into the accreted and *in situ* subsets, rather than selecting typical examples of individual (simulated) galaxies. Once again we notice that the peak metallicity shifts down from $[\text{Fe}/\text{H}] = 0$ to $[\text{Fe}/\text{H}] = -0.6$ for GC candidates compared to all stars, and we find that the mass at the peak lowers by roughly one dex. The mass budget of the GC candidates is dominated by the *in situ* subpopulation above $[\text{Fe}/\text{H}] = -1$, and by the accreted subset below this value. We show the MW (M31) GCS in purple (magenta) and that of M31 in magenta, and notice that the difference between the MW and M31 distributions is substantially larger than the scatter between different Auriga galaxies, particularly around $[\text{Fe}/\text{H}] = -1$. In addition, M31




Figure 5. Mass-weighted metallicity distribution of star particles in the Auriga L4 simulations. We show the median value of all Auriga haloes for all stars (green-beige dotted) and GC candidates (i.e. stars with age > 10 Gyr; green-beige solid). The latter sub-set is further split up between stars that formed *in situ* (blue-brown solid), and those that were accreted (red-teal solid). Shaded regions indicate the 1-sigma interval and 25-75 percentile range. The MW (M31) GCS is shown in purple (magenta). The middle (bottom) panel shows the ratio of the simulated mass to the mass in the MW (M31) GCS.

does host (a handful of) GCs with $[\text{Fe}/\text{H}] < -2.5$ as well as GCs with $[\text{Fe}/\text{H}] > 0$ while the MW does not.

We show the ratio of the simulated to the observed profiles in the middle (bottom) panel and bottom panels. This mass excess can be thought of the ‘mass budget’ that the Auriga GC candidates can ‘afford to lose’ (due to a combination of smaller than unity bound cluster formation efficiencies combined with a Hubble time of dynamical evolution), while still producing sufficient mass at the right metallicities. In particular, the cluster formation combined efficiency would have to decrease-increase with decreasing metallicity for Auriga GC candidates to produce a population of GC candidates that is consistent with the MW. For the GC candidates in M31 we find a constant mass ratio up to $[\text{Fe}/\text{H}] = -0.9$, above which the simulations produce a higher mass ratio with increasing metallicity. If dynamical evolution is not expected to more efficiently disrupt GCs of higher metallicity, then we would find that the efficiency to form bound star clusters would have to decrease with increasing metallicity. This trend could be consistent with the cluster formation efficiency model of Kruijssen (2012) as lower metallicity star particles formed at earlier times when the bound




Figure 6. Mean-Median and standard deviation of the logarithm of radial distribution of star particles in each of the thirty Auriga L4 haloes compared to the MW (M31) GCS shown in purple (magenta). The Auriga (M31) radii are divided by $r_{\text{vir,Au}}$ ($r_{\text{vir,M31}}$) and multiplied by $r_{\text{vir,MW}}$ to compensate for different intrinsic sizes of the galaxies.

cluster formation efficiencies are higher due to higher pressure birth environments.

4.2 Radial distribution

We repeat the analysis of the previous subsection for the distribution of galactocentric radius instead of metallicity. We divide the Auriga radii by $r_{\text{vir,Au}}$ and multiply by $r_{\text{vir,MW}}$ to compensate for the different intrinsic sizes of the Auriga galaxies. We look for general trends present in all Auriga L4 galaxies. Figure ?? shows the mean-6 shows the median and standard deviation of $\log_{10}(r_{\text{gal}})$ of star particles in all Auriga L4 simulations, respectively. The purple (magenta) cross again shows the mean and magenta cross show the median of the MW (and M31) GCs (GCs, which are to be compared to the green crosses), but solid (open) beige crosses. The solid dot now shows our calculation of the mean-median value of the ‘red’ metal-rich (population, and the open dot that of the ‘blue’ metal-poor) population. We split the radii up into the metal-rich/metal-poor groups by taking a metallicity cut at $[\text{Fe}/\text{H}] = -1$ as is done in the literature.

We notice that the *in situ* GC candidates are more centrally distributed, whereas the accreted subsets have a larger radial extent. This is not surprising because the classification of *in situ* requires star particles to have formed within the virial radius, thus they could naturally be expected to end up at small galactocentric radii. Accreted star particles, on the other hand, have formed in another (sub)halo beyond the virial radius, thus, they would first have to migrate inwards in order to populate the innermost radii. Moreover, we find that the simulations have a wider dispersion in $\log(r_{\text{gal}})$ than the MW GCS, while the dispersion of M31 seems to lie within the range of dispersions found in the Auriga galaxies. Furthermore, the mean-median of the MW GCS lies roughly within the range of values produced by the Auriga simulations, whereas the mean-median of M31 is somewhat larger and slightly closer to typical mean-median values of the accreted GC candidates in the Auriga simulations. The larger radial extent of the M31 GCS is generally believed to hint at a more accretion-dominated origin of the GCS,

8 *T. L. R. Halbesma et al.*




Figure 7. Mass-weighted radial distribution of star particles in the Auriga L4 simulations, where the Auriga (M31) radii are again divided by $r_{\text{vir,Au}}$ ($r_{\text{vir,M31}}$) and multiplied by $r_{\text{vir,MW}}$. We show the median value of all Auriga haloes for all stars (green-beige dotted) and GC candidates (i.e. stars with age > 10 Gyr; green-beige solid). The latter sub-set is further split up between stars that formed *in situ* (blue-brown solid), and those that were accreted (red-teal solid). Shaded regions indicate the 25-75 percentile range. The MW (M31) GCS is shown in purple (magenta). The middle (bottom) panel shows the ratio of the simulated mass to the mass in the MW (M31) GCS.

and may reflect a richer accretion history of M31 in comparison to the MW.

Figure 7 shows the mass-weighted radial distribution of the Auriga L4 haloes. We notice a subtle peak around 10 kpc for all star particles that is not present for the GC candidates, indicating that the stellar disc is no longer present when applying the latter selection criterion. Furthermore, we find that the dominant contribution to the total mass in GC candidates changes from those formed *in situ* to the accreted population around 10 kpc. Again we show the mass ratio of the simulations compared to the MW and M31 GCS. We find a decreasing mass ratio with increasing radius in the range 0.2 to ~ 5 kpc, followed by an increase attributed to the accreted sub-population. For the MW we notice that significantly fewer GCs are found beyond 40 kpc than for M31, and that accreted GC candidates contribute mostly to the Auriga GCS at these radii. We further investigate a breakdown of the total mass in Auriga GC candidates into bins of both metallicity and radius in the following section.




Figure 8. Mass-weighted $[\text{Fe}/\text{H}]$ - r_{gal} distribution of all thirty Auriga L4 haloes. Here we consider only the GC candidates and color-code by the median mass (values also shown in each bin). The numbers in parenthesis show how many star particles fall within the bin. Note that the range of the colourmap differs from both panels in Figure 2 (for improved contrast within the plot).

4.3 Total mass in metallicity-radial space

We investigate whether the Auriga simulations still produce sufficient mass when the GC candidates are two-dimensionally binned in $[\text{Fe}/\text{H}]$ and r_{gal} . First we sum the total mass of GC candidates in each bin (for an individual Auriga simulation), then we calculate the median total mass in each bin over all thirty Auriga L4 haloes. The result is shown in Figure 8, which can be compared to the observed total GC mass shown in Figure 2. We notice that the simulations produce relatively little mass in metal-poor GC candidates at small radii and in metal-rich GC candidates at large radii. However, there is still few times $10^7 M_{\odot}$ produced by the simulations in the uppermost 3 bins where no observed GCs are found.

To ease this comparison between the simulations and observations we take the ratio of the total mass of GC candidates in Auriga to the total GC mass in the MW (and M31) observations. We use a diverging colour map centered around the overall median of the mass ratio (i.e. we subtract the median of these twenty-five bins from the value in each bin). We show this in the top (bottom) and bottom panel of Figure 9 for the MW and M31, respectively. Note that the data in the three top-right bins is masked (3 bins in the top right are empty (white)) because the mass ratio cannot be calculated due to zero mass in the observations (therefore the bins are white). If the product of the bound cluster formation efficiency and consecutive GC disruption over a Hubble time would be constant, then the value in each bin would have been zero. The red and blue bins indicate deviations from such a scenario and indicate a higher and lower than average mass excess in the simulations. The median value used to shift the mass ratio is 2.68 and 1.69 dex for the MW and M31, respectively. This means that the median ‘mass budget’ that the simulations ‘can afford to lose’ is roughly a factor 480 and 50 for the MW and M31, respectively. The lowest mass surplus for both galaxies is the bin where $-2.5 < [\text{Fe}/\text{H}] < -1.5$ and $3 < r_{\text{gal}}/\text{kpc} < 8$ in $[-2.5, -1.5]$ and galactocentric radius $r_{\text{gal}} \in [3, 8]$ kpc, for which the simulations pro-

duce a factor $10^{2.68}/10^{1.33} \approx 22$ and $10^{1.69}/10^{0.75} \approx 9$ more mass in GC candidates than ~~what~~ is observed in the MW and M31 respectively. The highest factor for the MW is 2.1×10^5 , and 1.1×10^3 for M31.

We now focus on four quadrants of this diagram, split at $[\text{Fe}/\text{H}] = -1$ (metal-poor or metal-rich) and $r_{\text{gal}} = 8$ kpc (inside or outside the Solar radius). The majority of the mass in Auriga GC candidates is produced in metal-rich star particles inside the Solar radius. The total mass is dominated by the *in situ* component, for $[\text{Fe}/\text{H}]$ as well as for r_{gal} as can be seen in Figure 5 and Figure 7. The total mass excess in this region increases with decreasing metallicity for the MW and M31, and there is a weak trend of decreasing over-production with increasing radius for the *in situ* subsets but no trend with radius for the accreted GC candidates. As for the metal-rich GC candidates outside the Solar radius, we find that the mass excess with respect to the median is largest in this corner for the MW and M31. The main reason for this trend are the lower GC counts in the observations, whereas the Auriga simulations efficiently form metal-rich GC candidates and have no problem to produce them at large radii. The opposite is true for the metal-poor GC candidates inside the Solar radius, where the mass surplus with respect to the median is smallest for both galaxies. However, the disruption rates are expected to be particularly high for GCs in the inner regions of galaxies that are on which must have orbits with pericentres close to the galactic bulge or bar. In addition, disk crossings in this region will cause spikes in the tidal tensor due to steep density gradients that could be highly disruptive to the GCs. The bins with the lowest ‘mass budget’ are in this region (a factor 9 for the MW and 22 for M31). The bound cluster formation efficiency has to exceed 10%, and none of the GC candidates in this region can experience significant mass loss in order for Auriga to produce sufficient mass in this quadrant. Finally, the metal-poor GC candidates outside the Solar radius have lower than average mass ratios, which is not surprising given that the Auriga simulations overall underproduce low-metallicity stars particles.

5 DISCUSSION

5.1 Metallicity distribution

GCs are ~ 0.5 dex more metal-poor than spheroid stars observed at the same radius for almost all galaxies (Harris 1991). Our selection function (age cut) does lower the mean metallicity of GC candidates by 0.5 dex with respect to all star particles, but we still find that the GC candidates in the Auriga simulations are more metal-rich than the MW and M31 GCSs. Although the (old) star particles represent single stellar populations with a mass consistent with that of GCs, they are in fact statistical tracers of the stellar population of the galaxy as a whole. Therefore only a (small) fraction of the star particles may represent plausible formation sites of GCs, whereas the majority represents (halo) field stars - the disk component effectively falls outside our selection of star particles due to the age cut. It is implausible that all halo stars come from disrupted GCs because the total mass in GCSs at redshift zero is generally a few orders of magnitude lower than the total mass of stellar haloes of galaxies. If stellar haloes solely consist of disrupted GCs, then the GC disruption rate would have to be considerably higher than expected from theoretical models. Moreover, disrupted dwarf galaxies are more likely to contribute to the build up of the stellar halo than GCs (e.g. Deason et al. 2015; Conroy et al. 2019). Overall, our set of GC candidates most likely contains an overabundance of regular




Figure 9. The top (bottom) panel shows the logarithm of the ratio of simulated mass to mass in the MW (M31) GCSs, i.e. the logarithm of the mass ratio, minus the median value of these 25 bins. The values are 2.68 and 1.69 dex for the MW and M31, respectively. Red (blue) indicates a higher (lower) overall mass excess. The number in each bin shows the color-coded value that is plotted, and the (median) star particle count is given in parenthesis. Note that the three bins in the upper right corner are white because the observations have zero mass there.

stars. It could be that the resulting metallicity distribution is then biased towards higher values than ~~what~~ is observed for real GCs. An alternative explanation could overmixing of metals at early times.

In general, the colour (metallicity) distribution of most GCSs is appears to be bimodal with typical separating value $[\text{Fe}/\text{H}] \sim -1$ (Zinn 1985; Gebhardt & Kissler-Patig 1999; Larsen et al. 2001; Peng et al. 2006). Indeed, Ashman & Zepf (1998) and Harris (2001) find that the MW GCS has a bimodal $[\text{Fe}/\text{H}]$ distribution. Observations of GCs in M31, however, may be best split into three distinct metallicity groups (Caldwell & Romanowsky 2016). The numerical simulation of Renaud et al. (2017) does yield a bimodal metallicity distribution where the metal-poor population is dominated by accreted star particles and the metal-rich population by *in situ* stars.

However, we find that none of the Auriga simulations produces a bimodal metallicity distribution for age-selected GC candidates. Interestingly, the cross-over point above (below) which the mass-

weighted metallicity distribution of GC candidates is dominated by those that have formed *in situ* (were accreted) does coincide with the separation between the metal-rich and metal-poor populations of GCSs at $[\text{Fe}/\text{H}] = -1$, as for the MW. Moreover, we do find that the mean metallicity shifts when we split the GC candidates ~~up~~ according to birth location (accreted or *in situ*). In particular, the mean values of the *in situ* GC candidates are roughly consistent with the metal-rich MW GCs. On the other hand, we also notice that ~~a substantial number of simulations has similar mean values most of the simulations have a similar mean value~~ for all GC candidates, and the simulated *in situ* GC candidates have larger dispersions than ~~the is claimed for~~ metal-rich GCs ~~in the MW~~. Moreover, the offset between the ~~mean means~~ of the metal-rich and metal-poor populations in the MW is 1 dex, a factor 2 – 3 larger than the offset that we find between the *in situ* and accreted populations. ~~The Auriga simulations therefore do not support the view that the metal-poor GCs could have only formed in satellites that have later been accreted whereas the metal-rich GC subpopulation has only formed in situ. With this statement, however, we do~~ However, we caution that Kruijssen et al. (2019) suggests to reserve the ‘*ex situ*’ classification for accretion after $z = 2$, when the central galaxy has formed and accretion unambiguously contributes to the radially extended halo GC population. In fact, our classification might not hold for Au4-1.

Finally, we note that Brodie & Strader (2006) compares the number of metal-poor GCs to the stellar halo mass and find⁸ $T_{\text{blue}}^n \sim 100$, while the number of metal-rich GCs compared to the bulge mass yields $T_{\text{red}}^n \sim 5$, and therefore conclude that the formation efficiency of metal-poor GCs is ~~twenty~~ 20 times higher than the metal-rich GCs with respect to field stars. We find that a gradual increase in formation efficiency of GCs with respect to field stars would be required with decreasing metallicity for the GC candidates in the Auriga simulations to yield sufficient total mass to be consistent with the MW or M31 GCS. In addition, the metal-poor GC candidates would have to experience low mass ~~loss~~ ~~loss~~ rates, especially those within the Solar radius.

5.2 Radial distribution

The M31 GCS has a factor 2-3 more GCs at large radii compared to the MW GCS, which may indicate that M31 has a richer merger history than the MW (Caldwell & Romanowsky 2016). In the Auriga simulations we find that the GC candidates at radii larger than ~ 20 kpc are indeed dominated by accreted star particles. However, we select star particles that are bound to the main halo and main subhalo, which means that we include particles up to the virial radius r_{200} . The Auriga simulations have no problem populating the stellar halo up to the virial radius, even with our additional age cut. On the other hand, the MW and M31 have fewer GCs at large radii. We expect tidal disruption to be less efficient at larger radii, thus, the formation efficiencies of the accreted GC candidates would have to be lower ~~Alternatively, the fraction of old star particles in our subset of GC candidates may contain considerably more star particles that represent stars in the old stellar halo that have been accreted during hierarchical buildup of the galaxy.~~ Our classification ~~as of~~ GC candidates could ~~especially~~ be improved for the accreted subset ~~in particular~~. For example, we could ~~tag~~ ~~select~~ only a subset of the accreted star particles as GC candidates ~~where~~ where the number

number of GCs ~~is~~ ~~would be~~ determined by the satellite halo mass, given the $M_{\text{vir}} - N_{\text{gal}}$ correlation found by Burkert & Forbes (2019). ~~However, we did not include such an analysis in this study.~~

5.3 Lower age cut in the GC selection function

Our method is sensitive to the definition that we adopt to classify GC candidates (e.g. Forbes et al. 2018). Therefore we explore how our results change if we include all star particles older than 8 Gyr, and if we further lower the age cut to 6 Gyr. The average fraction of GC candidates that we select from the all the star particles increases to 35% and 55% for an age cut of 8 and 6 Gyr, respectively.

We find that including more recently formed star particles in our subset of GC candidates shifts the mean metallicity to higher values. Moreover, the total mass in metal-rich ($[\text{Fe}/\text{H}] > -1$) GC candidates increases for a lower age threshold and moves closer to the dotted line that shows the distribution of all star particles in Figure 5. However, we notice that the Auriga simulations, on average, do not form new stars with a metallicity below $[\text{Fe}/\text{H}] = -1$ more recent than 6 Gyr ago; the behaviour of Figure 5 does not change for metallicities below $[\text{Fe}/\text{H}] = -1$ when we change the age cut to 8 or 6 Gyr, and the slope of the mass ratio for metallicities above $[\text{Fe}/\text{H}] = -1$ is slightly steeper. In addition, we still find that the accreted GC candidates are more metal-poor than those that have formed *in situ*.

The qualitative behaviour of the median distribution of galactocentric radii (Figure 7) is unaffected by lowering the age cut other than including more star particles (more mass) which leads to a higher normalisation and a higher mass ratio with respect to the observed GC systems of the MW and M31. The qualitative behaviour presented in Figure 9 does not change when the age cut is lowered to 8 or 6 Gyr. GC candidates is not affected definition because our selection function mostly includes star particles that have formed before roughly $z = 2$. Overall, reducing the age cut worsens the agreement between the observed GCSs and the candidate GCs in the Auriga galaxies.

6 SUMMARY AND CONCLUSIONS

We investigate statistics of age-selected (> 10 Gyr) GC candidates in the Auriga simulations and compare the simulations to the MW and M31 GCSs ~~to test whether the star formation model implemented for the Auriga simulations could give rise to metallicity and radial distributions that are consistent with these two observed GC systems.~~ Based on our analysis we draw the following conclusions.

- The star formation model implemented in the Auriga simulations produces metallicity distributions that are more metal-rich than the MW and M31 GCSs. For most of the Auriga simulations we find that the subset of accreted GC candidates has a lower mean metallicity than the *in situ* subpopulation. However, the accreted subsets are still more metal-rich than the MW and M31 GCSs. Moreover, none of the simulated subsets has a mean metallicity as low as the mean of the ‘blue’ metal-poor component of the MW GCS.

- We find that the difference between the metallicity distributions of the MW and M31 GCSs is big compared to the scatter in the Auriga GC candidates of different Auriga runs. The radial distributions of the MW and M31 GCSs are somewhat more similar, but they are not consistent with being drawn from the same

⁸ The ~~quantity~~ ~~specific frequency~~ T is the number of GCs per $10^9 M_{\odot}$ of galaxy stellar mass.

distribution. The scatter between the radial distributions of different Auriga simulations is smaller than the difference between the MW and M31 GCSs.

- GC candidates in the Auriga simulations may be found out to r_{200} , given that our selection function of star particles selects all stars bound to the main subhalo ~~in the main of the~~ halo. The total mass of GC candidates is dominated by accreted star particles at radii beyond 20 kpc. The GCs in the MW and M31, on the other hand, have a much smaller radial extent.

- ~~Tidal forces are larger in the inner regions of galaxies than in the outer halo. Therefore, we expect dynamical evolution to more strongly affect GC candidates at small radii than at large radii. However, we find that the Auriga simulations produce more mass in GC candidates with high metallicities at large radii than with low metallicities at small radii.~~ The cluster formation efficiency would have to increase with decreasing metallicity for GC candidates in the Auriga simulations to be consistent with the MW GCS, ~~given that we expect dynamical evolution to more strongly affect GC candidates at smaller radii.~~ This trend of over-production of old star particles that are metal-rich and at large radii compared to observed GCs is less clear for the M31 GCS.

- Overall, we conclude that the metallicity and radial distribution of age-selected star particles in the Auriga simulations are not consistent with globular cluster system of the Milky Way or with that of the Andromeda galaxy.

ACKNOWLEDGEMENTS

We thank the anonymous referee for a thoughtful and elaborate report. TLRH acknowledges support from the International Max-Planck Research School (IMPRS) on Astrophysics. TLRH thanks the Max Planck Computing and Data Facility for maintaining the Freya compute cluster, where the simulations were post-processed. FM is supported by the program "Rita Levi Montalcini" of the Italian MIUR. FAG acknowledges financial support from CONICYT through the project FONDECYT Regular Nr. 1181264, and funding from the Max Planck Society through a Partner Group grant. This work has been supported by the European Research Council under ERC-StG grant EXAGAL- 308037. Part of the simulations of this paper used the SuperMUC system at the Leibniz Computing Centre, Garching, under the project PR85JE of the Gauss Centre for Supercomputing. This work used the DiRAC Data Centric system at Durham University, operated by the Institute for Computational Cosmology on behalf of the STFC DiRAC HPC Facility 'www.dirac.ac.uk'. This equipment was funded by BIS National E-infrastructure capital grant ST/K00042X/1, STFC capital grant ST/H008519/1 and STFC DiRAC Operations grant ST/K003267/1 and Durham University. DiRAC is part of the UK National E-Infrastructure.

The analysis in this work was performed using the Python (van Rossum & de Boer 1991) programming language, the IPython (Perez & Granger 2007) environment, the NumPy (van der Walt et al. 2011), SciPy (Jones et al. 2001), and Astropy (Astropy Collaboration et al. 2013) packages. Plots were created with Matplotlib (Hunter 2007), using the perceptually uniform colour maps from Kovesi (2015) for two-dimensional plots. This research has made use of NASA's Astrophysics Data System Bibliographic Services.

REFERENCES

Ashman K. M., Zepf S. E., 1992, *ApJ*, **384**, 50

Ashman K. M., Zepf S. E., 1998, *Globular Cluster Systems*
 Astropy Collaboration et al., 2013, *A&A*, **558**, A33
 Beasley M. A., Baugh C. M., Forbes D. A., Sharples R. M., Frenk C. S., 2002, *MNRAS*, **333**, 383
 Boley A. C., Lake G., Read J., Teyssier R., 2009, *ApJ*, **706**, L192
 Bondi H., 1952, *MNRAS*, **112**, 195
 Bondi H., Hoyle F., 1944, *MNRAS*, **104**, 273
 Boylan-Kolchin M., 2017, *MNRAS*, **472**, 3120
 Brodie J. P., Strader J., 2006, *ARA&A*, **44**, 193
 Burkert A., Forbes D., 2019, arXiv e-prints,
 Caldwell N., Romanowsky A. J., 2016, *ApJ*, **824**, 42
 Caldwell N., Schiavon R., Morrison H., Rose J. A., Harding P., 2011, *AJ*, **141**, 61
 Cen R., 2001, *ApJ*, **560**, 592
 Chiou Y. S., Naoz S., Burkhardt B., Marinacci F., Vogelsberger M., 2019, arXiv e-prints,
 Choksi N., Gnedin O. Y., 2019a, arXiv e-prints,
 Choksi N., Gnedin O. Y., 2019b, *MNRAS*, **486**, 331
 Choksi N., Gnedin O. Y., Li H., 2018, *MNRAS*, **480**, 2343
 Conn A. R., et al., 2012, *ApJ*, **758**, 11
 Conroy C., Naidu R. P., Zaritsky D., Bonaca A., Cargile P., Johnson B. D., Caldwell N., 2019, *ApJ*, **887**, 237
 Côté P., Marzke R. O., West M. J., Minniti D., 2000, *ApJ*, **533**, 869
 Côté P., West M. J., Marzke R. O., 2002, *ApJ*, **567**, 853
 Deason A. J., Belokurov V., Weisz D. R., 2015, *MNRAS*, **448**, L77
 Diemand J., Madau P., Moore B., 2005, *MNRAS*, **364**, 367
 El-Badry K., Quataert E., Weisz D. R., Choksi N., Boylan-Kolchin M., 2019, *MNRAS*, **482**, 4528
 Fall S. M., Rees M. J., 1985, *ApJ*, **298**, 18
 Fattahi A., et al., 2019, *MNRAS*, **484**, 4471
 Faucher-Giguère C.-A., Lidz A., Zaldarriaga M., Hernquist L., 2009, *ApJ*, **703**, 1416
 Forbes D. A., et al., 2018, *Proceedings of the Royal Society of London Series A*, **474**, 20170616
 Galletti S., Federici L., Bellazzini M., Fusi Pecci F., Macrina S., 2004, *A&A*, **416**, 917
 Gebhardt K., Kissler-Patig M., 1999, *AJ*, **118**, 1526
 Gnedin O. Y., 2003, in Kissler-Patig M., ed., *Extragalactic Globular Cluster Systems*. p. 224 (arXiv:astro-ph/0210556), doi:10.1007/10857603_35
 Gnedin O. Y., Lahav O., Rees M. J., 2001, arXiv Astrophysics e-prints,
 Grand R. J. J., et al., 2017, *MNRAS*, **467**, 179
 Grand R. J. J., et al., 2018, *MNRAS*, **474**, 3629
 Gratton R. G., Carretta E., Bragaglia A., 2012, *A&ARv*, **20**, 50
 Gunn J. E., 1980, in Hanes D., Madore B., eds, *Globular Clusters*. p. 301
 Harris W. E., 1991, *ARA&A*, **29**, 543
 Harris W. E., 1996, *AJ*, **112**, 1487
 Harris W. E., 2001, *Globular Cluster Systems*. Springer Berlin Heidelberg, p. 223–408, doi:10.1007/3-540-31634-5_2, https://doi.org/10.1007/3-540-31634-5_2
 Harris W. E., Pudritz R. E., 1994, *ApJ*, **429**, 177
 Holtzman J. A., et al., 1992, *AJ*, **103**, 691
 Holtzman J. A., et al., 1996, *AJ*, **112**, 416
 Hunter J. D., 2007, *Computing in Science and Engineering*, **9**, 90
 Huxor A. P., et al., 2014, *MNRAS*, **442**, 2165
 Johnson T. L., et al., 2017, *ApJ*, **843**, L21
 Jones E., Oliphant T., Peterson P., et al., 2001, *SciPy: Open source scientific tools for Python*, http://www.scipy.org/
 Kang H., Shapiro P. R., Fall S. M., Rees M. J., 1990, *ApJ*, **363**, 488
 Kim J.-h., et al., 2018, *MNRAS*, **474**, 4232
 Kovesi P., 2015, arXiv e-prints, p. arXiv:1509.03700
 Kruijssen J. M. D., 2012, *MNRAS*, **426**, 3008
 Kruijssen J. M. D., 2014, *Classical and Quantum Gravity*, **31**, 244006
 Kruijssen J. M. D., 2015, *MNRAS*, **454**, 1658
 Kruijssen J. M. D., Pelupessy F. I., Lamers H. J. G. L. M., Portegies Zwart S. F., Icke V., 2011, *MNRAS*, **414**, 1339
 Kruijssen J. M. D., Pelupessy F. I., Lamers H. J. G. L. M., Portegies Zwart S. F., Bastian N., Icke V., 2012, *MNRAS*, **421**, 1927

12 *T. L. R. Halbesma et al.*

- Kruijssen J. M. D., Pfeffer J. L., Crain R. A., Bastian N., 2019, *MNRAS*, **486**, 3134
- Lahén N., Naab T., Johansson P. H., Elmegreen B., Hu C.-Y., Walch S., 2019, arXiv e-prints,
- Larsen S. S., Brodie J. P., Huchra J. P., Forbes D. A., Grillmair C. J., 2001, *AJ*, **121**, 2974
- Larson R. B., 1996, in Morrison H. L., Sarajedini A., eds, *Astronomical Society of the Pacific Conference Series Vol. 92, Formation of the Galactic Halo...Inside and Out*. p. 241
- Li H., Gnedin O. Y., 2014, *ApJ*, **796**, 10
- Li H., Gnedin O. Y., Gnedin N. Y., Meng X., Semenov V. A., Kravtsov A. V., 2017, *ApJ*, **834**, 69
- Mackey A. D., et al., 2019, *MNRAS*, **484**, 1756
- Madau P., Lupi A., Diemand J., Burkert A., Lin D. N. C., 2019, arXiv e-prints,
- Marinacci F., Pakmor R., Springel V., 2014, *MNRAS*, **437**, 1750
- Marinacci F., Grand R. J. J., Pakmor R., Springel V., Gómez F. A., Frenk C. S., White S. D. M., 2017, *MNRAS*, **466**, 3859
- McConnachie A. W., Irwin M. J., Ferguson A. M. N., Ibata R. A., Lewis G. F., Tanvir N., 2005, *MNRAS*, **356**, 979
- McLaughlin D. E., van der Marel R. P., 2005, *ApJS*, **161**, 304
- Monachesi A., et al., 2019, *MNRAS*, **485**, 2589
- Muratov A. L., Gnedin O. Y., 2010, *ApJ*, **718**, 1266
- Murray S. D., Lin D. N. C., 1992, *ApJ*, **400**, 265
- Pakmor R., Springel V., 2013, *MNRAS*, **432**, 176
- Pakmor R., Marinacci F., Springel V., 2014, *ApJ*, **783**, L20
- Pakmor R., Springel V., Bauer A., Mocz P., Munoz D. J., Ohlmann S. T., Schaal K., Zhu C., 2016, *MNRAS*, **455**, 1134
- Patel E., Besla G., Sohn S. T., 2017, *MNRAS*, **464**, 3825
- Peebles P. J. E., 1984, *ApJ*, **277**, 470
- Peebles P. J. E., Dicke R. H., 1968, *ApJ*, **154**, 891
- Peng E. W., et al., 2006, *ApJ*, **639**, 95
- Perez F., Granger B. E., 2007, *Computing in Science and Engineering*, **9**, 21
- Pfeffer J., Kruijssen J. M. D., Crain R. A., Bastian N., 2018, *MNRAS*, **475**, 4309
- Portegies Zwart S. F., McMillan S. L. W., Gieles M., 2010, *ARA&A*, **48**, 431
- Recchi S., Wünsch R., Palouš J., Dinnbier F., 2017, *Ap&SS*, **362**, 183
- Renaud F., Agertz O., Gieles M., 2017, *MNRAS*, **465**, 3622
- Ricotti M., Parry O. H., Gnedin N. Y., 2016, *ApJ*, **831**, 204
- Ryu J., Lee M. G., 2018, *ApJ*, **863**, L38
- Schweizer F., 1987, in Faber S. M., ed., *Nearly Normal Galaxies. From the Planck Time to the Present*. pp 18–25
- Spitzer L., 1987, *Dynamical evolution of globular clusters*
- Springel V., 2010, *MNRAS*, **401**, 791
- Springel V., Hernquist L., 2003, *MNRAS*, **339**, 289
- Springel V., Di Matteo T., Hernquist L., 2005, *MNRAS*, **361**, 776
- VandenBerg D. A., Brogaard K., Leaman R., Casagrande L., 2013, *ApJ*, **775**, 134
- Vanzella E., et al., 2017, *MNRAS*, **467**, 4304
- Veljanoski J., et al., 2014, *MNRAS*, **442**, 2929
- Vietri M., Pesce E., 1995, *ApJ*, **442**, 618
- Vogelsberger M., Genel S., Sijacki D., Torrey P., Springel V., Hernquist L., 2013, *MNRAS*, **436**, 3031
- Wang S., Ma J., Liu J., 2019, arXiv e-prints
- West M. J., Côté P., Marzke R. O., Jordán A., 2004, *Nature*, **427**, 31
- Whitmore B. C., Schweizer F., 1995, *AJ*, **109**, 960
- Whitmore B. C., Zhang Q., Leitherer C., Fall S. M., Schweizer F., Miller B. W., 1999, *AJ*, **118**, 1551
- Zepf S. E., Ashman K. M., English J., Freeman K. C., Sharples R. M., 1999, *AJ*, **118**, 752
- Zinn R., 1985, *ApJ*, **293**, 424
- van Rossum G., de Boer J., 1991, *CWI Quarterly*, **4**, 283
- van der Walt S., Colbert S. C., Varoquaux G., 2011, *Computing in Science and Engineering*, **13**, 22

This paper has been typeset from a \LaTeX file prepared by the author.