

Tameez Latib

Problem 13

I, Tameez Latib, declare that this work is my own. I did this work honestly and can fully stand behind everything that I have written.

Suppose that osteoblasts secrete uncalcified bone matrix from $x = 0$ at a constant rate, so that all parts of the matrix move at speed $s > 0$ (along the $+x$ direction) Let's suppose the crystals are uniformly deposited and spaced, have a spherical shape, and grow at a rate proportional to their surface area.

From this, let's model the Volume (of a spherical crystal) as a function of x and t , $V(x, t)$. With V' (the growth) being proportional to surface area

$$(V(x, t))' \propto 4\pi r^2$$

Noting that $V(x, t) = 4/3\pi r^3$, Then $V^{2/3} \propto r^2$
Hence

$$(V(x, t))' \propto V^{2/3}$$

$$(V(x, t))' = c * V^{2/3}$$

Where c is a constant. We can also model the position x as a function of t ; $x(t) = st + x(0)$ That is, the crystals move at a fixed speed s from their initial position $x(0)$.

Now, by the chain rule, the left hand side is:

$$V' = V_t t' + V_x x' = V_t + sV_x$$

So now we have that

$$V_t + sV_x = cV^{2/3}$$

to non-dimensionalise,

$$V = v^* \bar{v} \quad t = t^* \bar{t} \quad x = x^* \bar{x}$$

Then:

$$\frac{\bar{v}}{\bar{t}} v_{t^*}^* + \frac{\bar{v}s}{\bar{x}} v_{x^*}^* = c \bar{v}^{2/3} v^{*2/3}$$

Letting $\bar{x} = s\bar{t}$, $\bar{v} = (c\bar{t})^3$,

$$v_{t^*}^* + \frac{\bar{t}s}{\bar{x}} v_{x^*}^* = c \frac{\bar{t}}{\bar{v}^{1/3}} v^{*2/3}$$

We obtain:

$$v_{t^*}^* + v_{x^*}^* = v^{*2/3}$$

For convenience, I will omit the '*'s, since there will be no further non-dimensionalising or converting back to dimensional form

Now to solve

$$v_t + v_x = v^{2/3}$$

To the constraints: $v(x, 0) = 0, v(0, t) = 1$

First, let $x = x(\tau, \sigma), t = t(\tau, \sigma)$, with σ parametrizing the boundary. Then

$$\frac{dv}{d\tau} = v_x \frac{\partial x}{\partial \tau} + v_t \frac{\partial t}{\partial \tau}$$

If we let

$$\frac{\partial x}{\partial \tau} = \frac{\partial t}{\partial \tau} = 1$$

Then

$$\frac{dv}{d\tau} = v^{2/3}$$

This differential equation can be solved, to get

$$v = \frac{(\tau + c)^3}{27}$$

Now to convert back, σ is the boundary, so

$$x(0, \sigma) = \begin{cases} 0 & \text{if } \sigma < 0 \\ \sigma & \text{if } \sigma > 0 \end{cases}$$

$$t(0, \sigma) = \begin{cases} -\sigma & \text{if } \sigma < 0 \\ 0 & \text{if } \sigma > 0 \end{cases}$$

Then we have that $x - t = \sigma$. Furthermore, since $\frac{\partial x}{\partial \tau} = 1, x = \tau + x(0, \sigma)$

$$\tau = x - x(0, \sigma) = \begin{cases} x - 0 & \text{if } \sigma < 0 \\ x - \sigma & \text{if } \sigma > 0 \end{cases}$$

$$\tau = \begin{cases} x & \text{if } x < t \\ t & \text{if } x > t \end{cases}$$

Then we have

$$v(x, t) = \begin{cases} (x + c)^3/27 & \text{if } x < t \\ (t + c)^3/27 & \text{if } x > t \end{cases}$$

For the initial conditions, $v(0, t) = 1$ and $v(x, 0) = 0$, This means (as $x = 0 < t$ and $t = 0 < x$)

$$v(x, t) = \begin{cases} (x + 3)^3/27 & \text{if } x < t \\ (t)^3/27 & \text{if } x > t \end{cases}$$

However, $x \nsubseteq t$ means that we are trying to find the volume of a crystal where there is no matrix. So the only solution should be

$$v(x, t) = \frac{(x + 3)^3}{27}$$