

MAT 442: Advanced Linear Algebra

Arizona State University

Fall 2020

Contents

1	Vector Spaces	3
1.1	Fields	3
1.2	Vector Spaces	4
1.3	Subspaces	6
1.4	Linear combinations and systems of linear equations	8
1.5	Linear dependence and linear independence	9
1.6	Bases and dimension	10
1.7	Maximal linearly independent subsets	13
2	Linear Transformations and Matrices	14
2.1	Linear transformations, null spaces, and ranges	14
2.2	The matrix representation of a linear transformation	18
2.3	Composition of linear transformations and matrix multiplication	19
2.4	Invertibility and isomorphisms	23
2.5	The change of coordinate matrix	27
2.6	Dual spaces	29
3	Elementary matrix operations and systems of linear equations	32
3.1	Elementary matrix operations and elementary matrices	32
3.2	The rank of a matrix and matrix inverse	33
3.3	Systems of linear equations (theoretical aspect)	39
4	Determinants	41
4.1	Determinants of order 2	41
4.2	Determinants of order n	44
4.3	Properties	49
5	Diagonalization	51
5.1	Eigenvalues and eigenvectors	51
5.2	Diagonalizability	54
5.4	Invariant subspaces and the Cayley-Hamilton theorem	59
6	Inner product spaces	62
6.1	Inner products and norms	62
6.2	The Gram-Schmidt orthogonalization	66
6.3	The adjoint of a linear operator	69
6.4	Normal and self-adjoint operators	71
6.5	Unitary and orthogonal operators and their matrices	75

1 Vector Spaces

1.1 Fields

Definition 1 A *field* is a triple $(\mathbb{F}, +, \cdot)$ with $+: \mathbb{F} \times \mathbb{F} \rightarrow \mathbb{F}$ and $\cdot: \mathbb{F} \times \mathbb{F} \rightarrow \mathbb{F}$ that satisfies the following axioms.

- (1) For all $a, b, c \in \mathbb{F}$, $(a + b) + c = a + (b + c)$ (Associativity of addition)
- (2) For all $a, b, c \in \mathbb{F}$, $(a \cdot b) \cdot c = a \cdot (b \cdot c)$ (Associativity of multiplication)
- (3) For all $a, b \in \mathbb{F}$, $a + b = b + a$ (Commutativity of addition)
- (4) For all $a, b \in \mathbb{F}$, $a \cdot b = b \cdot a$ (Commutativity of multiplication)
- (5) For all $a, b, c \in \mathbb{F}$, $a \cdot (b + c) = a \cdot b + a \cdot c$ (Distributive law)
- (6) There exists $0 \in \mathbb{F}$ such that $a + 0 = a$ for all $a \in \mathbb{F}$ (Identity element of addition)
- (7) There exists $1 \in \mathbb{F}$ such that $a \cdot 1 = a$ for all $a \in \mathbb{F}$ (Identity element of multiplication)
- (8) For all $a \in \mathbb{F}$, there exists $(-a) \in \mathbb{F}$ such that $a + (-a) = 0$ (Additive inverse)
- (9) For all $a \in \mathbb{F}$, there exists $a^{-1} \in \mathbb{F}$ such that $a \cdot a^{-1} = 1$ (Multiplicative inverse)

Examples of Fields

- (1) $(\mathbb{R}, +, \cdot)$
- (2) $(\mathbb{C}, +, \cdot)$
- (3) $(\mathbb{Q}, +, \cdot)$
- (4) $(\{a + b\sqrt{2} : a, b \in \mathbb{Q}\}, +, \cdot)$
- (5) $(\mathbb{Z}_2, +, \cdot)$
- (6) $(\mathbb{Z}_p, +, \cdot)$, where p prime

1.2 Vector Spaces

Definition 1 A *vector space* V over a field F is a triple $(V, +, \cdot)$ where $+: V \times V \rightarrow V$ (addition), $\cdot: F \times V \rightarrow V$ (scalar multiplication) and the following conditions hold.

- (VS 1) For $x, y \in V$, $x + y = y + x$ (Commutativity of addition)
- (VS 2) For $x, y, z \in V$, $(x + y) + z = x + (y + z)$ (Associativity of addition)
- (VS 3) There exists an element $0_V \in V$ such that $0_V + x = x$ for all $x \in V$ (Identity element of addition)
- (VS 4) For $x \in V$, there is $y_x \in V$ such that $x + y_x = 0_V$ (Inverse elements of addition)
- (VS 5) For $x \in V$, $1 \cdot x = x$ (Identity element of scalar multiplication)
- (VS 6) For $a, b \in \mathbb{F}$, $x \in V$, $(ab)x = a(bx)$ (Compatibility of scalar multiplication)
- (VS 7) For $a \in \mathbb{F}$, $x, y \in V$, $a(x + y) = ax + ay$ (Distributivity of scalar multiplication with respect to vector addition)
- (VS 8) For $a, b \in \mathbb{F}$, $x \in V$, $(a + b)x = ax + bx$ (Distributivity of scalar multiplication with respect to field addition)

Conventions

- We will often identify the set V with the vector space V .
- We will write av instead of $a \times v$.
- We will often write 0 for 0_V when V is clear from the context.

An $m \times n$ matrix with entries from a field F is a function $A: \{1, \dots, m\} \times \{1, \dots, n\} \rightarrow F$. We often organize the values in a rectangular array with m rows and n columns and use A_{ij} for $A(i, j)$.

Examples of Vector Fields

- (1) $\mathbb{F}^n = \{(a_1, \dots, a_n) : a_i \in \mathbb{F}\}$
- (2) $\{f: S \rightarrow \mathbb{F} : S \neq \emptyset\}$. Also notated $\mathcal{F}(S, F)$.
- (3) Set of all sequences over \mathbb{F}
- (4) $P(\mathbb{F})$. Set of all polynomials with coefficients in \mathbb{F} .
- (5) $M_{m \times n}(\mathbb{F})$

Example: Is the empty set a vector space?

Answer. No. By rule (VS 3), there must exist $0_V \in V$, such that $0_V + x = x$ for all $x \in V$, but since $V = \emptyset$, $\nexists 0_V \in V$.

Example: Decide if V is a vector space.

$$(1) \quad V = \{(a_1, a_2) : a_1, a_2 \in \mathbb{R}\}, (a_1, a_2) + (b_1, b_2) = (a_1 + b_1, a_2 - b_2), c(a_1, a_2) = (ca_1, ca_2)$$

Answer. No. V violates axiom (VS 1).

$$(2) \quad V = \{(a_1, a_2) : a_1, a_2 \in \mathbb{R}\}, (a_1, a_2) + (b_1, b_2) = (a_1 + b_1, 0), c(a_1, a_2) = (ca_1, 0)$$

Answer. No. V violates axiom (VS 5).

$$(3) \quad V = \{(a_1, a_2) : a_1, a_2 \in \mathbb{R}\}, (a_1, a_2) + (b_1, b_2) = (a_1 + b_1, a_2 b_2), c(a_1, a_2) = (ca_1, a_2)$$

Answer. No. V violates axiom (VS 4).

Theorem 1.1. (Cancellation Law for Vector Addition) Let $x, y, z \in V$. If $x + z = y + z$, then $x = y$.

Proof. Suppose $x + z = y + z$. Let w be such that $z + w = 0_V$. Then

$$\begin{aligned} x &= x + 0_V \\ &= x + (z + w) \\ &= (x + z) + w \\ &= (y + z) + w \\ &= y + (z + w) \\ &= y + 0_V \\ &= y. \end{aligned}$$

□

Corollary 2

- There is unique $0_V \in V$ such that $0_V + x = x$ for all $x \in V$.

Proof. Let $v \in V$. Suppose that 0_V and $0'_V$ are zero elements.

$$v + 0_V = v = v + 0'_V.$$

Thus, by cancellation law, $0_V = 0'_V$.

□

- For every $x \in V$, there is unique $y_x \in V$ such that $x + y_x = 0_V$. *Proof.* Let $x \in V$. Suppose that y_x and y'_x are inverses of x . Then

$$x + y_x = 0 = x + y'_x.$$

Thus, by cancellation law, $y_x = y'_x$. □

Theorem 1.2. Let $x \in V$, $a \in \mathbb{F}$. Then

- $0x = 0_V$;

Proof.

$$\begin{aligned} 0x + 0x &= (0 + 0)x \\ &= 0x \\ &= 0x + 0_V. \end{aligned}$$

Thus, by cancellation law, $0x = 0_V$. □

- $(-a)x = -(ax) = a(-x)$;

Proof. We can show

$$\begin{aligned} (-a)x + (ax) &= (-a + a)x \\ &= 0x \\ &= 0_V. \end{aligned}$$

Thus, the inverse of ax is unique, which implies $-ax = (-a)x$. We have $(-x) = (-1)x$, which implies $a(-x) = (a(-1))x = (-a)x$. □

- $a0_V = 0_V$.

Proof.

$$a0_V = a(0_V - 0_V) = a0_V - a0_V = 0_V.$$

□

1.3 Subspaces

Definition 2 Let $(V, +, \cdot)$ be a vector space and let $W \subseteq V$. Then W is called a *subspace* if $(W, +, \cdot)$ is a vector space.

Note: W is a subspace if

- $x + y \in W$ for $x, y \in W$

- $cx \in W$ for $x \in W$, $c \in \mathbb{F}$
- W has a zero vector 0_W
- For $x \in W$ there is $y_x \in W$ such that $x + y_x = 0_W$

Theorem 1.3. Let W be a subset of V a vector space $(V, +, \cdot)$. Then W is a subspace of V if and only if the following hold.

- (1) $0_V \in W$
- (2) if $x, y \in W$, then $x + y \in W$
- (3) if $c \in \mathbb{F}$ and $x \in W$, then $cx \in W$

Proof. (\implies): Suppose W is a subspace. Then (2) and (3) are satisfied. Then, $0_W + 0_W = 0_W = 0_V$. Thus, by cancellation law, $0_W = 0_V$.

(\impliedby): Suppose $x \in W$. We have $-x = (-1)x$ and $-1 \in \mathbb{F}$, $x \in W$. Thus, $-x \in W$. \square

The *transpose* of an $m \times n$ matrix A is the $n \times m$ matrix A^t such that $(A^t)_{ij} = A_{ji}$. Note, $(A + B)^t = A^t + B^t$. Also, $(cA^t) = cA^t$. This is proven by,

$$\begin{aligned} ((A + B)^t)_{ij} &= (A + B)_{ji} \\ &= A_{ji} + B_{ji} \\ &= (A^t)_{ij} + (B^t)_{ij}. \end{aligned}$$

A is called *symmetric* if $A^t = A$.

An $n \times n$ matrix A is called *diagonal* if $A_{ji} = 0$ whenever $j \neq i$.

Theorem 1.4. Any intersection of subspaces of a vector space V is a subspace of V .

Definition 3

- Let $S_1, S_2 \subseteq V$. Then $S_1 + S_2$ is the set $\{x + y : x \in S_1, y \in S_2\}$
- V is called a direct sum of W_1 and W_2 if W_1, W_2 are subspaces of V such that $W_1 \cap W_2 = \{0\}$ and $W_1 + W_2 = V$. We then write $V = W_1 \oplus W_2$.

Example: Let $V = \mathbb{R}^2$, $S_1 = \{(a, 0) : a \in \mathbb{R}\}$, $S_2 = \{(0, b) : b \in \mathbb{R}\}$. Then $S_1 + S_2$ generates V .

Examples of subspaces

- (1) $\mathcal{C} = \{f \in \mathcal{F}(\mathbb{R}, \mathbb{R}) : f \text{ is continuous}\}$ is a subspace of $\mathcal{F}(\mathbb{R}, \mathbb{R})$.
- (2) $P_n(\mathbb{R}) = \{f : f \text{ is a polynomial and } \deg(f) \leq n\}$ is a subspace of $P(\mathbb{R})$.
- (3) $D = \{A \in M_{n \times n}(\mathbb{R}) : A \text{ is diagonal}\}$ is a subspace of $M_{n \times n}(\mathbb{R})$.
- (4) $W = \{A \in M_{n \times n}(\mathbb{R}) : \text{tr}(A) = 0\}$ is a subspace of $M_{n \times n}(\mathbb{R})$.

1.4 Linear combinations and systems of linear equations

Definition 4 Let S be a subset of a vector space V and let $v \in V$. Then v is called a linear combination of vectors of S if there exists $u_1, \dots, u_n \in S$ and $a_1, \dots, a_n \in \mathbb{F}$ such that

$$v = \sum_{i=1}^n a_i v_i.$$

When solving a system of linear equations we reduced to what is called the reduction echelon form by performing the following three operations.

- Interchange two equations
- Multiply an equation by a non-zero element from F
- Add a scalar multiple of one equation to another

Example: $v = (2, 6, 8) \in \mathbb{R}^3$ is a linear combination of $v_1 = (1, 2, 1)$, $v_2 = (-2, -4, -2)$, $v_3 = (0, 2, 3)$, $v_4 = (2, 0, -3)$, $v_5 = (-3, 8, 16)$. Specifically, $v = a_1 v_1 + \dots + a_5 v_5$ where $(a_1, a_2, a_3, a_4, a_5) = (-4, 0, 7, 3, 0)$ is one solution.

Definition 5 Let S be a subset of a vector space V . If S is non-empty, we let $\text{span}(S)$ be the set of all linear combinations of vectors from S , and we set $\text{span}(\emptyset) = \{0_V\}$.

Theorem 1.5. Let S be a subset of a vector space V . Then $\text{span}(S)$ is a subspace of V . Moreover, if W is a subspace of V such that $S \subseteq W$, then $\text{span}(S) \subseteq W$.

Proof.

- (1) (Show $\text{span}(S)$ is a subspace): If $S = \emptyset$, then $\text{span}(S) = \{0_V\}$, which is a subspace. Assume $S \neq \emptyset$. Let $v \in S$. Then,

- (a) $0v = 0_V \in \text{span}(S)$
- (b) Suppose $x, y \in \text{span}(S)$. Then $x = \sum_{i=1}^n a_i v_i$, $y = \sum_{i=1}^m b_i w_i$ where $v_1, \dots, v_n, w_1, \dots, w_m \in S$ and $a_1, \dots, b_1, \dots, b_m \in \mathbb{F}$. Let

$$u_i = \begin{cases} v_i & i \leq n \\ w_{i-n} & i > n \end{cases}, \quad c_i = \begin{cases} a_i & i \leq n \\ b_{i-n} & i > n. \end{cases}$$

Then $x + y = \sum_{i=1}^{n+m} c_i u_i$ and $u_i \in S$, $c_i \in \mathbb{F}$ for $i = 1, \dots, n+m$. Thus, $x + y \in \text{span}(S)$.

- (c) Let $x \in \text{span}(S)$ and let $c \in \mathbb{F}$. Then $x = \sum_{i=1}^n a_i v_i$ where $v_1, \dots, v_n \in S$ and $a_1, \dots, a_n \in \mathbb{F}$. Then $cx = \sum_{i=1}^n (ca_i) v_i$ and $ca_i \in \mathbb{F}$.

Therefore $\text{span}(S)$ is a subspace.

(2) Let $x \in \text{span}(S)$. Then, $x = \sum_{i=1}^n a_i v_i$, where $a_i \in \mathbb{F}$ and $v_i \in S$. We will use induction on n .

- If $n = 1$, $x = a_1 v_1$, and $v_1 \in W$. Thus, $x \in W$.
- Suppose $x = \sum_{i=1}^n a_i v_i$. Then $x = \sum_{i=1}^{n-1} (a_i v_i) + a_n v_n$. Then $\sum_{i=1}^{n-1} (a_i v_i) \in W$ by inductive hypothesis and $a_n v_n \in W$. Thus $x \in W$.

Therefore, $\text{span}(S) \subseteq W$.

□

Note: In particular, $\text{span}(\text{span}(S)) = \text{span}(S)$.

Proof.

(a) $\text{span}(S) \subseteq \text{span}(\text{span}(S))$

(b) Let $x \in \text{span}(\text{span}(S))$. Then $\text{span}(S) \subseteq \text{span}(S)$ implies $\text{span}(S) \subseteq x$, which gives us $\text{span}(\text{span}(S)) \subseteq \text{span}(S)$. Therefore, $(\text{span}(\text{span}(S))) = \text{span}(S)$.

□

Definition 6 A subset S of V *generates* (spans) V if $\text{span}(S) = V$.

Example: Let $V = M_{2 \times 2}(\mathbb{R})$ and $S = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \right\}$. Then $\text{span}(S) = V$.

1.5 Linear dependence and linear independence

Definition 7 A subset S of a vector space V is called *linearly dependent* if there exists distinct vectors $u_1, \dots, u_n \in S$ and $a_1, \dots, a_n \in \mathbb{F}$ not all zero such that $a_1 u_1 + \dots + a_n u_n = 0$.

Notes:

- The empty set is linearly dependent.
- $S = \{u\}$ is linearly dependent if and only if $u \neq 0$.
- A set is called linearly independent if and only if the only representation of 0 as linear combinations of its vectors are trivial.

Theorem 1.6. Let $S_1 \subseteq S_2 \subseteq V$. If S_1 is linearly dependent, then so is S_2 .

Example: Let $S = \{(1, 0, 0, -1), (0, 1, 0, -1), (0, 0, 1, -1), (0, 0, 0, 1)\} \subseteq \mathbb{R}^4$. We'll show S is linearly independent. Suppose

$$a_1(1, 0, 0, -1) + a_2(0, 1, 0, -1) + a_3(0, 0, 1, -1) + a_4(0, 0, 0, 1) = (0, 0, 0, 0).$$

We get $a_1 = 0, a_2 = 0, a_3 = 0, -a_1 - a_2 - a_3 + a_4 = 0$, which implies $a_1 = a_2 = a_3 = a_4 = 0$.

Example: $P_k(x) = x^k + \cdots + x^n$ where $1 \leq k \leq n$. We'll show that $\{P_0(x), \dots, P_n(x)\}$ is linearly independent. Suppose $a_0P_0(x) + \cdots + a_nP_n(x) = 0$. The coefficient of x^i on the left hand side is $a_0 + \cdots + a_i$. Then we have

$$\begin{aligned} a_0 + \cdots + a_n &= 0 \\ a_0 + \cdots + a_{n-1} &= 0 \\ &\vdots \\ a_0 &= 0. \end{aligned}$$

Thus, $a_0 = \cdots = a_n = 0$.

Corollary 8 If $S_1 \subseteq S_2 \subseteq V$ and S_2 is linearly independent, then so is S_1 .

Theorem 1.7. Let S be a linearly independent subset of V and let $v \in V \setminus S$. Then $S \cup \{v\}$ is linearly dependent if and only if $v \in \text{span}(S)$.

Proof. Let S be linearly independent, $v \notin S$. We'll show $S \cup \{v\}$ is linearly dependent if and only if $v \in \text{span}(S)$.

(\implies): Suppose $T = S \cup \{v\}$ is linearly dependent. Then, there exists a_1, \dots, a_n not all zero and there exists $u_1, \dots, u_n \in T$ such that $a_1u_1 + \cdots + a_nu_n = 0_V$. Since S is linearly independent, $u_i = v$ for some $1 \leq i \leq n$. Say $u_1 = v$ and $a_i \neq 0$. Then $a_1u_1 = -a_2u_2 - \cdots - a_nu_n$. Thus $u_1 = -\frac{a_2}{a_1}u_2 - \cdots - \frac{a_n}{a_1}u_n$. Also $u_2, \dots, u_n \in S$. Therefore $u_1 \in \text{span}(S)$.

(\impliedby): Let $v \in \text{span}(S)$. Then there exists a_1, \dots, a_n not all zero and there exists $v_1, \dots, v_n \in S$ such that $v = a_1v_1 + \cdots + a_nv_n$. Then $v - a_1v_1 - \cdots - a_nv_n = 0_V$. The coefficient on v is $1 \neq 0$. Thus, $\{v\} \cup S$ is linearly dependent. \square

1.6 Bases and dimension

Definition 8 A *basis* β for a vector space V is a linearly independent subset of V that generates V .

Theorem 1.8. Let V be a vector space and let $\beta = \{u_1, \dots, u_n\}$. Then β is a basis of V if and only if each v can be uniquely written as

$$v = a_1u_1 + \cdots + a_nu_n$$

where $a_1, \dots, a_n \in \mathbb{F}$.

Proof. (\Rightarrow): Suppose $\beta = \{u_1, \dots, u_n\}$ is a basis for a vector space V . Then $\text{span}(\beta) = V$ and every $v \in V$ can be written as a linear combination of u_1, \dots, u_n . Suppose $v = \sum_{i=1}^n a_i u_i$ and $v = \sum_{i=1}^n b_i u_i$. Then $0_V = \sum_{i=1}^n (a_i - b_i) u_i$. Thus $a_i = b_i = 0$ for every $i = 1, \dots, n$.

(\Leftarrow): Suppose every $v \in V$ can be written uniquely as $v = \sum_{i=1}^n a_i u_i$. Then $\text{span}(\{u_1, \dots, u_n\}) = V$. Suppose $a_1 u_1 + \dots + a_n u_n = 0_V$. Note, $0_V = u_1 \cdot 0 + \dots + u_n \cdot 0$. Since every vector has a unique representation as a linear vector, then $a_1 = \dots = a_n = 0$. Thus, $\{u_1, \dots, u_n\}$ is linearly independent and a basis for V . \square

Theorem 1.9. Let S be a finite set that generates V . Then there is a subset of S which is a basis for V .

Proof. If $S = \emptyset$ or $S = \{0_V\}$, then $V = \{0_V\}$ and \emptyset is a basis. Assume S contains a finite set of at least one nonzero vector v , which generates V . Then $\{v\}$ is linearly independent. Let $\{u_1, \dots, u_k\}$ is a maximal linearly independent subset of S . Then $S \subseteq \text{span}(\{u_1, \dots, u_k\})$. Then $\text{span}(S) \subseteq \text{span}(\text{span}(\{u_1, \dots, u_k\}))$. So, $\text{span}(S) = V \subseteq \text{span}(\{u_1, \dots, u_k\})$. Therefore $\{u_1, \dots, u_k\}$ is a basis for V . \square

Theorem 1.10. Let $G \subset V$, $|G| = n$ and $V = \text{span}(G)$. Suppose further that $L \subseteq V$, $|L| = m$ and L is linearly independent. Then $m \leq n$ and there exists a subset H of G such that $|H| = n - m$ and $L \cup H$ generates V .

Proof. (Outline)

- Induction on m . For the inductive step consider $L = \{v_1, \dots, v_{m+1}\}$
- Apply induction to $\{v_1, \dots, v_m\}$. Thus $m \leq n$ and there is a subset $H' \dots$
- H' can't be empty, say $H' = \{u_1, \dots, u_{n-m}\}$. Thus $m \leq n$.
- To find H for L , use the fact $H' \cup \{v_1, \dots, v_m\}$ generates V and substitute one of vectors from H' with v_{m+1} .

Proof. Let $|G| = n$, $\text{span}(G) = V$, $|L| = m$, L is linearly independent. Then, using induction on m ,

(Base Case): If $m = 0$, then $L = \emptyset$, $|L| = 0$ and $H = G$ and $|H| = n = n - 0$.

(Inductive Step): Suppose $L = \{v_1, \dots, v_{m+1}\}$ is linearly independent. Let $L' = \{v_1, \dots, v_m\}$. By inductive hypothesis, there exists $m \leq n$ and there exists $H' \subseteq G$, such that $|H'| = n - m$ and $\text{span}(L' \cup H') = V$. Note, $H' \neq \emptyset$. Otherwise, $\text{span}(L') = V$. In particular, $v_{m+1} \in \text{span}(\{v_1, \dots, v_m\})$, but L is linearly independent, contradiction. Therefore, $H' \neq \emptyset$ and so $n - m > 0$. Thus, $m + 1 \leq n$. Say $H' = \{u_1, \dots, u_{n-m}\}$. Since $\text{span}(L' \cup H') = V$,

$$v_{m+1} = \sum_{i=1}^m a_i v_i + \sum_{i=1}^{n-m} b_i u_i.$$

Now, at least one of scalars b_i is nonzero, say $b_1 \neq 0$. Then, $b_1 u_1 = v_{m+1} - \sum_{i=2}^m a_i v_i - \sum_{i=2}^{n-m} b_i u_i$. So,

$$u_1 = \frac{1}{b_1} v_{m+1} - \sum_{i=2}^m \frac{a_i}{b_1} v_i - \sum_{i=2}^{n-m} \frac{b_i}{b_1} u_i,$$

which implies $u_1 \in \text{span}(\{v_1, \dots, v_{m+1}\} \cup \{u_2, \dots, u_{n-m}\})$. Thus, $u_1 \in \text{span}(L \cup \{u_2, \dots, u_{n-m}\})$. Let $H = \{u_2, \dots, u_{n-m}\}$. Then $|H| = n - m - 1 = n - (m + 1)$ and $H \subseteq G$. We have $u_i \in \text{span}(L \cup H)$ for $i = 1, \dots, n - m$ and $L \subseteq \text{span}(L \cup H)$. Thus, $V \subseteq \text{span}(L \cup \{u_1, \dots, u_{n-m}\}) \subseteq \text{span}(L \cup H)$. \square

Corollary 13 Suppose V has finite basis. Then every basis for V has the same cardinality.

Proof. Let β and γ be bases for V .

- β is linearly independent, $\text{span}(\gamma) = V$, $|\beta| \leq |\gamma|$ by Theorem 1.10.
- So, γ is linearly independent and $\text{span}(\beta) = V$. Again, by Theorem 1.10, $|\gamma| \leq |\beta|$.

\square

Definition 9 A vector space is called *finite-dimensional* if it has a finite basis. The number of vectors in a basis, is called the *dimension* of V , notated $\dim(V)$. A vector space which is not finite-dimensional is called *infinite-dimensional*.

Corollary 14 Suppose $\dim(V) = n$.

- (a) If S is finite and $\text{span}(S) = V$, then $n \leq |S|$. If $|S| = n$, then S is a basis.

Proof. Let S be finite and $\text{span}(S) = V$. Let β be a basis for V . Then, β is linearly independent and $|\beta| = n$. By Theorem 1.10, $|S| \geq |\beta| = n$. Suppose $|S| = n$. Then S contains a subset T such that T is linearly independent and $\text{span}(T) = V$. Consequently, T is a basis for V . Thus, $|T| = \dim(V) = n$, $T \subseteq S$, $|T| = n$. Therefore, $S = T$, which implies S is a basis. \square

- (b) If $|S| = n$ and S is linearly independent, then S is a basis.

Proof. Suppose L is linearly independent and $|L| = \dim(V)$. Let β be a basis for V . Then $|L| \leq |\beta|$, in addition, there exists $H \subseteq \beta$ such that $\text{span}(L \cup H) = V$ and $|H| = |\beta| - |L| = \dim(V) - \dim(V) = 0$. Thus, $H = \emptyset$. Thus $\text{span}(L) = V$. \square

- (c) Every linearly independent set can be extended to a basis.

Proof. Let L be linearly independent. Suppose $|L| < n$. Let β be a basis. Then there exists $H \subseteq \beta$ such that $\text{span}(L \cup H) = V$ and $|H| = n - |L|$. So, $|L \cup H| \leq n$. As before, $L \cup H$ contains an independent subset T such that $\text{span}(T) = V$. Then T is a basis and so $|T| = n$. Therefore, $T = L \cup H$. Thus $L \cup H$ is a basis. \square

Example:

- (1) \mathbb{F}^n , $\dim(\mathbb{F}^n) = n$.
- (2) $V = M_{m \times n}(\mathbb{F})$, $\dim(V) = mn$.
- (3) $V = \{A \in M_{n \times n} : A \text{ is symmetric}\}$, $\dim(V) = \frac{n(n+1)}{2}$.

Theorem 1.11. Let W be a subspace of a finite-dimensional space V . Then W is finite-dimensional and $\dim(W) \leq \dim(V)$. Moreover, if $\dim(W) = \dim(V)$, then $W = V$.

1.7 Maximal linearly independent subsets

Definition 10 A collection \mathcal{C} of sets is called a *chain* if for every $A, B \in \mathcal{C}$, $A \subseteq B$ or $B \subseteq A$.

Maximal Principle: Let \mathcal{F} be a family of sets. If for every chain \mathcal{C} in \mathcal{F} there is a set in \mathcal{F} which contains all members of \mathcal{C} , then \mathcal{F} contains a maximal element.

Theorem 1.12. Every vector space has a basis.

Proof. (Outline)

- Start with an arbitrary (finite) linearly independent set S in V and consider the family \mathcal{F} of all independent sets contains S . Argue that Maximal Principle applies and take the element β in \mathcal{F} .
- β generates V .

Cauchy's functional equation:

$$f(x + y) = f(x) + f(y)$$

What type of functions can f be?

- (1) $f : \mathbb{Q} \rightarrow \mathbb{Q}$ where $f(x) = \alpha \cdot x$ and $\alpha = f(1)$.
- (2) $f : \mathbb{R} \rightarrow \mathbb{R}$ allows for other fairly exotic functions.

For example, \mathbb{R} is a vector space over \mathbb{Q} . Let H be a basis of this vector space, commonly called Hamel basis. Then for every element $x \in \mathbb{R}$ there exists unique $h_1, \dots, h_n \in H$ and unique scalars $c_1, \dots, c_n \in \mathbb{Q}$ with $c_1, \dots, c_n \neq 0$ such that $x = \sum_{i=1}^n c_i h_i$. Then, for any $g : H \rightarrow \mathbb{R}$, we can extend g to $\bar{g} : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$\bar{g}(x) = \sum_{i=1}^n c_i g(h_i).$$

So,

$$x + y = \sum_{i=1}^n d_i h_i + \sum_{i=1}^n a_i h_i = \sum_{i=1}^n c_i h_i.$$

2 Linear Transformations and Matrices

2.1 Linear transformations, null spaces, and ranges

Definition 1 A function $T : V \rightarrow W$ is called a *linear transformation* from V to W if for all $x, y \in V$ and $c \in \mathbb{F}$ the following hold.

- $T(x + y) = T(x) + T(y)$
- $T(cx) = cT(x)$

Observations:

- T is linear if and only if $x_1, \dots, x_n \in V, a_1, \dots, a_n \in \mathbb{F}$

$$T\left(\sum_{i=1}^n a_i x_i\right) = \sum_{i=1}^n a_i T(x_i).$$

- T is linear if and only if $T(cx + y) = cT(x) + T(y)$ for $x, y \in V, c \in \mathbb{F}$.

Example:

- (1) Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ to be $T(a_1, a_2) = (2a_1 + a_2, a_1)$. Then

$$\begin{aligned} T((a_1, a_2) + (b_1, b_2)) &= T(a_1 + b_1, a_2 + b_2) \\ &= (2(a_1 + b_1) + a_2 + b_2, a_1 + b_1) \\ &= (2a_1 + a_2, a_1) + (2b_1 + b_2, b_1) \\ &= T(a_1, a_2) + T(b_1, b_2). \end{aligned}$$

It's also easy to show,

$$T(c(a_1, a_2)) = cT(a_1, a_2).$$

- (2) Let $T_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ where $T_\theta(a_1, a_2)$ is the vector obtained by (a_1, a_2) by the angle θ .

- (3) Let $V = \mathcal{C}(\mathbb{R})$, and $a, b \in \mathbb{R}$, with $a < b$. Let $T : V \rightarrow \mathbb{R}$, where

$$T(f) = \int_a^b f(t) dt.$$

Then, $T(f + g) = T(f) + T(g)$. Also, $T(cf) = cT(f)$.

Definition 2 Let $T : V \rightarrow W$ be linear.

- The *null space* (kernel) $N(T) = \{x \in V : T(x) = 0\}$

- The *range* (image) $R(T) = \{T(x) : x \in V\}$

Theorem 2.1. Let $T : V \rightarrow W$ be linear. Then $N(T)$ and $R(T)$ are subspaces of V and W .

Proof.

- (1) We have $0_V \in N(T)$ because $T(0_V) = 0_W$.
- (2) Suppose $x, y \in N(T)$. Then $T(x) = 0_W$, $T(y) = 0_W$. Thus, $T(x + y) = T(x) + T(y) = 0_W + 0_W = 0_W$. Thus $x + y \in N(T)$.
- (3) Suppose $x \in N(T)$ and $c \in \mathbb{F}$. then $T(x) = 0_W$ and so $T(cx) = cT(x) = 0_W$. Thus, $cx \in N(T)$.

Therefore $N(T)$ is a subspace of V . $R(T)$ can be shown to be a subspace of W in a similar manner. \square

Theorem 2.2. Let $T : V \rightarrow W$ be linear and let $\beta = \{v_1, \dots, v_n\}$ be a basis for V . Then $R(T) = \text{span}(T(\beta)) = \text{span}(\{T(v_1), \dots, T(v_n)\})$.

Proof.

- (1) Let $w \in R(T)$. Then $w = T(v)$ for some $v \in V$. Then $v = \sum_{i=1}^n c_i v_i$ for some $c_1, \dots, c_n \in \mathbb{F}$. Thus, $T(v) = \sum_{i=1}^n c_i T(v_i)$. Therefore, $w = T(v) \in \text{span}(T(\beta))$.
- (2) Let $w \in \text{span}(T(\beta))$. Then $w = \sum_{i=1}^n c_i T(v_i)$ for some $c_1, \dots, c_n \in \mathbb{F}$. Thus $w = T(\sum_{i=1}^n c_i v_i)$ and $\sum_{i=1}^n c_i v_i \in V$. Thus $w \in R(T)$.

\square

- $\text{nullity}(T) = \dim(N(T))$
- $\text{rank}(T) = \dim(R(T))$

Theorem 2.3. (Dimension Theorem) Let $T : V \rightarrow W$ be linear. If V is finite-dimensional, then

$$\text{nullity}(T) + \text{rank}(T) = \dim(V).$$

Proof. (Outline)

- Start with a basis for $N(T)$, $\{v_1, \dots, v_k\}$ and extend it to a basis of V , $\{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$.
- Prove that $\{T(v_{k+1}), \dots, T(v_n)\}$ is a basis for $R(T)$.

Proof. Let $\{v_1, \dots, v_k\}$ be a basis for $N(T)$. Extend this basis to a basis β for V , say $\beta = \{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$. Note, $k = \dim(N(T))$ and $n = \dim(V)$. We claim that $\{T(v_{k+1}), \dots, T(v_n)\}$ is a basis for $R(T)$.

- (1) Clearly, $\text{span}(\{T(v_{k+1}), \dots, T(v_n)\}) \subseteq R(T)$. Let $w \in R(T)$. Then $w = T(v)$ for some $v \in V$ and $v = \sum_{i=1}^n c_i v_i$ for some $c_1, \dots, c_n \in \mathbb{F}$. Then $w = T(v) = \sum_{i=1}^n c_i T(v_i) = \sum_{i=k+1}^n c_i T(v_i)$ since $T(v_i) = 0$ for $i \leq k$. Therefore, $w \in \text{span}(\{T(v_{k+1}), \dots, T(v_n)\})$. So, $\text{span}(\{T(v_{k+1}), \dots, T(v_n)\}) = R(T)$.
- (2) We'll show $\{T(v_{k+1}), \dots, T(v_n)\}$ is linearly independent. Suppose $\sum_{i=k+1}^n c_i T(v_i) = 0$. Then, $T(\sum_{i=k+1}^n c_i v_i) = 0$. Thus, $\sum_{i=k+1}^n c_i v_i \in N(T)$. So, $\sum_{i=k+1}^n c_i v_i = \sum_{i=1}^k d_i v_i$ for some $d_1, \dots, d_k \in \mathbb{F}$. Therefore, $\sum_{i=k+1}^n c_i v_i - \sum_{i=1}^k d_i v_i = 0$. Since β is linearly independent, $c_{k+1} = \dots = c_n = 0$. In addition, $d_1 = \dots = d_k = 0$. Thus, $\{T(v_{k+1}), \dots, T(v_n)\}$ is linearly independent. □

Theorem 2.4. Let $T : V \rightarrow W$ be linear. Then T is injective if and only if $N(T) = \{0\}$.

Proof.

(\implies): Suppose $T : V \rightarrow W$ is injective. If $T(x) = 0$ then since $T(0) = 0$, we have $T(x) = T(0)$ and so $x = 0$.

(\impliedby): Suppose $N(T) = \{0\}$. If $T(x) = T(y)$, then $T(x) - T(y) = 0$. Thus, $T(x - y) = 0$. Thus, $x - y \in N(T)$, which implies $x - y = 0$ and thus $x = y$. □

Theorem 2.5. Let V, W be finite-dimensional vector spaces such that $\dim(V) = \dim(W)$ and let $T : V \rightarrow W$ be linear. Then the following are equivalent:

- (a) T is injective;
- (b) T is surjective;
- (c) $\text{rank}(T) = \dim(V) = \dim(W)$.

Proof. Note, T being injective is equivalent to $N(T) = \{0\}$, which is equivalent to $\text{nullity}(T) = 0$. By the dimension theorem, $\text{rank}(T) = \dim(V)$. So, $\text{rank}(T) = \dim(W)$, which is equivalent to $\dim(R(T)) = \dim(W)$. This is equivalent to $R(T) = W$ since $R(T) \subseteq W$. Thus, T is surjective. □

Theorem 2.6. Suppose $\{v_1, \dots, v_n\}$ is a basis for V . For $w_1, \dots, w_n \in W$ there exists exactly one linear transformation $T : V \rightarrow W$ such that $T(v_i) = w_i$ for every i .

Proof. (Outline)

- For $x \in V$ we can write $x = \sum_{i=1}^n a_i v_i$ uniquely.
- Let $T : V \rightarrow W$ be $T(x) = \sum_{i=1}^n a_i w_i$.

Proof. Let $\{v_1, \dots, v_n\}$ be a basis for V and $w_1, \dots, w_n \in W$. Then there exists unique $T : V \rightarrow W$ defined by $T(v_i) = w_i$.

- (1) Let $x \in V$. Then x can be uniquely written as $x = \sum_{i=1}^n a_i v_i$ where $a_i \in \mathbb{F}$. Let $T(x) := \sum_{i=1}^n a_i w_i$. $T : V \rightarrow W$ is well-defined, and thus a function.
- (2) Let T be linear and $u, v \in V$. Let $c \in \mathbb{F}$. We will show $T(cu + v) = cT(u) + T(v)$. Then, $u = \sum_{i=1}^n a_i v_i$ and $v = \sum_{i=1}^n b_i v_i$ for unique $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{F}$. Then $cu + v = \sum_{i=1}^n (ca_i + b_i) v_i$. So,

$$\begin{aligned} T(cu + v) &= \sum_{i=1}^n (ca_i + b_i) w_i \\ &= c \sum_{i=1}^n a_i w_i + \sum_{i=1}^n b_i w_i \\ &= cT(u) + T(v). \end{aligned}$$

- (3) Suppose $U : V \rightarrow W$ is linear and $U(v_i) = w_i$. Let $x \in V$. Then $x = \sum_{i=1}^n a_i v_i$. Then

$$\begin{aligned} U(x) &= U\left(\sum_{i=1}^n a_i v_i\right) \\ &= \sum_{i=1}^n a_i U(v_i) \\ &= \sum_{i=1}^n a_i w_i \\ &= T(x). \end{aligned}$$

- (4) Note, $T(v_i) = w_i$ because $v_i = 1 \cdot v_i$. Thus, $T(v_i) = 1 \cdot w_i = w_i$.

□

Corollary 7 Let $\{v_1, \dots, v_n\}$ be a basis of V . If $U, T : V \rightarrow W$ are linear and $T(v_i) = U(v_i)$, then $T = U$.

2.2 The matrix representation of a linear transformation

Definition 3 Let V be a finite-dimensional vector space. An *ordered basis* for V is a basis equipped with an ordering.

Example: Let $\beta = \left\{ e_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, e_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$. Let $x = \begin{pmatrix} 5 \\ 6 \\ 7 \end{pmatrix} = 5e_1 + 6e_2 + 7e_3$.

Then $[x]_\beta = \begin{pmatrix} 5 \\ 6 \\ 7 \end{pmatrix}$.

Definition 4 Let $\beta = \{u_1, \dots, u_n\}$ be an ordered basis for V . Then for every $x \in V$, $x = \sum a_i u_i$ and a_1, \dots, a_n are unique. The coordinate vector of x relative to β , $[x]_\beta = (a_1, \dots, a_n)^T$.

Let $T : V \rightarrow W$, $\beta = \{v_1, \dots, v_n\}$, $\gamma = \{w_1, \dots, w_m\}$ be ordered bases for V and W . We have

$$T(v_j) = \sum_{i=1}^m a_{ij} w_i.$$

Then $A = [a_{ij}]$ is called the matrix representation of T in the ordered bases β and γ and we write

$$A = [T]_\beta^\gamma.$$

If $V = W$ and $\beta = \gamma$, we say $A = [T]_\beta$.

Observations:

- The j th column of A is $[T(v_j)]_\gamma$.
- If $T, U : V \rightarrow W$ and $[U]_\beta^\gamma = [T]_\beta^\gamma$, then $T = U$.

Example: Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $\beta = \{(1, 0), (0, 1)\}$ and $\gamma = \{(1, 1), (1, -1)\}$. Let $T((1, 0)) = (1, 0)$ and $T((0, 1)) = (1, 1)$. Find $[T]_\beta^\gamma$.

We can see $T((1, 0)) = (1, 0) = \frac{1}{2}(1, 1) + \frac{1}{2}(1, -1)$. Also, $T((0, 1)) = (1, 1) = 1(1, 1) + 0(1, -1)$. Therefore, $[T]_\beta^\gamma = \begin{pmatrix} \frac{1}{2} & 1 \\ \frac{1}{2} & 0 \end{pmatrix}$.

Theorem 2.7. Let $T, U : V \rightarrow W$ are linear and let $a \in \mathbb{F}$. Then,

- $aT + U$ is linear.

- The set of all linear transformations from V to W (with addition of functions and scalar multiplication) forms a vector space over F .

We notate $\mathcal{L}(V, W)$ as the vector space of all linear transformations from V to W . Also, $\mathcal{L}(V) = \mathcal{L}(V, V)$.

Theorem 2.8. Let $T, U : V \rightarrow W$ be linear. Then,

- $[T + U]_{\beta}^{\gamma} = [T]_{\beta}^{\gamma} + [U]_{\beta}^{\gamma}$
- $[aT]_{\beta}^{\gamma} = a [T]_{\beta}^{\gamma}$.

2.3 Composition of linear transformations and matrix multiplication

Theorem 2.9. Let $T : V \rightarrow W$, $U : W \rightarrow Z$ be linear. Then $UT : V \rightarrow Z$ is linear.

Proof. Let $T : V \rightarrow W$, $U : W \rightarrow Z$ be linear. Then,

$$\begin{aligned} UT(au + v) &= U(T(au + v)) \\ &= U(aT(u) + T(v)) \\ &= aU(T(u)) + U(T(v)) \\ &= aUT(u) + UT(v). \end{aligned}$$

□

Theorem 2.10. Let $T, U_1, U_2 \in \mathcal{L}(V)$. Then

- $T(U_1 + U_2) = TU_1 + TU_2$ and $(U_1 + U_2)T = U_1T + U_2T$
- $T(U_1U_2) = (TU_1)U_2$
- $TI = IT = T$
- $a(U_1U_2) = (aU_1)U_2 = U_1(aU_2)$

Definition 5 Let A be an $m \times n$ matrix, B be an $n \times p$ matrix. We define AB to be the $m \times p$ matrix such that

$$(AB)_{ij} = \sum_{k=1}^n A_{ik}B_{kj}.$$

Note: $(AB)^t = B^tA^t$.

Theorem 2.11. Let $T : V \rightarrow W$. $U : W \rightarrow Z$ be linear and let α, β, γ be ordered bases in V, W, Z . Then

$$[UT]_{\alpha}^{\gamma} = [U]_{\beta}^{\gamma} [T]_{\alpha}^{\beta}.$$

Proof. (Outline)

- $\alpha = \{v_1, \dots, v_m\}$, $\beta = \{w_1, \dots, w_n\}$, $\gamma = \{z_1, \dots, z_p\}$.
- $A = [U]_{\beta}^{\gamma}$, $B = [T]_{\alpha}^{\beta}$
- $U(T(v_j)) = U(\sum_{k=1}^n B_{kj} w_k) = \sum_{k=1}^n B_{kj} U(w_k) = \sum_{k=1}^n B_{kj} \sum_{i=1}^p A_{ik} z_i = \sum_{i=1}^p C_{ij} z_i$
- Thus, by definition, the i, j th entry in $[UT]_{\alpha}^{\gamma}$ is $C_{ij} = \sum_{k=1}^n A_{ik} B_{kj}$.

Proof. Let $\alpha = \{v_1, \dots, v_m\}$, $\beta = \{w_1, \dots, w_n\}$ and $\gamma = \{z_1, \dots, z_p\}$. Let $A = [U]_{\beta}^{\gamma}$ and $B = [T]_{\alpha}^{\beta}$. Consider $[UT]_{\alpha}^{\gamma}$. Then

$$\begin{aligned} (UT)(v_j) &= U(T(v_j)) \\ &= U\left(\sum_{i=1}^m B_{ij} w_i\right) \\ &= \sum_{i=1}^m B_{ij} U(w_i) \\ &= \sum_{i=1}^m B_{ij} \cdot \sum_{k=1}^p A_{ki} z_k \\ &= \sum_{k=1}^p \left(\sum_{i=1}^m A_{ki} B_{ij}\right) z_k. \end{aligned}$$

If $C = [UT]_{\alpha}^{\gamma}$, then

$$C_{kj} = \sum_{i=1}^m A_{ki} B_{ij}.$$

□

Example: Let $\alpha = \{1, x, x^2\}$ and $\beta = \{1, x, x^2, x^3\}$ be standard bases for $P_2(\mathbb{R})$ and $P_3(\mathbb{R})$. Let $U : P_3(\mathbb{R}) \rightarrow P_2(\mathbb{R})$ and $T : P_2(\mathbb{R}) \rightarrow P_3(\mathbb{R})$ defined by $U(f(x)) = f'(x)$ and $T(f(x)) = \int_0^x f(t) dt$. From calculus, $UT = I$. We can see

$$\begin{aligned} U(1) &= 0 = 0 \cdot 1 + 0 \cdot x + 0 \cdot x^2 \\ U(x) &= 1 = 1 \cdot 1 + 0 \cdot x + 0 \cdot x^2 \\ U(x^2) &= 2x = 0 \cdot 1 + 2 \cdot x + 0 \cdot x^2 \\ U(x^3) &= 3x^2 = 0 \cdot 1 + 0 \cdot x + 3 \cdot x^2. \end{aligned}$$

Also,

$$\begin{aligned} T(1) &= x = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 \\ T(x) &= \frac{1}{2}x^2 = 0 \cdot 1 + 0 \cdot x + \frac{1}{2}x^2 + 0 \cdot x^3 \\ T(x^2) &= \frac{1}{3}x^3 = 0 \cdot 1 + 0 \cdot x + 0 \cdot x^2 + \frac{1}{3}x^3. \end{aligned}$$

Thus,

$$\begin{aligned} [UT]_{\alpha}^{\alpha} &= [U]_{\beta}^{\alpha} [T]_{\alpha}^{\beta} \\ &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{3} \end{pmatrix} \\ &= I. \end{aligned}$$

Theorem 2.12. Let A be an $m \times n$ matrix, B and C be $n \times p$ matrices, D, E be $q \times m$ matrices. Then

- $A(B + C) = AB + AC$, $(D + E)A = DA + EA$
- $a(AB) = (aA)B = A(aB)$
- $I_m A = A = A I_n$
- If V is n -dimensional with ordered basis β , then $[I_V]_{\beta} = I_n$

Theorem 2.13. Let A be an $m \times n$ matrix, B be an $n \times p$ matrix. Let u_j, v_j denote the j th columns of AB and B . Then

- (a) $u_j = Av_j$
- (b) $v_j = Be_j$

Theorem 2.14. Let V, W be finite-dimensional vector spaces with ordered bases β and γ and let $T : V \rightarrow W$ be linear. Then for $u \in V$

$$[T(u)]_{\gamma} = [T]_{\beta}^{\gamma} [u]_{\beta}.$$

Proof. Let V, W be finite-dimensional vector spaces with ordered bases β and γ and let $T : V \rightarrow W$ be linear. Fix $u \in V$. Let $f : \mathbb{R} \rightarrow V$ and $g : \mathbb{R} \rightarrow W$ where $f(a) = au$

and $g(a) = aT(u)$. Let $\alpha = \{1\}$ be the standard basis for \mathbb{R} . Note that, g and f are linear transformations. Then,

$$\begin{aligned}
 [T(u)]_\gamma &= [g(1)]_\gamma \\
 &= [g]_\alpha^\gamma \\
 &= [Tf]_\alpha^\gamma \\
 &= [T]_\beta^\gamma [f]_\alpha^\beta \\
 &= [T]_\beta^\gamma [f(1)]_\beta \\
 &= [T]_\beta^\gamma [u]_\beta.
 \end{aligned}$$

□


Let $L_A : \mathbb{F}^n \rightarrow \mathbb{F}^m$ be given by $L_A(x) = Ax$ where A is an $m \times n$ matrix.

Theorem 2.15. Let A be an $m \times n$ matrix. Then L_A is linear. Moreover, if $B \in M_{m \times n}(\mathbb{F})$ and β and γ are the standard ordered bases of \mathbb{F}^n and \mathbb{F}^m , then

- (a) $[L_A]_\beta^\gamma = A$
- (b) $L_A = L_B$ if and only if $A = B$
- (c) $L_{A+B} = L_A + L_B$, $L_{aA} = aL_A$
- (d) If $T : \mathbb{F}^n \rightarrow \mathbb{F}^m$ is linear, then there exists unique matrix C (namely $C = [T]_\beta^\gamma$) such that $T = L_C$.
- (e) If E is an $n \times p$ matrix, then $L_{AE} = L_A L_E$
- (f) If $m = n$, $L_{I_n} = I_{\mathbb{F}^n}$.

Theorem 2.16. Let A, B, C be such that $A(BC)$ is defined then $(AB)C = A(BC)$.

Tournaments: There are n players $\{1, \dots, n\}$ for every $i \neq j$ there is exactly one game between i and j which results in i winning or j winning. Let A be the incidence matrix of a tournament where we put $A_{ij} = 1$ if i wins with j and 0 otherwise. Show that $A^2 + A$ contains a column such that each entry but the diagonal is at least one.



2.4 Invertibility and isomorphisms

Definition 6 Let $T : V \rightarrow W$ be linear. A function $U : W \rightarrow V$ is called an *inverse* of T if $UT = I_V$, $TU = I_W$. If T has an inverse, then it's called *invertible*.

Note:

- If T is invertible, then the inverse is unique, denoted by T^{-1} .
- $(UT)^{-1} = T^{-1}U^{-1}$
- $(T^{-1})^{-1} = T$
- T is invertible if and only if T is a bijection.
- $T : V \rightarrow W$, V, W are finite-dimensional of equal dimensions.
- T is invertible if and only if $\text{rank}(T) = \dim(V)$.

Theorem 2.17. Let $T : V \rightarrow W$ be linear and invertible. Then $T^{-1} : W \rightarrow V$ is linear.

Proof. Suppose $T : V \rightarrow W$ is linear and invertible. Then $T^{-1} : W \rightarrow V$. Let $y_1, y_2 \in W$. Since T is bijective, there exists unique $x_1, x_2 \in V$ such that $T(x_1) = y_1$ and

$T(x_2) = y_2$. Let $c \in \mathbb{F}$. Then

$$\begin{aligned} T^{-1}(cy_1 + y_2) &= T^{-1}(cT(x_1) + T(x_2)) \\ &= T^{-1}(T(cx_1 + x_2)) \\ &= cx_1 + x_2 \\ &= cT^{-1}(y_1) + T^{-1}(y_2). \end{aligned}$$

□

Definition 7 Let A be an $n \times n$ matrix. Then A is *invertible* if there exists an $n \times n$ matrix B such that $AB = BA = I$. Then B is called the inverse of A denoted by A^{-1} .

Lemma 19. Let T be an invertible transformation from V to W . Then V is finite-dimensional if and only if W is. In this case $\dim(V) = \dim(W)$.

Proof. Let T be an invertible transformation from V to W . Then T must be bijective. Suppose V is finite-dimensional. Let $\beta = \{v_1, \dots, v_n\}$ be a basis for V . Then $\dim(V) = n$. Then $T(\beta) = \{T(v_1), \dots, T(v_n)\}$ generates W because T is surjective. Let $w \in W$. Then there exists $v \in V$ such that $T(v) = w$. Then $v = \sum_{i=1}^n a_i v_i$. Thus, $T(v) = \sum_{i=1}^n a_i T(v_i)$. As a result, $\dim(W) \leq n = \dim(V)$. However, $T^{-1} : W \rightarrow V$ is also linear. Then the same argument implies $\dim(V) \leq \dim(W)$. Therefore, $\dim(V) = \dim(W)$. □

Theorem 2.18. Let V, W be finite-dimensional with ordered bases β and γ . Let $T : V \rightarrow W$ be linear. Then T is invertible if and only if $[T]_{\beta}^{\gamma}$ is invertible. Furthermore $[T^{-1}]_{\gamma}^{\beta} = ([T]_{\beta}^{\gamma})^{-1}$.

Proof. (Outline)

- If T is invertible, then by the lemma $\dim(V) = \dim(W)$ and the fact that $[T]_{\beta}^{\gamma}$ follows from previous facts and $T^{-1}T = I_V$.
- If $[T]_{\beta}^{\gamma}$ is invertible then it has an inverse B and so there is a transformation $U : W \rightarrow V$ with $B = [U]_{\gamma}^{\beta}$. Now check that $UT = I_V$.

Proof. Suppose T is invertible. Then by the lemma $\dim(V) = \dim(W) = n$, and in addition, there exists $U : W \rightarrow V$ such that $TU = I_W$ and $UT = I_V$. Let β, γ be bases for V, W respectively. Then

$$I = [I_V]_{\beta} = [UT]_{\beta} = [U]_{\gamma}^{\beta} [T]_{\beta}^{\gamma}.$$

In the same way,

$$[I_W]_{\gamma} = [TU]_{\gamma} = [T]_{\beta}^{\gamma} [U]_{\gamma}^{\beta}.$$

Thus, $[U]_\gamma^\beta = \left([T]_\beta^\gamma\right)^{-1}$.

Now, assume $[T]_\beta^\gamma$ is invertible. Let $A = [T]_\beta^\gamma$. Then A is an $n \times n$ matrix. Let $\beta = \{v_1, \dots, v_n\}$ and $\gamma = \{w_1, \dots, w_n\}$. Since A is invertible, there exists B such that $AB = I_n = BA$. Then there exists unique linear transformation $U : W \rightarrow V$ such that $U(w_j) = \sum_{i=1}^n B_{ij}v_i$. Then $[U]_\gamma^\beta = B$. Thus $[TU]_\gamma = [T]_\beta^\gamma [U]_\gamma^\beta = A \cdot B = I_n = [I_W]_\gamma$. Also, $[UT]_\beta = B \cdot A = I_n = [I_V]_\beta$. Thus $TU = I_W$ and $UT = I_V$. \square

Definition 8 Vector space V is isomorphic to W if there is an invertible linear transformation $T : V \rightarrow W$.

Example: Let $T : P_3(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ be defined by

$$T(f) = \begin{pmatrix} f(1) & f(2) \\ f(3) & f(4) \end{pmatrix}.$$

Then

(1) T is linear.

$$\begin{aligned} T(f+g) &= T(f) + T(g) \\ T(cf) &= cT(f) \end{aligned}$$

(2) T is injective. (This is because $N(T) = \{0\}$.)

(3) $\dim(P_3(\mathbb{R})) = 4$ and $\dim(M_{2 \times 2}(\mathbb{R})) = 4$

(4) T is surjective

(5) T is bijective

(6) $P_3(\mathbb{R}) \cong M_{2 \times 2}(\mathbb{R})$

Theorem 2.19. Let V, W be finite-dimensional over F . Then V is isomorphic to W if and only if $\dim(V) = \dim(W)$.

Proof.

(\implies): Suppose $V \cong W$. By the lemma, $\dim(V) = \dim(W)$.

(\impliedby): Suppose $\dim(V) = \dim(W)$. Let $\beta = \{v_1, \dots, v_n\}$ be a basis for V and let $\gamma = \{w_1, \dots, w_n\}$ be a basis for W . Then there exists a unique linear transformation $T : V \rightarrow W$ such that $T(v_i) = w_i$. Then T is surjective. Since $\dim(V) = \dim(W)$, T is injective. Thus T is invertible. Therefore, $V \cong W$. \square

Theorem 2.20. Let V, W be finite-dimensional over F of dimensions n and m . Let β, γ be ordered bases for V and W . Then $\Phi : \mathcal{L}(V, W) \rightarrow M_{m \times n}(\mathbb{F})$ given by $\Phi(T) = [T]_{\beta}^{\gamma}$ for $T \in \mathcal{L}(V, W)$ is an isomorphism.

Proof. Let $\Phi(T) = [T]_{\beta}^{\gamma}$.

(1) Let $\phi : \mathcal{L}(V, W) \rightarrow M_{m \times n}(\mathbb{F})$. Then Φ is linear.

$$\begin{aligned}\Phi(aT + U) &= [aT + U]_{\beta}^{\gamma} \\ &= [aT]_{\beta}^{\gamma} + [U]_{\beta}^{\gamma} \\ &= a [T]_{\beta}^{\gamma} + [U]_{\beta}^{\gamma}\end{aligned}$$

(2) For every $A \in M_{m \times n}(\mathbb{F})$ there exists unique transformation T such that $\Phi(T) = A = [A_{ij}]$. Let $\beta = \{v_1, \dots, v_n\}$ and $\gamma = \{w_1, \dots, w_m\}$. Then there exists unique transformation $T : V \rightarrow W$ such that

$$T(v_j) = \sum_{i=1}^m A_{ij} w_i,$$

for $j = 1, \dots, n$ and $i = 1, \dots, m$. Thus

$$[T]_{\beta}^{\gamma} = A.$$

Thus $\Phi(T) = [T]_{\beta}^{\gamma} = A$.

□

Definition 9 Let β be an ordered basis for an n -dimensional vector space V over F . The *standard representation of V with respect to β* is the function $\phi_{\beta} : V \rightarrow \mathbb{F}^n$ defined as $\phi_{\beta}(x) = [x]_{\beta}$.

Theorem 2.21. Let V be a finite-dimensional vector space with ordered basis β . Then ϕ_{β} is an isomorphism.

Note: For vector fields V and W with basis β and γ , respectively, we have the following schematic representation:

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \phi_{\beta} \downarrow & & \downarrow \phi_{\gamma} \\ \mathbb{F}^n & \xrightarrow{L_A} & \mathbb{F}^m \end{array}$$

Where $L_A(x) = Ax$ and $A = [T]_{\beta}^{\gamma}$, we have

$$L_A \phi_{\beta} = \phi_{\gamma} T.$$

That is,

$$[T]_{\beta}^{\gamma} [u]_{\beta} = [T(u)]_{\gamma}.$$

2.5 The change of coordinate matrix

Theorem 2.22. Let β, β' be two ordered bases for a finite-dimensional vector space V and let $Q = [I_V]_{\beta'}^{\beta}$. Then

- (a) Q is invertible.
- (b) For any $v \in V$, $[v]_{\beta} = Q [v]_{\beta'}$.

Proof.

- (a) We know $[I_V]_{\beta'}^{\beta}$ is invertible because $I_V : V \rightarrow V$ is invertible. So, $I_V^{-1} = I_V$.
- (b) We have $[u]_{\beta} = Q \cdot [u]_{\beta'}$. So, $[u]_{\beta} = [I_V(u)]_{\beta} = [I_V]_{\beta'}^{\beta} [u]_{\beta'}$.

□

Example: Let $\beta = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$ and $\beta' = \left\{ \begin{pmatrix} 2 \\ 4 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\}$. Then

$$\begin{aligned} \begin{pmatrix} 2 \\ 4 \end{pmatrix} &= 3 \begin{pmatrix} 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \\ \begin{pmatrix} 3 \\ 1 \end{pmatrix} &= 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \\ [I_V]_{\beta'}^{\beta} &= \begin{pmatrix} 3 & 2 \\ -1 & 1 \end{pmatrix}, \\ \left[\begin{pmatrix} 2 \\ 4 \end{pmatrix} \right]_{\beta} &= Q \left[\begin{pmatrix} 2 \\ 4 \end{pmatrix} \right]_{\beta'} \\ &= Q \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} 3 \\ -1 \end{pmatrix}. \end{aligned}$$

Q is called the change of coordinate matrix. A linear operator on V is the linear transformation from V to V .

Theorem 2.23. Let T be a linear operator on a finite-dimensional vector space V . Let β, β' be ordered bases for V . Suppose $Q = [I_V]_{\beta'}^{\beta}$. Then

$$[T]_{\beta'} = Q^{-1} [T]_{\beta} Q.$$

Proof. Suppose $Q = [I_V]_{\beta'}^{\beta}$. Note that $I_V Q = Q = Q I_V$. Then,

$$\begin{aligned} Q [T]_{\beta'} &= [I_V]_{\beta'}^{\beta} [T]_{\beta'} \\ &= [I_V T]_{\beta'}^{\beta} \\ &= [T I_V]_{\beta'}^{\beta} \\ &= [T]_{\beta} [I_V]_{\beta'}^{\beta} \\ &= [T]_{\beta} \cdot Q. \end{aligned}$$

So, $Q^{-1} Q [T]_{\beta'} = Q^{-1} [T]_{\beta} Q$. Therefore, $[T]_{\beta'} = Q^{-1} [T]_{\beta} Q$. \square

Corollary 26 Let $A \in M_{n \times n}(\mathbb{F})$ and let $\gamma = \{u_1, \dots, u_n\}$ be an ordered basis for \mathbb{F}^n . Then $[L_A]_{\gamma} = Q^{-1} A Q$ where Q is the matrix with the j th column equal to u_j .

Note: We say B is **similar to** A if there exists an invertible matrix C such that

$$B = C^{-1} A C.$$

Example: Let $A = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{pmatrix}$. Find A^{100} .

Solution. Let $\gamma = \left\{ \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\}$.

(1) Find $[L_A]_{\gamma}$.

$$\begin{aligned} L_A \left(\begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix} \right) &= A \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}, \\ L_A \left(\begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right) &= A \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \\ 2 \end{pmatrix}, \\ L_A \left(\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right) &= A \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}. \end{aligned}$$

Thus

$$[L_A]_{\gamma} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

(2) Note, $Q = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$ and $Q^{-1} = \begin{pmatrix} 1 & 0 & 1 \\ -1 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}$. Then $[L_A]_\gamma = Q^{-1}AQ$. So, $A = Q[L_A]_\gamma Q^{-1}$. Then we have

$$\begin{aligned} A^{100} &= \left(Q[L_A]_\gamma Q^{-1} \right)^{100} \\ &= Q[L_A]_\gamma Q^{-1} Q[L_A]_\gamma Q^{-1} \dots Q[L_A]_\gamma Q^{-1} \\ &= Q[L_A]_\gamma^{100} Q^{-1} \\ &= Q \begin{pmatrix} 1^{100} & 0 & 0 \\ 0 & 2^{100} & 0 \\ 0 & 0 & 2^{100} \end{pmatrix} Q^{-1}. \end{aligned}$$

2.6 Dual spaces

- Linear functional on V – linear transformation from V to F .
- $V^* = \mathcal{L}(V, F)$
- $V^{**} = (V^*)^*$

$$\dim(V^*) = \dim(V)$$

Let $\beta = \{x_1, \dots, x_n\}$ for $v \in V$ let $[v]_\beta = (a_1, \dots, a_n)^T$. Define

$$f_i(v) = a_i.$$

Example:

(1) Let $V =$ continuous functions $f : [0, 2\pi] \rightarrow \mathbb{R}$. Let $g \in V$. Define $h : V \rightarrow \mathbb{R}$ by

$$h(x) = \frac{1}{2\pi} \int_0^{2\pi} x(t)g(t)dt.$$

Note, h is linear. If $g(t) = \sin(nt)$ or $g(t) = \cos(nt)$, then $h(x)$ is called the n th Fourier coefficient of x .

(2) Let $V = M_{n \times n}(\mathbb{F})$ and $f : V \rightarrow \mathbb{F}$ where $f(A) = \text{tr}(A)$.

Theorem 2.24. Let V be a finite-dimensional vector space with ordered basis $\beta = \{x_1, \dots, x_n\}$ and let $\beta^* = \{f_1, \dots, f_n\}$. Then β^* is an ordered basis for V^* and for $f \in V^*$ where

$$f = \sum_{i=1}^n f(x_i)f_i.$$

Note, $f_i : V \rightarrow \mathbb{F}$ is defined as $f_i(x_j) = \delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{else} \end{cases}$.

Proof. (Outline)

- Enough to show $f = \sum f(x_i)f_i$.
- Let $F := \sum f(x_i)f_i$. Then $F(x_j) = f(x_j)$ for every j .

Proof. Let V be a finite-dimensional vector space with ordered basis $\beta = \{x_1, \dots, x_n\}$ and let $\beta^* = \{f_1, \dots, f_n\} \subseteq V^*$ where $n = \dim(V) = \dim(V^*)$. Since $|\beta^*| = n = \dim(V^*)$, it is enough to show that β^* generates V^* . To that end, we will argue that for $f \in V^*$,

$$f = \sum_{i=1}^n f(x_i)f_i.$$

Let $g = \sum_{i=1}^n f(x_i)f_i$. Then

$$\begin{aligned} g(x_j) &= \left(\sum_{i=1}^n f(x_i)f_i \right) (x_j) \\ &= \sum_{i=1}^n f(x_i)f_i(x_j) \\ &= \sum_{i=1}^n f(x_i)\delta_{ij} \\ &= f(x_j). \end{aligned}$$

Thus, $g(x_j) = f(x_j)$ for every $x_j \in \beta$. Therefore $g = f$. □

Definition 10 An ordered basis $\beta^* = \{f_1, \dots, f_n\}$ for V^* such that $f_i(x_j) = \delta_{ij}$ is called the *dual basis* of $\beta = \{x_1, \dots, x_n\}$.

Theorem 2.25. Let V, W be finite-dimensional vector spaces over \mathbb{F} with ordered bases β and γ . Let $T : V \rightarrow W$ be linear. Then $T^t : W^* \rightarrow V^*$ given by $T^t(g) = gT$ is linear and

$$[T^t]_{\gamma^*}^{\beta^*} = ([T]_{\beta}^{\gamma})^t.$$

Proof. (Outline)

- It's easy to see that T^t is a linear transformation from W^* to V^* .
- Let $\beta = \{x_1, \dots, x_n\}$, $\gamma = \{y_1, \dots, y_m\}$, $\beta^* = \{f_1, \dots, f_n\}$, $\gamma^* = \{g_1, \dots, g_m\}$, $A = [T]_{\beta}^{\gamma}$.

- The j th column of $[T^t]_{\gamma^*}^{\beta^*}$ is $T^t(g_j)$ which is

$$\sum_{k=1}^n (g_j T)(x_k) f_k.$$

- Thus the i, j -th entry is $(T^t(g_j))(x_i)$ which is A_{ji} .

Proof. Let $T^t : W^* \rightarrow V^*$, $T^t(g) = gT$.

(1) Note that $gT : V \rightarrow \mathbb{F}$ and gT is a linear transformation. Thus $T^t(g) \in V^*$.

(2) We have T^t is linear. So,

$$\begin{aligned} T^t(cg + h) &= (cg + h)T \\ &= cgT + hT \\ &= cT^t(g) + T^t(h). \end{aligned}$$

(3) Lastly, $[T^t]_{\gamma^*}^{\beta^*} = \left([T]_{\beta}^{\gamma}\right)^t$. Let $\beta = \{x_1, \dots, x_n\}$, $\gamma = \{y_1, \dots, y_m\}$, $\beta^* = \{f_1, \dots, f_n\}$, and $\gamma^* = \{g_1, \dots, g_m\}$. Let $A = [T]_{\beta}^{\gamma}$. To obtain the j th column of $[T]_{\gamma^*}^{\beta^*}$,

$$\begin{aligned} T^t(g_j) &= g_j T \\ &= \sum_{k=1}^n (g_j T)(x_k) f_k. \end{aligned}$$

Thus, the i, j th entry of $[T]_{\gamma^*}^{\beta^*}$ is

$$\begin{aligned} (g_j T)(x_i) &= g_j(T(x_i)) \\ &= g_j \left(\sum_{k=1}^m A_{ki} y_k \right) \\ &= \sum_{k=1}^m A_{ki} g_j(y_k) & (g_j(y_k) = \delta_{kj}) \\ &= A_{ji}. \end{aligned}$$

□

For $x \in V$ let $\hat{x} : V^* \rightarrow \mathbb{F}$ given by $\hat{x}(f) = f(x)$.

Theorem 2.26. Let V be finite-dimensional and let $\psi : V \rightarrow V^{**}$ be given by $\psi(x) = \hat{x}$. Then ψ is an isomorphism.

3 Elementary matrix operations and systems of linear equations

3.1 Elementary matrix operations and elementary matrices

Definition 1 Let A be a matrix. Elementary row operations:

- Interchange any two rows of A
- Add a scalar multiple of a row of A to another row
- Multiply any row of A by a non-zero scalar

Note: The same can be done for columns.

Definition 2 An $n \times n$ elementary matrix is a matrix obtained from I_n by an elementary operation. Its type is the type of the operation performed.

Theorem 3.1. Let $A \in M_{m \times n}(\mathbb{F})$ and suppose B is obtained by performing an elementary row (column) operation. Then there exists an $m \times m$ ($n \times n$) elementary matrix E such that $B = EA$ ($B = AE$).

Example: Note, $\begin{pmatrix} 1 & -7 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ is an elementary matrix.

$$\begin{pmatrix} a-7x & b-7y & c-7z \\ x & y & z \\ u & v & w \end{pmatrix} = \begin{pmatrix} 1 & -7 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Theorem 3.2. Elementary matrices are invertible and the inverse of an elementary matrix is an elementary matrix of the same type.

Example: Let $A = \begin{pmatrix} 1 & -7 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$. Then

$$\begin{aligned} AA^{-1} &= \begin{pmatrix} 1 & -7 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 7 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

3.2 The rank of a matrix and matrix inverse

Definition 3 Let $A \in M_{m \times n}(\mathbb{F})$. The $\text{rank}(A)$ is the rank of $L_A : \mathbb{F}^n \rightarrow \mathbb{F}^m$. Also $\text{rank}(L_A) = \dim(R(L_A))$.

Theorem 3.3. Let $T : V \rightarrow W$ be linear and let β, γ be ordered bases for V and W . Then

$$\text{rank}(T) = \text{rank} \left([T]_{\beta}^{\gamma} \right).$$

Recall, where V and W are vector spaces with bases β and γ , respectively, and $A = [T]_{\beta}^{\gamma}$, we have

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \updownarrow & & \updownarrow \\ \mathbb{F}^n & \xrightarrow{L_A} & \mathbb{F}^m \end{array}$$

Corollary 3.4.1 Elementary operations are rank-preserving.

Theorem 3.4. Let A be an $m \times n$ matrix. If P and Q are invertible $m \times m$ and $n \times n$ matrices, then

- $\text{rank}(AQ) = \text{rank}(A)$
- $\text{rank}(PA) = \text{rank}(A)$

and so $\text{rank}(PAQ) = \text{rank}(A)$.

Proof. (Outline)

- $R(L_{AQ}) = R(L_A)$ because $L_Q(\mathbb{F}^n) = \mathbb{F}^n$.
- $\dim(L_P(L_A(\mathbb{F}^n))) = \dim(L_A(\mathbb{F}^n))$ because $L_P : \mathbb{F}^m \rightarrow \mathbb{F}^m$ is an isomorphism.

Proof.

(1) Note,

$$\begin{aligned} \text{rank}(AQ) &= \text{rank}(L_{AQ}) \\ &= \dim(R(L_{AQ})). \end{aligned}$$

So,

$$\begin{aligned} R(L_{AQ}) &= R(L_A L_Q) \\ &= L_A L_Q(\mathbb{F}^n) \\ &= L_A(L_Q(\mathbb{F}^n)) \\ &= L_A(\mathbb{F}^n) && (\text{Since } L_Q(\mathbb{F}^n) = \mathbb{F}^n) \\ &= R(L_A). \end{aligned}$$

Therefore, $\dim(R(L_{AQ})) = \dim(R(L_A)) = \text{rank}(A)$.

(2) We have to show $\text{rank}(PA) = \text{rank}(A)$. So,

$$\begin{aligned}\text{rank}(PA) &= \dim(R(L_{PA})) \\ &= \dim(L_{PA}(\mathbb{F}^n)) \\ &= \dim(L_P(L_A(\mathbb{F}^n))).\end{aligned}$$

Since $L_P : L_A(\mathbb{F}^n) \rightarrow L_P(L_A(\mathbb{F}^n))$, then L_P is an isomorphism. Thus, $\dim(L_P(L_A(\mathbb{F}^n))) = \dim(L_A(\mathbb{F}^n)) = \text{rank}(A)$. Therefore, $\text{rank}(PA) = \text{rank}(A)$. □

Theorem 3.5. The rank of a matrix equals the maximum number of its linearly independent columns.

Proof. (Outline)

- $R(L_A) = \text{span}(L_A(\{e_1, \dots, e_n\}))$ and $L_A(e_j)$ is the j th column of A .

Proof. Let $A \in M_{m \times n}(\mathbb{F}^n)$. Then

$$\begin{aligned}\text{rank}(A) &= \dim(R(L_A)) \\ &= \dim(L_A(\mathbb{F}^n)).\end{aligned}$$

Let β be the standard basis for \mathbb{F}^n . We have $\text{span}(\beta) = \mathbb{F}^n$. Therefore,

$$\begin{aligned}R(L_A) &= \text{span}(L_A(\beta)) \\ &= \text{span}(\{L_A(e_1), \dots, L_A(e_n)\}).\end{aligned}$$

We have $L_A(e_i) = a_i$ where a_i is the i th column of A . Thus, $R(L_A) = \text{span}(\{a_1, \dots, a_n\})$. Thus, $\dim(R(L_A))$ is the maximum number of linearly independent columns. □

Example: Find the rank of

$$A = \begin{pmatrix} 1 & 2 & 1 \\ 1 & 0 & 3 \\ 1 & 1 & 1 \end{pmatrix}.$$

Solution.

$$\begin{aligned}
 A &\xrightarrow{\text{Row Op.}} \begin{pmatrix} 1 & 2 & 1 \\ 0 & -2 & 2 \\ 0 & -1 & 0 \end{pmatrix} \\
 &\xrightarrow{\text{Col. Op.}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & -1 & 0 \end{pmatrix} \\
 &\xrightarrow{\text{Row Op.}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 2 \end{pmatrix} \\
 &\xrightarrow{\text{Row Op.}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.
 \end{aligned}$$

Therefore, $\text{rank}(A) = 3$.

Theorem 3.6. Let A be an $m \times n$ matrix of rank r . Then $r \leq \min\{m, n\}$ and A can be transformed to

$$D = \begin{pmatrix} I_r & 0_1 \\ 0_2 & 0_3 \end{pmatrix}$$

using a finite number of elementary row and column operations.

Proof. (Outline) Induction on m . In the inductive step use row and column operations to reduce to

$$\left(\begin{array}{c|ccc} 1 & 0 & \dots & 0 \\ 0 & & & \\ \hline & & B' & \\ 0 & & & \end{array} \right).$$

Proof. If A is the zero matrix, then $\text{rank}(A) = 0$ and $D = A$, thus $r = 0$. Suppose A is non-zero. We will use induction on m .

(Base Step) Let $m = 1$. Then A has one row. Then by applying elementary column operations, we can transform A to $(1 \ 0 \ \dots \ 0)$ and so $r = 1$ and $\text{rank}(A) = 1$.

(Induction Step) Suppose $n \geq 2$. If $n = 1$, then A can be transformed into $\begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$.

So, $r = 1 = \text{rank}(A)$. Let $n \geq 2$. So, there exists A_{ij} such that $A_{ij} \neq 0$ and we can

transform A so that A_{ij} is in position $(1,1)$. Therefore, we can transform A to

$$\left(\begin{array}{c|ccc} 1 & 0 & \dots & 0 \\ \hline 0 & & & \\ & B & & \\ 0 & & & \end{array} \right)$$

and $\text{rank}(B) = \text{rank}(A) - 1 = r - 1$. By the inductive hypothesis, $r - 1 \leq m - 1$ and $r - 1 \leq n - 1$ and B can be transformed to

$$\begin{pmatrix} I_{r-1} & 0_4 \\ 0_5 & 0_6 \end{pmatrix}.$$

Thus, A can be transformed to

$$\begin{pmatrix} I_r & 0_1 \\ 0_2 & 0_3 \end{pmatrix}$$

for some 0_i .

□

Note, if $m = n$ and $\text{rank}(A) = n$, then $B = I_n$. The converse of this statement is also true.

Corollary 7 Let A be an $m \times n$ matrix of rank r . Then there exist invertible matrices B and C of sizes $m \times m$ and $n \times n$ such that $D = BAC$.

As a consequence of the above corollary, A is invertible if and only if $\text{rank}(A) = n$.

Corollary 8 Let A be an $m \times n$ matrix. Then

- $\text{rank}(A^t) = \text{rank}(A)$
- $\text{rank}(A)$ is equal to the dimension of the row space of A
- dimension of the row space is equal to the dimension of the column space.

Proof. We will show $\text{rank}(A^t) = \text{rank}(A)$. By the Theorem 3.6, there exists invertible matrices B and C such that $D = BAC$. Then $D^t = (BAC)^t = C^t A^t B^t$. Also, B^t and C^t are invertible. Recall, $(B^t)^{-1} = (B^{-1})^t$. We have $\text{rank}(D^t) = r = \text{rank}(D)$ and $\text{rank}(A) = \text{rank}(D) = \text{rank}(D^t) = \text{rank}(A^t)$. □

Corollary 9 Every invertible matrix is a product of elementary matrices.

Proof. Suppose A is invertible. Then there exists invertible matrices B and C such that $D = BAC$. Thus, D is invertible, which implies $D = I_n$. Also, $B = E_1 \dots E_p$ and $C = G_1 \dots G_q$, where E_i and G_j are elementary. Therefore, $BAC = I_n$ and $A = B^{-1}C^{-1}$. Thus,

$$\begin{aligned} A &= (E_1 \dots E_p)^{-1}(G_1 \dots G_q)^{-1} \\ &= E_p^{-1} \dots E_1^{-1} G_q^{-1} \dots G_1^{-1}. \end{aligned}$$

□

Theorem 3.7. Let $T : V \rightarrow W$ and $U : W \rightarrow Z$ be linear transformations on finite-dimensional vector spaces. Let A, B be matrices such that AB is defined. Then

- (a) $\text{rank}(UT) \leq \text{rank}(U)$
- (b) $\text{rank}(UT) \leq \text{rank}(T)$
- (c) $\text{rank}(AB) \leq \text{rank}(A)$
- (d) $\text{rank}(AB) \leq \text{rank}(B)$

Proof. (Outline)

- For (a), $R(UT) = U(R(T)) \subseteq U(W) = R(U)$
- (c) and (d) follow from (a) and discussion of the transpose
- (b) follows from the previous by considering matrix representations.

Proof. Let $T : V \rightarrow W$ and $U : W \rightarrow Z$ be linear transformations on finite-dimensional vector spaces. Let A, B be matrices such that AB is defined. For (a), we have $R(T) = T(V) \subseteq W$. Then

$$\begin{aligned} R(UT) &= (UT)(V) \\ &= U(T(V)) \\ &\subseteq U(W) \\ &= R(U). \end{aligned}$$

Thus, $\text{rank}(UT) = \dim(R(UT)) \leq \dim(R(U)) = \text{rank}(U)$.

Now, for (c),

$$\begin{aligned} \text{rank}(AB) &= \text{rank}(L_{AB}) \\ &= \text{rank}(L_A L_B) \\ &\leq \text{rank}(L_A) && \text{(By (a))} \\ &= \text{rank}(A). \end{aligned}$$

Now, for (d),

$$\begin{aligned}
 \text{rank}(AB) &= \text{rank}((AB)^t) \\
 &= \text{rank}(B^t A^t) \\
 &\leq \text{rank}(B^t) \\
 &= \text{rank}(B).
 \end{aligned}
 \tag{By (c)}$$

Now, for (b), let α, β, γ be ordered bases in V, W , and Z . Let $A = [U]_\beta^\gamma$ and $B = [T]_\alpha^\beta$. Then $AB = [UT]_\alpha^\gamma$. Thus,

$$\begin{aligned}
 \text{rank}(UT) &= \text{rank}([UT]_\alpha^\gamma) \\
 &= \text{rank}(AB) \\
 &\leq \text{rank}(B) \\
 &= \text{rank}([T]_\alpha^\beta) \\
 &= \text{rank}(T).
 \end{aligned}
 \tag{By (b)}$$

□

Observations: A is an invertible $n \times n$ matrix if and only if $(A|I_n)$ can be transformed into $(I_n|B)$ by elementary row operations, in this case $B = A^{-1}$.

So, $C = (A|I_n)$. Then $A^{-1}C = (A^{-1}A|A^{-1}) = (I_n|A^{-1})$. Consequently, $A^{-1} = E_1 \dots E_p$ where E_i is elementary. Thus, $(E_1 \dots E_p)C = (I_n|A^{-1})$. So, C can be converted to $(I_n|A^{-1})$ by elementary row operations.

Further, suppose we can transform C to $(I_n|B)$ by using elementary row operations. So, $E_1 \dots E_p(A|I_n) = (I_n|B)$. Let $M = E_1 \dots E_p$. Then, $MA = I_n$ and $M = B$. So, $MA = I_n$ implies $M = A^{-1} = B$. Finally, if A is not invertible, then by Theorem 3.6, $r < n$.

Example: Determine if $\begin{pmatrix} 4 & 0 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ is invertible and find its inverse.

Solution.

$$\begin{aligned}
 \left(\begin{array}{ccc|ccc} 4 & 0 & 1 & 1 & 0 & 0 \\ 2 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 \end{array} \right) &\xrightarrow{\text{Row Op.}} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 0 & 0 & 1 \\ 2 & 1 & 1 & 0 & 1 & 0 \\ 4 & 0 & 1 & 1 & 0 & 0 \end{array} \right) \\
 &\xrightarrow{\text{Row Op.}} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 0 & 0 & 1 \\ 0 & -1 & -1 & 0 & 1 & -2 \\ 0 & -4 & -3 & 1 & 0 & -4 \end{array} \right) \\
 &\xrightarrow{\text{Row Op.}} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & -1 & 2 \\ 0 & 4 & 3 & -1 & 0 & 4 \end{array} \right) \\
 &\vdots \\
 &\xrightarrow{\text{Row Op.}} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & 0 & -1 & 3 & -2 \\ 0 & 0 & 1 & 1 & -4 & 4 \end{array} \right)
 \end{aligned}$$

Example: Let $T : P_2(\mathbb{R}) \rightarrow P_2(\mathbb{R})$ be defined by $T(f) = f + f' + f''$. Find T^{-1} .

Solution. Let $\beta = \{1, x, x^2\}$ be the standard ordered basis for $P_2(\mathbb{R})$. Then

$$\begin{aligned}
 T(1) &= 1 + 0 + 0 = 1 \\
 T(x) &= x + 1 = 1 + x \\
 T(x^2) &= x^2 + 2x + 2 = 2 + 2x + x^2.
 \end{aligned}$$

We have

$$[T^{-1}]_{\beta} = [T]_{\beta}^{-1} = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}.$$

Therefore,

$$\begin{aligned}
 T^{-1}(a_0 + a_1x + a_2x^2) &= \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_0 \\ a_1 \\ a_2 \end{pmatrix} \\
 &= \begin{pmatrix} a_0 - a_1 \\ a_1 - 2a_2 \\ a_2 \end{pmatrix} \\
 &= (a_0 - a_1) + (a_1 - 2a_2)x + a_2x^2.
 \end{aligned}$$

3.3 Systems of linear equations (theoretical aspect)

- $Ax = b$ is consistent if it has at least one solution and inconsistent otherwise.

- $Ax = 0$ is called a homogeneous system and $Ax = b$ for $b \neq 0$ a nonhomogeneous system.

Theorem 3.8. Let $Ax = 0$ be a homogeneous system over F in n unknowns and let K be the set of solutions. Then $K = N(L_A)$ and so K is a subspace of \mathbb{F}^n and $\dim(K) = n - \text{rank}(A)$.

Corollary 12 If $m < n$, then $Ax = 0$ has a non-zero solution.

Proof. We have $\dim(K) = \dim(N(L_A)) = n - \text{rank}(A)$. We know $\text{rank}(A) \leq m$. So, $\dim(K) = n - \text{rank}(A) \geq n - m > 0$. \square

Theorem 3.9. Let K be the solution set to $Ax = b$ and K_H be the solution set to $Ax = 0$. Then for any $s \in K$, $K = \{s\} + K_H = \{s + k : k \in K_H\}$.

Proof. Let $s \in K$. Then $K = s + K_H$.

- (1) Let $w \in K$. Then $Aw = b$. So, $A(w - s) = Aw - As = b - b = 0$. Thus, $w - s \in K_H$ and we have $w = s + (w - s)$. So, $w \in \{s\} + K_H$.
- (2) Let $w \in \{s\} + K_H$. Then $w = s + k$ for some $k \in K_H$, and so $Aw = A(s + k) = As + Ak = b + 0 = b$. Thus, $w \in K$.

\square

Theorem 3.10. Let $Ax = b$ be a system of n linear equations in n unknowns. Then A is invertible if and only if the system has exactly one solution. Namely, $x = A^{-1}b$.

Proof. Let $Ax = b$ be a system of n linear equations in n unknowns.

- (\implies): Suppose A is invertible. Then $x = A^{-1}b$ is a solution to $Ax = b$. Clearly, $AA^{-1}b = b$. If s is a solution to $Ax = b$, then $As = b$. So, $A^{-1}As = A^{-1}b$ implies $s = A^{-1}b$.
- (\impliedby): Suppose $Ax = b$ has exactly one solution. Let s be this solution. Let K_H be the set of solutions to $Ax = 0$. Then, by theorem 3.9, $\{s\} = \{s\} + K_H$. Thus, $K_H = \{0\}$. Therefore, $N(L_A) = K_H = \{0\}$. Therefore, L_A is injective and surjective. Thus, L_A has an inverse. So, A has an inverse.

\square

Theorem 3.11. The system $Ax = b$ is consistent if and only if $\text{rank}(A) = \text{rank}(A|b)$.

Proof. Note, $R(L_A) = \text{span}(\{a_1, \dots, a_n\})$ where a_i is the i th column of A .

$$\begin{aligned}
 Ax = b \text{ is consistent} &\iff b \in R(L_A) \\
 &\iff b \in \text{span}(\{a_1, \dots, a_n\}) \\
 &\iff \text{span}(\{a_1, \dots, a_n, b\}) = \text{span}(\{a_1, \dots, a_n\}) \\
 &\iff \dim(\text{span}(\{a_1, \dots, a_n, b\})) = \dim(\text{span}(\{a_1, \dots, a_n\})) \\
 &\iff \text{rank}(A|b) = \text{rank}(A).
 \end{aligned}$$

\square

4 Determinants

4.1 Determinants of order 2

Definition 1 If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then the determinant of A is $ad - bc$.

Notation: The determinant of A will be denoted by $\det(A)$ or $|A|$.

Theorem 4.1. For $u, v, w \in \mathbb{F}^2$ and $k \in \mathbb{F}$

$$\begin{aligned} \det \begin{pmatrix} u + kv \\ w \end{pmatrix} &= \det \begin{pmatrix} u \\ w \end{pmatrix} + k \det \begin{pmatrix} v \\ w \end{pmatrix} \\ \det \begin{pmatrix} w \\ u + kv \end{pmatrix} &= \det \begin{pmatrix} w \\ u \end{pmatrix} + k \det \begin{pmatrix} w \\ v \end{pmatrix}. \end{aligned}$$

Theorem 4.2. Let $A \in M_{2 \times 2}(\mathbb{F})$. Then the determinant of A is non-zero if and only if A is invertible. If A is invertible, then $A^{-1} = \frac{1}{\det(A)} \begin{pmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{pmatrix}$.

Proof. Let $A \in M_{2 \times 2}(\mathbb{F})$.

(\Rightarrow): Let $M = \frac{1}{\det(A)} \begin{pmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{pmatrix}$. Then,

$$\begin{aligned} AM &= \frac{1}{\det(A)} \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{pmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{pmatrix} \\ &= I_2 \\ &= MA. \end{aligned}$$

(\Leftarrow): Suppose A is invertible. Then $\text{rank}(A) = 2$. If $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$, then $A_{11} \neq 0$ or $A_{12} \neq 0$.

(a) Suppose $A_{11} \neq 0$. Then we can transform A into

$$\begin{pmatrix} A_{11} & A_{12} \\ 0 & A_{22} - \frac{A_{21}A_{12}}{A_{11}} \end{pmatrix},$$

which has a rank of 2. Therefore, $A_{22} - \frac{A_{21}A_{12}}{A_{11}} \neq 0$. Thus, $\det(A) \neq 0$.

(b) Suppose $A_{11} = 0$. Then $A_{12} \neq 0$ and $A_{21} \neq 0$. So, $\det(A) = -A_{12} - A_{21} \neq 0$.

□

Observation: Let $\delta : M_{2 \times 2}(\mathbb{F}) \rightarrow \mathbb{F}$ be such that

$$(1) \quad \delta \begin{pmatrix} u + kv \\ w \end{pmatrix} = \delta \begin{pmatrix} u \\ w \end{pmatrix} + k\delta \begin{pmatrix} v \\ w \end{pmatrix}$$

$$\delta \begin{pmatrix} w \\ u + kv \end{pmatrix} = \delta \begin{pmatrix} w \\ u \end{pmatrix} + k\delta \begin{pmatrix} w \\ v \end{pmatrix};$$

$$(2) \quad \delta \begin{pmatrix} u \\ u \end{pmatrix} = 0;$$

$$(3) \quad \delta(I_2) = 1.$$

Then $\delta = \det$.

Proof. We have

$$\delta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \delta \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix} = \delta \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\delta \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \delta \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} = \delta \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$

and

$$\delta \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} + \delta \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} = \delta \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} = 0.$$

Thus,

$$\delta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \delta \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = 0$$

and


$$\delta \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = -1.$$

Then,

$$\begin{aligned} \delta \begin{pmatrix} a & b \\ c & d \end{pmatrix} &= \delta \begin{pmatrix} a & 0 \\ c & d \end{pmatrix} + \delta \begin{pmatrix} 0 & b \\ c & d \end{pmatrix} \\ &= a\delta \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} + b\delta \begin{pmatrix} 0 & 1 \\ c & d \end{pmatrix} \\ &= a \left(c\delta \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} + d\delta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) + b \left(c\delta \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + d\delta \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix} \right) \\ &= ad - bc \\ &= \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}. \end{aligned}$$

□

Observation: When $\mathbb{F} = \mathbb{R}$ and $M_{2 \times 2}(\mathbb{R})$ and for $\begin{pmatrix} u \\ v \end{pmatrix}$, then we have the following diagram.



We will define the area of the parallelogram to be $A \begin{pmatrix} u \\ v \end{pmatrix}$. If u and v are linearly dependent, then $A \begin{pmatrix} u \\ v \end{pmatrix} = 0$. We can show $A \begin{pmatrix} u \\ v \end{pmatrix} = \text{sign} \left(\det \begin{pmatrix} u \\ v \end{pmatrix} \right) \det \begin{pmatrix} u \\ v \end{pmatrix}$.

Proof. Define $O : M_{2 \times 2}(\mathbb{R}) \rightarrow \{-1, 1\}$ by

$$O \begin{pmatrix} u \\ v \end{pmatrix} = \begin{cases} \frac{\det \begin{pmatrix} u \\ v \end{pmatrix}}{\left| \det \begin{pmatrix} u \\ v \end{pmatrix} \right|} & \text{if } u, v \text{ are linearly independent} \\ 1 & \text{if } u, v \text{ are linearly dependent.} \end{cases}$$

Thus, we can show $\det \begin{pmatrix} u \\ v \end{pmatrix} = O \begin{pmatrix} u \\ v \end{pmatrix} A \begin{pmatrix} u \\ v \end{pmatrix}$. We will show $\delta \begin{pmatrix} u \\ v \end{pmatrix} = O \begin{pmatrix} u \\ v \end{pmatrix} A \begin{pmatrix} u \\ v \end{pmatrix}$ satisfies the previous observation. Note,

$$\delta \begin{pmatrix} u \\ u \end{pmatrix} = O \begin{pmatrix} u \\ u \end{pmatrix} A \begin{pmatrix} u \\ u \end{pmatrix} = 0 \quad \text{and} \quad \delta(I_2) = \delta \begin{pmatrix} e_1 \\ e_2 \end{pmatrix} A \begin{pmatrix} e_1 \\ e_2 \end{pmatrix} = 1.$$

We now have three steps.

- (a) If $c = 0$, then $\delta \begin{pmatrix} u \\ 0 \end{pmatrix} = 0$. If $c \neq 0$, then $\delta \begin{pmatrix} u \\ cv \end{pmatrix} = O \begin{pmatrix} u \\ cv \end{pmatrix} A \begin{pmatrix} u \\ cv \end{pmatrix}$ and $A \begin{pmatrix} u \\ cv \end{pmatrix} = |c| A \begin{pmatrix} u \\ v \end{pmatrix}$ and $O \begin{pmatrix} u \\ cv \end{pmatrix} = \frac{\det \begin{pmatrix} u \\ cv \end{pmatrix}}{\left| \det \begin{pmatrix} u \\ cv \end{pmatrix} \right|} = \frac{c}{|c|}$. So, $\delta \begin{pmatrix} u \\ cv \end{pmatrix} = c \delta \begin{pmatrix} u \\ v \end{pmatrix}$.

(b) Note, $\delta \begin{pmatrix} u \\ u+v \end{pmatrix} = \delta \begin{pmatrix} u \\ v \end{pmatrix}$. If $a = 0$, then $\delta \begin{pmatrix} u \\ au+bv \end{pmatrix} = \delta \begin{pmatrix} u \\ bv \end{pmatrix} = b\delta \begin{pmatrix} u \\ v \end{pmatrix}$. If $a \neq 0$, then

$$\begin{aligned} \delta \begin{pmatrix} u \\ au+bv \end{pmatrix} &= a\delta \begin{pmatrix} u \\ u+\frac{b}{a}v \end{pmatrix} \\ &= a\delta \begin{pmatrix} u \\ \frac{b}{a}v \end{pmatrix} \\ &= b\delta \begin{pmatrix} u \\ v \end{pmatrix}. \end{aligned}$$

(c) Assume $u \neq 0$. Let $w \in \mathbb{R}^2$ such that uw is linearly independent. Then $v_1 = a_1u + b_1w$ and $v_2 = a_2u + b_2w$. So,

$$\begin{aligned} \delta \begin{pmatrix} u \\ v_1+v_2 \end{pmatrix} &= \delta \begin{pmatrix} u \\ (a_1+a_2)u + (b_1+b_2)w \end{pmatrix} \\ &= (b_1+b_2) \delta \begin{pmatrix} u \\ w \end{pmatrix}. \end{aligned} \quad (\text{by (b)})$$

Also, by (b), $\delta \begin{pmatrix} u \\ v_1 \end{pmatrix} + \delta \begin{pmatrix} u \\ v_2 \end{pmatrix} = b_1 \delta \begin{pmatrix} u \\ w \end{pmatrix} + b_2 \delta \begin{pmatrix} u \\ w \end{pmatrix}$. Therefore,

$$\delta \begin{pmatrix} u \\ v_1+v_2 \end{pmatrix} = \delta \begin{pmatrix} u \\ v_1 \end{pmatrix} + \delta \begin{pmatrix} u \\ v_2 \end{pmatrix}.$$

□

4.2 Determinants of order n

Let \tilde{A}_{ij} be the matrix obtained from A by deleting the i th row and the j th column.

Definition 2 Let $A \in M_{n \times n}(\mathbb{F})$.

- If $n = 1$, then $\det(A) = A_{11}$.
- If $n \geq 2$, then $\det(A) = \sum_{j=1}^n (-1)^{1+j} A_{1j} \det(\tilde{A}_{1j})$.

The cofactor of the i, j th entry of A ,

$$c_{ij} = (-1)^{i+j} \det(\tilde{A}_{ij}).$$

Example:

$$\begin{aligned}
 \begin{vmatrix} 0 & 1 & 3 \\ -2 & -3 & -5 \\ 4 & -4 & 4 \end{vmatrix} &= 0 \cdot (-1)^2 \cdot \begin{vmatrix} -3 & -5 \\ -4 & 4 \end{vmatrix} + 1 \cdot (-1)^3 \cdot \begin{vmatrix} -2 & -5 \\ 4 & 4 \end{vmatrix} + 3 \cdot (-1)^4 \cdot \begin{vmatrix} -2 & -3 \\ 4 & -4 \end{vmatrix} \\
 &= 0 - 12 + 60 \\
 &= 48.
 \end{aligned}$$

Theorem 4.3. Let $a_1, \dots, a_n \in \mathbb{F}^n$, let $k \in \mathbb{F}$ and suppose $a_r = u + kv$ for some $u, v \in \mathbb{F}^n$. Then

$$\begin{vmatrix} a_1 \\ a_{r-1} \\ a_r \\ a_{r+1} \\ a_n \end{vmatrix} = \begin{vmatrix} a_1 \\ a_{r-1} \\ u \\ a_{r+1} \\ a_n \end{vmatrix} + k \begin{vmatrix} a_1 \\ a_{r-1} \\ v \\ a_{r+1} \\ a_n \end{vmatrix}.$$

Proof. Let $a_r = u + kv$.

(Base Case): Let $n = 1$. Then clearly, $\det(A) = A_{11}$.

(Inductive Step): Let $n \geq 2$.

- If $r = 1$, then $A_{1j} = u_j + kv_j$. Let B and C be matrices obtained from A by replacing row r by u and v . Then

$$\det(\tilde{A}_{1j}) = \det(\tilde{B}_{1j}) = \det(\tilde{C}_{1j}).$$

Thus,

$$\begin{aligned}
 \det(A) &= \sum_{j=1}^n (-1)^{1+j} A_{1j} \det(\tilde{A}_{1j}) \\
 &= \sum_{j=1}^n (-1)^{1+j} (u_j + kv_j) \det(\tilde{A}_{1j}) \\
 &= \sum_{j=1}^n (-1)^{1+j} u_j \det(\tilde{B}_{1j}) + \sum_{j=1}^n (-1)^{1+j} kv_j \det(\tilde{C}_{1j}) \\
 &= \sum_{j=1}^n (-1)^{1+j} B_{1j} \det(\tilde{B}_{1j}) + k \sum_{j=1}^n (-1)^{1+j} C_{1j} \det(\tilde{C}_{1j}) \\
 &= \det(B) + k \det(C).
 \end{aligned}$$

- If $r > 1$, then $\tilde{A}_{1j} = \tilde{B}_{1j} = \tilde{C}_{1j}$ are the same except row $r - 1$, which is $(u_1 + kv_1, \dots, u_{j-1} + kv_{j-1}, u_{j+1} + kv_{j+1}, \dots, u_n + kv_n)$ in A , $(u_1, \dots, u_{j-1}, u_{j+1}, \dots, u_n)$ in B , and $(v_1, \dots, v_{j-1}, v_{j+1}, \dots, v_n)$ in C . Therefore, by the inductive hypothesis

$$\det(\tilde{A}_{1j}) = \det(\tilde{B}_{1j}) + k \cdot \det(\tilde{C}_{1j}).$$

Thus,

$$\begin{aligned} \det(A) &= \sum_{j=1}^n (-1)^{1+j} A_{1j} \det(\tilde{A}_{1j}) \\ &= \sum_{j=1}^n (-1)^{1+j} B_{1j} \det(\tilde{B}_{1j}) + k \cdot \sum_{j=1}^n (-1)^{1+j} C_{1j} \det(\tilde{C}_{1j}) \\ &= \det(B) + k \cdot \det(C). \end{aligned}$$

Therefore,

$$\det(A) = \det(B) + k \cdot \det(C).$$

□

Corollary 4 If A has a row consisting of zeroes, then $\det(A) = 0$.

Lemma 5. Let $B \in M_{n \times n}(\mathbb{F})$ and $n \geq 2$. Suppose that the i th row of B is e_k for some $1 \leq k \leq n$. Then $\det(B) = (-1)^{i+k} \det(\tilde{B}_{ik})$.

Theorem 4.4. For $A \in M_{n \times n}(\mathbb{F})$ and $i \in \{1, \dots, n\}$ and $n \geq 2$

$$\det(A) = \sum_{j=1}^n (-1)^{i+j} A_{ij} \det(\tilde{A}_{ij}).$$

Proof. Let B_j denote the matrix obtained from A by replacing the i th row with e_j . Then

$$A = \sum_{j=1}^n A_{ij} B_j.$$

Thus,

$$\begin{aligned} \det(A) &= \sum_{j=1}^n A_{ij} \det(B_j) && \text{(By Theorem 4.3)} \\ &= \sum_{j=1}^n A_{ij} (-1)^{i+j} \det(\tilde{A}_{ij}). && \text{(By Lemma 5)} \end{aligned}$$

□

Corollary 7 If $A \in M_{n \times n}(\mathbb{F})$ has two identical rows, then $\det(A) = 0$.

Proof. Let $A \in M_{n \times n}(\mathbb{F})$.

(Base Case): Let $n = 2$. Then $\begin{pmatrix} a & b \\ a & b \end{pmatrix} = ab - ab = 0$.

(Inductive Step): Let $n \geq 3$. Suppose rows r and s are identical. Let $i \in \{1, \dots, n\}$ such that $i \neq r$ and $i \neq s$. By Theorem 4.4,

$$\det(A) = \sum_{j=1}^n A_{ij}(-1)^{i+j} \det(\tilde{A}_{ij})$$

and \tilde{A}_{ij} has two identical rows. So, by the inductive hypothesis, $\det(\tilde{A}_{ij}) = 0$. Therefore, $\det(A) = 0$.

□

Theorem 4.5. If $A \in M_{n \times n}(\mathbb{F})$ and B is obtained from A by interchanging two rows, then $\det(B) = -\det(A)$.

Proof. Let $A \in M_{n \times n}(\mathbb{F})$. Suppose B is obtained from A by interchanging rows r and s

with $r < s$, without loss of generality. Let a_i denote the i th row of A . Then

$$\begin{aligned}
 0 &= \det \begin{pmatrix} a_1 \\ \vdots \\ a_r + a_s \\ \vdots \\ a_r + a_s \\ \vdots \\ a_n \end{pmatrix} \\
 &= \det \begin{pmatrix} a_1 \\ \vdots \\ a_r \\ \vdots \\ a_r + a_s \\ \vdots \\ a_n \end{pmatrix} + \det \begin{pmatrix} a_1 \\ \vdots \\ a_s \\ \vdots \\ a_r + a_s \\ \vdots \\ a_n \end{pmatrix} \\
 &= \det \begin{pmatrix} a_1 \\ \vdots \\ a_r \\ \vdots \\ a_r \\ \vdots \\ a_n \end{pmatrix} + \det \begin{pmatrix} a_1 \\ \vdots \\ a_r \\ \vdots \\ a_s \\ \vdots \\ a_n \end{pmatrix} + \det \begin{pmatrix} a_1 \\ \vdots \\ a_s \\ \vdots \\ a_r \\ \vdots \\ a_n \end{pmatrix} + \det \begin{pmatrix} a_1 \\ \vdots \\ a_s \\ \vdots \\ a_s \\ \vdots \\ a_n \end{pmatrix} \\
 &= 0 + \det(A) + \det(B) + 0.
 \end{aligned}$$

Therefore, $\det(B) = -\det(A)$. □

Theorem 4.6. If $A \in M_{n \times n}(\mathbb{F})$ and B is obtained from A by adding a multiple of one row to another, then $\det(B) = \det(A)$.

Proof. Let $A \in M_{n \times n}(\mathbb{F})$. Let B be obtained from A by adding ka_r to a_s . Let C be

obtained from A by replacing row s with a_r . Then $C = \begin{pmatrix} a_1 \\ \vdots \\ a_r \\ \vdots \\ a_r \\ \vdots \\ a_n \end{pmatrix}$ and so, $\det(C) = 0$. In

addition, row s of B is equal to row s of A plus k times row s of C . Thus,

$$\begin{aligned} \det(B) &= \det(A) + k\det(C) \\ &= \det(A). \end{aligned}$$

□

Corollary 10 If $A \in M_{n \times n}(\mathbb{F})$ and $\text{rank}(A) < n$, then $\det(A) = 0$.

Proof. Let $A \in M_{n \times n}(\mathbb{F})$ and $\text{rank}(A) < n$. Then the dimension of the row space of A is less than n . Thus, the rows of A are linearly dependent. Therefore, for some $r \in \{1, \dots, n\}$, $a_r = \sum_{j \neq r} c_j a_j$ for some $c_j \in \mathbb{F}$. For every $j \neq r$, if $c_j \neq 0$, then adding $-c_j a_j$ to row r results in the zero vector. Therefore, $\det(A) = 0$. □

Example:

$$\begin{aligned} \begin{vmatrix} 0 & 2 & 1 & 5 \\ 1 & 5 & 0 & 6 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & -2 \end{vmatrix} &\xrightarrow{\text{row op.}} - \begin{vmatrix} 1 & 5 & 0 & 6 \\ 0 & 2 & 1 & 5 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & -2 \end{vmatrix} \\ &= -1 \cdot 2 \cdot 4 \cdot (-2) \\ &= 16. \end{aligned}$$

4.3 Properties

Summary of properties

- Let $A, B \in M_{n \times n}(\mathbb{F})$. Then $\det(AB) = \det(A)\det(B)$.

Proof. Let $A, B \in M_{n \times n}(\mathbb{F})$.

(1) We will show this is the case when A is an elementary matrix.

- * If A is of Type I (interchange), then $\det(A) = -\det(I) = -1$ and by Theorem 4.5, $\det(AB) = -\det(B)$.
- * If A is Type II (multiply by k), then $\det(A) = k$ and $\det(AB) = k\det(B)$.
- * If A is of type III (adding), then $\det(A) = 1$ and $\det(AB) = \det(B)$.

So, the fact holds when A is an elementary matrix.

(2) If $\text{rank}(A) < n$, then by Corollary 10, $\det(A) = 0$. In addition, $\text{rank}(AB) \leq \text{rank}(A) < n$. Thus, $\det(A) = 0$.

- (3) If $\text{rank}(A) = n$, then A is invertible and a product of elementary matrices. Say $A = E_1 E_2 \dots E_m$ where E_i are elementary matrices. Then

$$\begin{aligned}
 \det(AB) &= \det(E_1 \dots E_m \cdot B) \\
 &= \det(E_1) \det(E_2 \dots E_m \cdot B) \\
 &\vdots \\
 &= \det(E_1) \det(E_2) \dots \det(E_m) \det(B) \\
 &= \det(E_1 \dots E_m) \det(B) \\
 &= \det(A) \det(B).
 \end{aligned}$$

□

- $A \in M_{n \times n}(\mathbb{F})$ is invertible if and only if $\det(A) \neq 0$ and $\det(A^{-1}) = \frac{1}{\det(A)}$.

Proof.

\implies If A is not invertible, then $\text{rank}(A) < n$ and so $\det(A) = 0$.

\impliedby If A is invertible, then $A \cdot A^{-1} = I_n$. Thus, by the fact above, $1 = \det(I_n) = \det(A) \det(A^{-1})$. So, $\det(A) \neq 0$.

□

- Let $A \in M_{n \times n}(\mathbb{F})$. Then $\det(A^t) = \det(A)$.

Proof.

(1) If A is not invertible, then $\text{rank}(A) < n$ and $\det(A) = 0$. Also, $\text{rank}(A^t) = \text{rank}(A) < n$. Thus, $\det(A^t) = 0$.

(2) If A is invertible, then A is a product of elementary matrices, say $A = E_1 E_2 \dots E_m$. Then $A^t = (E_1 \dots E_m)^t = E_m^t \dots E_1^t$. Thus,

$$\begin{aligned}
 \det(A^t) &= \det(E_m^t) \dots \det(E_1^t) \\
 &= \det(E_m) \dots \det(E_1) \\
 &= \det(E_1) \dots \det(E_m) \\
 &= \det(A).
 \end{aligned}$$

□

- (Cramer's Rule) Let $Ax = b$ where $A \in M_{n \times n}(\mathbb{F})$ and let M_k be the $n \times n$ matrix obtained from A by replacing column k with b . Then $x_k = \frac{\det(M_k)}{\det(A)}$.

Proof. Let $A \in M_{n \times n}(\mathbb{F})$ be nonzero. Thus, A is invertible, and so, $Ax = b$ has a unique solution. Fix k . Let X be obtained from I_n by replacing the k th column with

x . Then the i th column of AX is $Ae_i = a_i$ if $i \neq k$ and $Ax = b$ if $i = k$. Therefore $AX = M_k$ and so

$$\det(A)\det(X) = \det(AX) \\ \det(M_k).$$

So, $\det(X) = x_k$. Therefore, $x_k = \frac{\det(M_k)}{\det(A)}$. □

5 Diagonalization

5.1 Eigenvalues and eigenvectors

Definition 1 Let V be finite-dimensional and let $T : V \rightarrow V$ be linear. T is called *diagonalizable* if there is an ordered basis β such that $[T]_\beta$ is a diagonal matrix.

Matrix A is diagonalizable if L_A is.

Definition 2 A non-zero vector $v \in V$ such that $T(v) = \lambda v$ for some $\lambda \in \mathbb{F}$ is called an *eigenvector*. The scalar λ is called an *eigenvalue* of T .

Theorem 5.1. Let V be finite-dimensional. A linear operator on V is diagonalizable if and only if there is an ordered basis β for V consisting of eigenvectors of T .

Proof.

\Rightarrow Suppose T is diagonalizable. Then there exists an ordered basis $\beta = \{v_1, \dots, v_n\}$ for V such that $[T]_\beta = D$ where $D = [D_{ij}]$ and $D_{ij} = 0$ if $i \neq j$. Then

$$T(v_j) = \sum_{i=1}^n D_{ij}v_i = D_{jj}v_j.$$

Thus, v_j is an eigenvector of T since $v_j \neq 0$.

\Leftarrow Suppose $\beta = \{v_1, \dots, v_n\}$ is a basis for V consisting of eigenvectors of T , say $T(v_j) = \lambda_j v_j$. Then $[T]_\beta = D = [D_{ij}]$ where

$$D_{ij} = \begin{cases} \lambda_j & i = j \\ 0 & i \neq j. \end{cases}$$

Thus, D is diagonal. □

Theorem 5.2. Let $A \in M_{n \times n}(\mathbb{F})$. Then $\lambda \in \mathbb{F}$ is an eigenvalue of A if and only if $\det(A - \lambda I_n) = 0$.

Proof.

$$\begin{aligned}
 \lambda \text{ is an eigenvalue of } A &\iff \exists v \neq 0 \text{ such that } Av - \lambda v = 0 \\
 &\iff \exists v \neq 0 \text{ such that } (A - \lambda I_n)v = 0 \\
 &\iff \exists v \neq 0 \text{ such that } v \in N(A - \lambda I_n) \\
 &\iff A - \lambda I_n \text{ is not invertible} \\
 &\iff \det(A - \lambda I_n) = 0
 \end{aligned}$$

□

Definition 3

- Let $A \in M_{n \times n}(\mathbb{F})$. Then $f(t) = \det(A - tI_n)$ is called the *characteristic polynomial* of A .
- If T is a linear operator V , then the characteristic polynomial of T is the characteristic polynomial of $A = [T]_\beta$ where β is an ordered basis for V .

Note: The characteristic polynomial of T is well-defined.

Proof. Let β and α be ordered bases for V . Then $f_T(t) = \det([T]_\beta - tI) = \det([T]_\alpha - tI)$. Let $B = [T]_\beta$ and $A = [T]_\alpha$. Then there exists Q invertible such that $B = Q^{-1}AQ$. So,

$$\begin{aligned}
 \det(B - tI) &= \det(Q^{-1}AQ - tI) \\
 &= \det(Q^{-1}(A - tI)Q) \\
 &= \det(Q^{-1})\det(A - tI)\det(Q) \\
 &= \det(A - tI).
 \end{aligned}$$

□

Theorem 5.3. Let $A \in M_{n \times n}(\mathbb{F})$.

- $f_A(t)$ is a polynomial of degree n with the leading coefficient $(-1)^n$.
- A has at most n distinct eigenvalues.

Theorem 5.4. Let T be a linear operator on V and let λ be an eigenvalue of T . A vector $v \in V$ is an eigenvector of T corresponding to λ if and only if $v \neq 0$ and $v \in N(T - \lambda I)$.

Example: Let $C^\infty(\mathbb{R}) = \{f : \mathbb{R} \rightarrow \mathbb{R} : f^{(n)} \text{ exists for every } n \in \mathbb{Z}^+\}$. Then $C^\infty(\mathbb{R})$ is a subspace of $\mathcal{F}(\mathbb{R}, \mathbb{R})$. Let $T : C^\infty(\mathbb{R}) \rightarrow C^\infty(\mathbb{R})$, $T(f) = f'$. Find all eigen values of T .

Solution. We know f is an eigenvector if $f \neq 0$ and $Tf = \lambda f$ for some $\lambda \in \mathbb{R}$. So, $f' = \lambda f$ implies $\frac{df}{dx} = \lambda f$ implies $\frac{df}{f} = \lambda dx$ implies $\ln(f) = \lambda x + C$ implies $|f| = e^{\lambda x + C}$ implies $f(x) = Le^{\lambda x}$ for some $L \in \mathbb{R}$ and $L \neq 0$. All $\lambda \in \mathbb{R}$ are eigenvalues.

Example: Let $T : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ be defined by $T(A) = A^T$. Is T diagonalizable?

Solution. Let $\alpha = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\} = \{e_1, e_2, e_3, e_4\}$. So,

$$[T]_\alpha = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

is not diagonalizable. We need to find the eigenvalues of T . If λ is an eigenvalue, then $T(A) = \lambda A$ for some $A \neq 0$. Then, $T^2(A) = T(T(A)) = \lambda^2 A$. Thus, $\lambda^2 A = A \neq 0$. So, $\lambda = \pm 1$.

($\lambda = 1$): Then $T(A) = A$ and $T(A) = A^T$. Thus, $A = A^T$ is a symmetric matrix. Let $\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\} = \{f_1, f_2, f_3\}$.

($\lambda = -1$): Then $T(A) = -A$ and $T(A) = A^T$. So, $A^T = -A$ is a skew-symmetric matrix. Let $\left\{ \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\} = \{f_4\}$.

Then $\beta = \{f_1, f_2, f_3, f_4\}$ is a basis of eigenvectors of T . Then $T(f_1) = f_1$, $T(f_2) = f_2$, $T(f_3) = f_3$, and $T(f_4) = -f_4$ implies

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}.$$

Example: Let $A \in M_{2 \times 2}(\mathbb{F})$ be defined as $\begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$. Find the eigenvectors of A .

Solution.

- Finding the eigenvalues of A , we have

$$\begin{aligned} f_A(t) &= \begin{vmatrix} 1-t & 1 \\ 4 & 1-t \end{vmatrix} \\ &= (1-t)^2 - 4 \\ &= (t-3)(t+1). \end{aligned}$$

So, the eigenvalues of A are $\lambda_1 = 3$ and $\lambda_2 = -1$.

- Finding the eigenvectors of A , we have,

λ_1 : We have $A - 3I = \begin{pmatrix} -2 & 1 \\ 4 & -2 \end{pmatrix}$. So, $(A - 3I) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0$ implies $\begin{pmatrix} -2 & 1 \\ 4 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0$ implies $x_1 = t$ and $x_2 = 2t$. Thus, $N(A - \lambda_1 I) = \left\{ t \begin{pmatrix} 1 \\ 2 \end{pmatrix} : t \in \mathbb{R} \right\}$.

λ_2 : We have $A + I = \begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix}$. So, $(A + I) \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0$ implies $\begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 0$ implies $x_1 = t$ and $x_2 = -2t$. Thus, $N(A - \lambda_1 I) = \left\{ t \begin{pmatrix} 1 \\ -2 \end{pmatrix} : t \in \mathbb{R} \right\}$.

Now, is A diagonalizable? Yes. For $\beta = \left\{ \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ -2 \end{pmatrix} \right\}$, we have $[L_A]_\beta = \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix}$.

5.2 Diagonalizability

Theorem 5.5. Let T be a linear operator with distinct eigenvalues $\lambda_1, \dots, \lambda_k$. If v_1, \dots, v_k are eigenvectors of T such that v_i corresponds to λ_i , then $\{v_1, \dots, v_k\}$ is linearly independent.

Proof. Let $\lambda_1, \dots, \lambda_k$ be distinct eigenvalues.

Base Step: If $k = 1$, then $v_1 \neq 0$ implies v_1 is linearly independent.

Induction Step: Suppose v_1, \dots, v_{k-1} is linearly independent. Suppose $\sum_{i=1}^k c_i v_i = 0$ for some $c_i \in \mathbb{F}$. Then

$$(T - \lambda_k I) \left(\sum_{i=1}^k c_i v_i \right) = (T - \lambda_k I)(0) = 0.$$

Thus,

$$\begin{aligned} 0 &= \sum_{i=1}^k c_i (T - \lambda_k I)(v_i) \\ &= \sum_{i=1}^k c_i (\lambda_i - \lambda_k)(v_i) && \text{(Since } T(v_i) = \lambda_i v_i \text{)} \\ &= \sum_{i=1}^{k-1} c_i (T - \lambda_k I)(v_i). \end{aligned}$$

Since v_1, \dots, v_{k-1} is linearly independent, then $c_i (\lambda_i - \lambda_k) = 0$. Thus, $c_i = 0$ for $i = 1, \dots, k-1$ since $\lambda_i \neq \lambda_k$ if $i < k$. Thus, $c_k v_k = 0$ and $v_k \neq 0$ implies $c_k = 0$ and v_1, \dots, v_k is linearly independent. \square

Corollary 6 If T has n distinct eigenvalues, then T is diagonalizable.

Definition 4 A polynomial $f(t) \in P(\mathbb{F})$ splits over \mathbb{F} if $f(t) = c(t - a_1) \dots (t - a_n)$ for some $c, a_1, \dots, a_n \in \mathbb{F}$.

Theorem 5.6. Let T be a linear operator. If T is diagonalizable, then its characteristic polynomial splits.

Proof. Suppose T is diagonalizable. Then there exists $\beta = \{v_1, \dots, v_n\}$ such that v_i is an eigenvector of T , say $T(v_j) = \lambda_j v_j$. Consequently, $A = [T]_\beta = I\lambda_i$ and so

$$\begin{aligned} f_T(t) &= f_A(t) \\ &= (\lambda_1 - t) \dots (\lambda_n - t) \\ &= (-1)^{n+1} (t - \lambda_1) \dots (t - \lambda_n). \end{aligned}$$

Therefore, $f_T(t)$ splits. □

Lemma 8. Let $X \in M_{n \times n}(\mathbb{F})$ such that $X = \begin{pmatrix} A & B \\ 0 & C \end{pmatrix}$ for $A \in M_{m \times m}(\mathbb{F})$, $B \in M_{m \times p}(\mathbb{F})$, and $C \in M_{p \times p}(\mathbb{F})$. Then $\det(X) = \det(A)\det(C)$.

Proof.

- Suppose $C = I_p$. We will use induction on p . The base case is clearly true. Suppose $p \geq 2$. Use the expansion of the last row. Then $\det(X) = (-1)^{n+n} \cdot 1 \cdot \det \left(\begin{pmatrix} A & B' \\ 0 & I_{p-1} \end{pmatrix} \right)$ where B' is obtained from B by deleting the last column. Then, by the inductive hypothesis, $\det(X) = \det(A)\det(I_{p-1}) = \det(A)$. Thus, $\det(X) = \det(A)\det(C)$.
- Suppose $A = I_m$, then use the first column. Thus, $\det(X) = \det(A)\det(C)$.
- For the general case,
 - If $\det(A) = 0$, then the columns of A are linearly dependent. Then the first m columns of X are linearly dependent. Thus, $\det(X) = 0 = \det(A)\det(C)$.
 - If $\det(A) \neq 0$, then A is invertible. Then,

$$X = \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & I_p \end{pmatrix} \begin{pmatrix} I_m & A^{-1}B \\ 0 & C \end{pmatrix}.$$

Therefore, $\det(X) = \det(A)\det(C)$. □

Definition 5 The (algebraic) *multiplicity* of an eigenvalue λ is the largest positive integer k such that $(t - \lambda)^k | f(t)$.

The eigenspace of T with respect to λ is

$$E_\lambda = N(T - \lambda I_V) = \{x \in V | T(x) = \lambda x\}.$$

Theorem 5.7. Let λ be an eigenvalue of T of multiplicity m . Then

$$1 \leq \dim(E_\lambda) \leq m.$$

Proof. (Outline)

- Start with an ordered basis $\{v_1, \dots, v_p\}$ for E_λ and extend it to an ordered basis β for V .
- Then $A = [T]_\beta$ will have the following form

$$\begin{pmatrix} \lambda I_p & B \\ 0 & C \end{pmatrix}.$$

Proof. Let $k = \dim(E_\lambda)$ and let $\{v_1, \dots, v_k\}$ be a basis for E_λ . Extend the basis to a basis $\beta = \{v_1, \dots, v_k, v_{k+1}, \dots, v_n\}$ for V . Then $T(v_j) = \lambda v_j$ for $j = 1, \dots, k$. So,

$$A = [T]_\beta = \begin{pmatrix} \lambda I_k & B \\ 0 & C \end{pmatrix}.$$

Therefore,

$$\begin{aligned} f_T(t) &= f_A(t) \\ &= \begin{vmatrix} (\lambda - t)I_k & B \\ 0 & C - tI_{n-k} \end{vmatrix} \\ &= \det((\lambda - t)I_k) \det(C - tI_{n-k}) && \text{(By Lemma 8)} \\ &= (-1)^k (t - \lambda)^k g(t). \end{aligned}$$

Thus, $(t - \lambda)^k | f_T(t)$ and $k \leq m$. □

Example: Let $T : P_3(\mathbb{R}) \rightarrow P_3(\mathbb{R})$ be defined as $T(f) = f'$. Is T diagonalizable?

Solution. Let $\beta = \{1, x, x^2, x^3\}$ be a basis for $P_3(\mathbb{R})$. Then

$$[T]_\beta = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Thus,

$$f_T(t) = \begin{vmatrix} -t & 1 & 0 & 0 \\ 0 & -t & 2 & 0 \\ 0 & 0 & -t & 3 \\ 0 & 0 & 0 & -t \end{vmatrix} = t^4.$$

So, there is only one eigenvalue, $\lambda = 0$. Thus,

$$\begin{aligned} E_0 &= N(T - 0I) \\ &= \{f : Tf = 0\} \\ &= \{f : f' = 0\} \\ &= P_0(\mathbb{R}). \end{aligned}$$

Thus, $\dim(E_0) = 1$. Therefore, it is not possible to find a basis consisting of eigenvectors.

Lemma 9. Let $\lambda_1, \dots, \lambda_k$ be distinct eigenvalues of T and let $v_i \in E_{\lambda_i}$. If $v_1 + \dots + v_k = 0$, then $v_i = 0$ for every i .

Proof. Renumerate v_1, \dots, v_k so that for some m , $v_i = 0$ for $i > m$ and $v_i \neq 0$ for $i \leq m$. Then $v_1 + \dots + v_m = 0$ but, v_1, \dots, v_m are linearly independent. Therefore, $m = 0$. \square

Theorem 5.8. Let $\lambda_1, \dots, \lambda_k$ be distinct eigenvalues of T and let $S_i \subseteq E_{\lambda_i}$ be finite and linearly independent. Then $S_1 \cup \dots \cup S_k$ is linearly independent.

Proof. (Outline) Say $S_i = \{v_{i1}, \dots, v_{in_i}\}$ and suppose $\sum_{i=1}^k \sum_{j=1}^{n_i} a_{ij}v_{ij} = 0$. Consider $w_i = \sum_{j=1}^{n_i} a_{ij}v_{ij}$.

Proof. Suppose $S_i = \{v_{i1}, v_{i2}, \dots, v_{in_i}\}$ and $|S_i| = n_i$. Suppose $\sum_{i=1}^k \sum_{j=1}^{n_i} a_{ij}v_{ij} = 0$. Let $w_i = \sum_{j=1}^{n_i} a_{ij}v_{ij}$. Then $w_i \in E_{\lambda_i}$ and we have $w_1 + \dots + w_k = 0$. By Lemma 9, $w_i = 0$ for every $i = 1, \dots, k$. Thus, $\sum_{j=1}^{n_i} a_{ij}v_{ij} = 0$. Since S_i is linearly independent, $a_{ij} = 0$ for all i and j . \square

Theorem 5.9. Let T be a linear operator on a finite-dimensional vector space V such that the characteristic polynomial of T splits. Let $\lambda_1, \dots, \lambda_k$ be distinct eigenvalues of T . Then

- (a) T is diagonalizable if and only if the multiplicity of each λ_i equals $\dim(E_{\lambda_i})$.
- (b) If T is diagonalizable and β_i is an ordered basis for E_{λ_i} , then $\beta = \beta_1 \cup \dots \cup \beta_k$ is an ordered basis for V .

Proof. (Outline)

- Part (b) follows from the proof.

- (\implies): m_i - the multiplicity of λ_i , $d_i = \dim(E_{\lambda_i})$, $\beta_i = \beta \cap E_{\lambda_i}$, $n_i = |\beta_i|$. We have $n_i \leq d_i \leq m_i$ and $\sum n_i = n = \sum m_i$. It follows that $d_i = m_i$ for every i .
- (\impliedby): Let β_i be an ordered basis for E_{λ_i} ; let $\beta = \beta_1 \cup \dots \cup \beta_k$. Then β is a basis for V consisting of eigenvectors.

Proof. Let $\lambda_1, \dots, \lambda_k$ be distinct eigenvalues. Let m_i denote the multiplicity of λ_i . Let $n = \dim(V)$. Let $d_i = \dim(E_{\lambda_i})$. Then $d_i \leq m_i$ by Theorem 5.7. Note, $m_1 + \dots + m_k = n$.

\implies Suppose T is diagonalizable. Then there exists a basis β consisting of eigenvectors. Let $\beta_i = \beta \cap E_{\lambda_i}$. Then β_i is linearly independent and so $|\beta_i| \leq d_i$. In addition,

$$\begin{aligned}
 n &= |\beta| \\
 &= \sum_{i=1}^k |\beta_i| \\
 &\leq \sum_{i=1}^k d_i \\
 &\leq \sum_{i=1}^k m_i. \qquad \qquad \qquad (\text{By Theorem 5.7})
 \end{aligned}$$

Thus, $m_i = d_i$ for all i .

\impliedby Suppose $m_i = d_i$ for all i . Let β_i be a basis for E_{λ_i} . Thus, $|\beta_i| = d_i$. Let $\beta = \cup_{i=1}^k \beta_i$. Thus, β is linearly independent. In addition,

$$\begin{aligned}
 |\beta| &= \sum_{i=1}^k |\beta_i| \\
 &= \sum_{i=1}^k d_i \\
 &= \sum_{i=1}^k m_i \\
 &= n \\
 &= \dim(V).
 \end{aligned}$$

Thus, β is a basis for V .

□

Note: T is diagonalizable if and only if

- The characteristic polynomial splits and
- the multiplicity of λ_i is $\text{nullity}(T - \lambda_i I) = n - \text{rank}(T - \lambda_i I)$.

Example: Test if $A = \begin{pmatrix} 3 & 1 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix}$ is diagonalizable over \mathbb{R} .

Solution.

- We know $f_A(t) = \begin{vmatrix} 3-t & 1 & 0 \\ 0 & 3-t & 0 \\ 0 & 0 & 4-t \end{vmatrix} = (3-t)^2(4-t)$ splits.
- For $\lambda_1 = 4$, we have $\text{rank}(A - \lambda_1 I) = \text{rank} \left(\begin{pmatrix} -1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right) = 2$ and multiplicity of $\lambda_1 = 1$ and $3 - \text{rank}(A - \lambda_1 I) = 3 - 2 = 1$.
- For $\lambda_2 = 3$, we have $\text{rank}(A - \lambda_2 I) = \text{rank} \left(\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) = 2$ and multiplicity of $\lambda_2 = 2$ and $3 - 2 = 1$.

So, A is not diagonalizable.

5.4 Invariant subspaces and the Cayley-Hamilton theorem

Definition 6

- Let $T : V \rightarrow V$. A subspace $W \subseteq V$ is called a T -invariant subspace of V if $T(W) \subseteq W$.

Examples:

- $\{0\}$
- $N(T)$: if $x \in N(T)$, then $T(x) = 0$.
- $R(T)$: if $y \in R(T)$, then $y \in V$ and so, $T(y) \in R(T)$.
- E_λ : Let $v \in E_\lambda$. Then $T(v) = \lambda v$. Thus, $T(T(v)) = T(\lambda v) = \lambda T(v)$. Thus $T(v) \in E_\lambda$.
- The T -cyclic subspace of V generated by x is

$$\text{span}(\{x, T(x), T^2(x), \dots\}).$$

Note: T -cyclic subspace is a minimal subspaces which is T -invariant and contains x .

Note: If W is T -invariant and $v \in W$, then $T^i(v) \in W$ for all $i \in \mathbb{Z}^*$ and W contains a T -cyclic subspace generated by x .

Notation: If W is T -invariant, then $T_W = T|_W : W \rightarrow W$.

Theorem 5.21. Let T be a linear operator on a finite-dimensional vector space V and let W be a T -invariant subspace of V . Then the characteristic polynomial of the restriction of T to W , T_W , divides the characteristic polynomial of T .

Proof. Let $\alpha = \{v_1, \dots, v_k\}$ be a basis for W and extend it to a basis $\beta = \{v_1, \dots, v_n\}$ for V . Then

$$[T]_{\beta} = \begin{pmatrix} B_1 & B_2 \\ 0 & B_3 \end{pmatrix}$$

for some $B_1 \in M_{k \times k}(\mathbb{F})$ and $[T_W] = B_1$. Therefore, $f_T(t) = f_{T_W}(t)g(t)$ and so, $f_{T_W}(t) | f_T(t)$. \square

Example: Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be defined as $T(a, b, c, d) = (a + b + 5c - d, b + d, c - d, c + d)$. Let $W = \{x, y, 0, 0 : x, y \in \mathbb{R}\}$. Thus, $T(W) \subseteq W$. So, W is T -invariant. Let $\beta = \{e_1, e_2\}$ be a basis for W . Then $T(e_1) = e_1$ and $T(e_2) = (1, 1, 0, 0) = e_1 + e_2$. Thus,

$$[T_W]_{\beta} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Let $\alpha = \{e_1, e_2, e_3, e_4\}$ be a basis for V . So, $T(e_3) = (5, 0, 1, 1)$ and $T(e_4) = (-1, 1, -1, 1)$. Thus,

$$[T]_{\alpha} = \left(\begin{array}{cc|cc} 1 & 1 & 5 & -1 \\ 0 & 1 & 0 & 1 \\ \hline 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{array} \right).$$

Therefore,

$$\begin{aligned} f_T(t) &= \left| \begin{array}{cc|cc} 1-t & 1 & 5 & -1 \\ 0 & 1-t & 0 & 1 \\ \hline 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{array} \right| \\ &= (1-t)^2((1-t)^2 + 1) \\ &= f_{T_W}(t)g(t). \end{aligned}$$

So, W is T -invariant. Now, is W T -cyclic? We know $T(e_1) = e_1$ and $T(e_2) = e_1 + e_2$. Thus, $e_1 \in \text{span}(\{e_2, T(e_2)\})$. Therefore, W is T -cyclic.

Example: Is it possible to have T and W such that W is T -invariant, but is not T -cyclic?

Solution. Yes. Consider $N(T)$ such that $\dim(N(T)) > 1$. Then it is easy to show that W is T -invariant, but not T -cyclic.

Theorem 5.22. Let T be a linear operator on a vector space V of dimension n and let W be a T -cyclic subspace of V generated by a non-zero vector v of dimension k . Then

- (a) $\{v, T(v), \dots, T^{k-1}(v)\}$ is a basis for W .
- (b) If $a_0v + a_1T(v) + \dots + a_{k-1}T^{k-1}(v) + T^k(v) = 0$, then the characteristic polynomial of T_W is $f(t) = (-1)^k(a_0 + a_1t + \dots + a_{k-1}t^{k-1} + t^k)$.

Proof. (Outline)

- (a) Let j be the largest integer such that $\beta = \{v, T(v), \dots, T^{j-1}(v)\}$ is linearly independent. Then $\text{span}(\beta)$ is T -invariant and it follows that $j = k$.
- (b) Let $\beta = v, T(v), \dots, T^{k-1}(v)$ and suppose $a_0v + a_1T(v) + \dots + a_{k-1}T^{k-1}(v) + T^k(v) = 0$. Use the form of $[T_W]_\beta$ to notice that $f_{T_W}(t) = (-1)^k(a_0 + a_1t + \dots + a_{k-1}t^{k-1} + t^k)$.

Proof. Let W be T -cyclic generated by v and $\dim(W) = k$. Since $v \neq 0$, $\{v\}$ is linearly independent. Let j be the largest positive integer such that $\{v, T(v), \dots, T^{j-1}(v)\}$ is linearly independent. Then $j \leq k$. We will show that $j = k$. Let $Z = \text{span}(\{v, T(v), \dots, T^{j-1}(v)\})$. We claim that Z is T -invariant. Let $w \in Z$. Then, $w = a_0v + a_1T(v) + \dots + a_{j-1}T^{j-1}(v)$. So, $T(w) = a_0T(v) + \dots + a_{j-1}T^j(v)$. Thus, $T^j(v)$ is linearly dependent on $\{v, T(v), \dots, T^{j-1}(v)\}$. Thus, $T^j(v) \in Z$ and $T(v), \dots, T^{j-1}(v) \in Z$ and so, $T(w) \in Z$. Therefore, Z is T -invariant. Since Z is T -invariant and $v \in Z$, then $W \subseteq Z$. So, $W = Z$ and $j = k$. Thus, $\{v, T(v), \dots, T^{k-1}(v)\}$ is a basis for W . Then

$$[T_W]_\beta = \begin{pmatrix} 0 & 0 & \dots & \dots & 0 & -a_0 \\ 1 & 0 & \ddots & \ddots & \vdots & -a_1 \\ 0 & 1 & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & 1 & 0 & \vdots \\ 0 & 0 & \dots & 0 & 1 & -a_{k-1} \end{pmatrix}.$$

Thus, $f_{T_W}(t) = (-1)^k(a_0 + a_1t + \dots + a_{k-1}t^{k-1} + t^k)$. □

Theorem 5.23, Cayley-Hamilton. Let V be finite-dimensional and let $T : V \rightarrow V$ be linear with the characteristic polynomial $f(t)$. Then $f(T) = T_0$.

Proof. (Outline) Note that $f(T)$ is a linear operator. We will show $f(T)(v) = 0$ for every v .

- Take $v \neq 0$ and consider the T -cycle subspace generated by v , W .

- Let $k = \dim(W)$. Then $T^k \in \text{span}(\{v, T(v), \dots, T^{k-1}(v)\})$ by 5.22(a) and we are done by 5.22(b).

Proof. We will show $f(T)v = 0_V$ for all $v \in V$. Note, $f(T)$ is a linear operator. Let $v = 0$. Then $f(T)v = 0$. Let $v \neq 0$. Consider the T -cyclic subspace generated by v , say W . Suppose that $\dim(W) = k$. Then by Theorem 5.22, $\{v, T(v), \dots, T^{k-1}(v)\}$ is a basis for W . So, $T^k(v) \in \text{span}(\{v, \dots, T^{k-1}(v)\})$. So, $a_0v + a_1T(v) + \dots + a_{k-1}T^{k-1}(v) + T^k(v) = 0$. Again, by Theorem 5.22, the characteristic polynomial of T_W satisfies $f_{T_W}(t) = (-1)^k(a_0 + \dots + a_{k-1}t^{k-1} + t^k)$. Therefore, $f_{T_W}(T) = (-1)^k(a_0I + \dots + a_{k-1}T^{k-1} + T^k)$. Thus, $f_{T_W}v = (-1)^k(a_0v + \dots + a_{k-1}T^{k-1}(v) + T^k(v)) = 0$. By Theorem 5.21, $F_T(t) = g(t)f_{T_W}(t)$. Thus, $f_T(T) = g(T)f_{T_W}(T)$. Therefore,

$$\begin{aligned} f_T(T)v &= g(T)f_{T_W}(T)v \\ &= g(T)(0_V) \\ &= 0_V. \end{aligned}$$

□

Corollary 15 Let A be an $n \times n$ matrix and let $f(t)$ be its characteristic polynomial. Then $f(A) = 0_n$, the $n \times n$ zero matrix.

Example: Let $A = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$. Then $f_A(t) = \begin{vmatrix} 2-t & 1 \\ 0 & 1-t \end{vmatrix} = (t-1)(t-2) = t^2 - 3t + 2$. Thus, $f(A) = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}^2 - 3 \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix} + 2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

6 Inner product spaces

6.1 Inner products and norms

Definition 1 Let V be a vector space over \mathbb{F} . An *inner product* on V is a function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ such that the following conditions hold.

- $\langle x + z, y \rangle = \langle x, y \rangle + \langle z, y \rangle$
- $\langle cx, y \rangle = c\langle x, y \rangle$
- $\overline{\langle x, y \rangle} = \langle y, x \rangle$
- $\langle x, x \rangle > 0$ if $x \neq 0$ and $\langle 0, 0 \rangle = 0$.

Examples:

- Let $x = (a_1, \dots, a_n)$, $y = (b_1, \dots, b_n) \in \mathbb{F}^n$. Then $\langle x, y \rangle = \sum_{i=1}^n a_i \overline{b_i}$ is called the standard inner product.
- For $V = C[0, 1] = \{f : [0, 1] \rightarrow \mathbb{R} : f \text{ is continuous}\}$, $\langle f, g \rangle = \int_0^1 f(t)g(t)dt$ is an inner product on V .
- (Frobenius inner product)

Let $V = M_{n \times n}(\mathbb{F})$. For $A, B \in V$, let $\langle A, B \rangle = \text{tr}(B^* A)$ where B^* is the adjoint of B . The four properties of an inner product are satisfied on V as follows:

(1)

$$\begin{aligned}
 \langle A_1 + A_2, B \rangle &= \text{tr}(B^*(A_1 + A_2)) \\
 &= \text{tr}(B^* A_1 + B^* A_2) \\
 &= \text{tr}(B^* A_1) + \text{tr}(B^* A_2) \\
 &= \langle A_1, B \rangle + \langle A_2, B \rangle
 \end{aligned}$$

(2)

$$\begin{aligned}
 \langle cA, B \rangle &= \text{tr}(B^* cA) \\
 &= c \cdot \text{tr}(B^* A) \\
 &= c \langle A, B \rangle
 \end{aligned}$$

(3)

$$\begin{aligned}
 \overline{\langle A, B \rangle} &= \overline{\text{tr}(B^* A)} \\
 &= \text{tr}(A^*) \\
 &= \langle B, A \rangle
 \end{aligned}$$

(4)

$$\begin{aligned}
 \langle A, A \rangle &= \text{tr}(A^* A) \\
 &= \sum_{i=1}^n \sum_{j=1}^n \overline{A_{ji}} A_{ji} \\
 &= \sum_{i,j=1}^n |A_{ji}|^2 \\
 &> 0
 \end{aligned}$$

(When $A \neq 0$)

Definition 2 The *adjoint* of an $m \times n$ matrix A is the $n \times m$ matrix A^* such that $(A^*)_{ij} = \overline{A_{ji}}$.

Theorem 6.1.

- $\langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$
- $\langle x, cy \rangle = \bar{c}\langle x, y \rangle$
- $\langle x, 0 \rangle = \langle 0, x \rangle = 0$
- $\langle x, x \rangle = 0$ if and only if $x = 0$
- If $\langle x, y \rangle = \langle x, z \rangle$ for every $x \in V$, then $y = z$.

Definition 3 Let V be an inner product space. For $x \in V$, the *norm* of x is $\|x\| = \sqrt{\langle x, x \rangle}$.

Theorem 6.2. Let V be an inner product space over F . For $x, y \in V$ and $c \in \mathbb{F}$.

- (a) $\|cx\| = |c| \cdot \|x\|$
- (b) $\|x\| = 0$ if and only if $x = 0$ and $\|x\| \geq 0$ for any x .
- (c) (Cauchy-Schwarz Inequality) $|\langle x, y \rangle| \leq \|x\| \|y\|$
- (d) (Triangle Inequality) $\|x + y\| \leq \|x\| + \|y\|$

Proof. (Outline)

- (c) Expand $\|x - cy\|^2$ and apply with $c = \frac{\langle x, y \rangle}{\langle y, y \rangle}$
- (d) Note that $\langle x, y \rangle + \langle y, x \rangle = 2\operatorname{Re}\langle x, y \rangle$ and $\operatorname{Re}\langle x, y \rangle \leq |\langle x, y \rangle|$.

Proof.

- (c) Note, the inequality holds if $y = 0$. Suppose $y \neq 0$. Let $c \in \mathbb{F}$. Then

$$0 \leq \langle x - cy, x - cy \rangle = \langle x, x \rangle - c\langle y, x \rangle - \bar{c}\langle x, y \rangle + c\bar{c}\langle y, y \rangle.$$

Let $c = \frac{\langle x, y \rangle}{\langle y, y \rangle}$. Then

$$\begin{aligned} 0 &\leq \langle x, x \rangle - \frac{\langle x, y \rangle \langle y, x \rangle}{\langle y, y \rangle} \\ &= \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2}. \end{aligned}$$

Thus, $|\langle x, y \rangle|^2 \leq \|x\|^2 \|y\|^2$.

(d)

$$\begin{aligned}
\|x + y\|^2 &= \langle x + y, x + y \rangle \\
&= \langle x, x \rangle + \langle y, x \rangle + \langle x, y \rangle + \langle y, y \rangle \\
&= \langle x, x \rangle + 2\operatorname{Re}\langle x, y \rangle + \langle y, y \rangle \\
&\leq \|x\|^2 + 2|\langle x, y \rangle| + \|y\|^2 \\
&= \|x\|^2 + 2\|x\|\|y\| + \|y\|^2 && \text{(By Cauchy-Schwartz)} \\
&= (\|x\| + \|y\|)^2.
\end{aligned}$$

□

Definition 4

- Vectors $x, y \in V$ are called *orthogonal* if $\langle x, y \rangle = 0$.
- A set of vectors $S \subseteq V$ is called *orthogonal* if any two distinct vectors are orthogonal.
- S is called *orthonormal* if it is orthogonal and $\|x\| = 1$ for every $x \in S$.

Example: Let $H = \{f : [0, 2\pi] \rightarrow \mathbb{C}\}$. Let $\langle f, g \rangle = \frac{1}{2\pi} \int_0^{2\pi} f(t) \overline{g(t)} dt$. Let $f_n(t) = e^{int} = \cos(nt) + i \sin(nt)$. Then $S = \{f_n(t) : n \in \mathbb{N}\}$ is orthonormal.

Proof. If $n \neq m$, then

$$\begin{aligned}
\langle f_n, f_m \rangle &= \frac{1}{2\pi} \int_0^{2\pi} e^{int} \overline{e^{imt}} dt \\
&= \frac{1}{2\pi} \int_0^{2\pi} e^{int} e^{-imt} dt \\
&= \frac{1}{2\pi} \int_0^{2\pi} e^{it(n-m)} dt \\
&= \frac{1}{2\pi} \int_0^{2\pi} \cos(t(n-m)) + i \sin(t(n-m)) dt \\
&= \frac{1}{2\pi} [\sin(t(n-m)) - \cos(t(n-m))]_0^{2\pi} \\
&= 0.
\end{aligned}$$

If $n = m$, then

$$\langle f_n, f_n \rangle = \frac{1}{2\pi} \int_0^{2\pi} e^{int} e^{-int} dt = 1.$$

□

Odd Town Problem: There are n people living in an odd town and they form clubs. A club must contain an odd number of members and for any two distinct clubs there must be an even (possibly zero) number of people in both of them. What is the maximum number of clubs that can be formed?

Proof. Let S_1, \dots, S_m denote these clubs. Let $v_i = \begin{pmatrix} v_{i1} \\ \vdots \\ v_{in} \end{pmatrix}$ when $v_{ij} = \begin{cases} 1 & j \in S_i \\ 0 & \text{else} \end{cases}$.

Then $v_i \in \mathbb{F}_2^n$. Then v_1, \dots, v_m are linearly independent. Suppose $\sum_{i=1}^m c_i v_i = 0$. Then

$$0 = \left\langle \sum_{i=1}^m c_i v_i, v_j \right\rangle = \sum_{i=1}^m c_i \langle v_i, v_j \rangle = c_i.$$

Then, $m \leq \dim(\mathbb{F}_2^n) = n$. Therefore, the number of clubs is at most n . \square

6.2 The Gram-Schmidt orthogonalization

Definition 5 An ordered basis which is orthonormal is called an *orthonormal basis*.

Theorem 6.3. Suppose $S = \{v_1, \dots, v_k\}$ is an orthogonal subset of V such that $v_i \neq 0$. For $y \in \text{span}(S)$,

$$y = \sum_{i=1}^k \frac{\langle y, v_i \rangle}{\|v_i\|^2} v_i.$$

Proof. Let $y \in \text{span}(S)$. Then $y = \sum_{i=1}^k a_i v_i$ and so $\langle y, v_j \rangle = \sum_{i=1}^k a_i \langle v_i, v_j \rangle = a_j \langle v_j, v_j \rangle$. Thus, $a_j = \frac{\langle y, v_j \rangle}{\langle v_j, v_j \rangle} = \frac{\langle y, v_j \rangle}{\|v_j\|^2}$. \square

Corollary 4 Any orthogonal set of non-zero vectors is linearly independent.

Proof. Suppose $\sum_{i=1}^k a_i v_i = 0$. Then $a_i = \frac{\langle 0, v_i \rangle}{\|v_i\|^2} = 0$ by Theorem 6.3. \square

Theorem 6.4 (Graham-Schmidt algorithm). Let $S = \{w_1, \dots, w_n\}$ be a linearly independent subset. Define $S' = \{v_1, \dots, v_n\}$ as follows, $v_1 := w_1$ and for $k \geq 2$

$$v_k = w_k - \sum_{j=1}^{k-1} \frac{\langle w_k, v_j \rangle}{\|v_j\|^2} v_j.$$

Then S' is orthogonal and $\text{span}(S') = \text{span}(S)$.

Proof. Let $S_k = \{w_1, \dots, w_k\}$. We will use induction on k .

- Base Case: For $k = 1$, clearly $\text{span}(S'_1) = \text{span}(S_1)$.

- Induction Step: Suppose $S'_{k-1} = \{v_1, \dots, v_{k-1}\}$ has been constructed so that S'_{k-1} is orthogonal and $\text{span}(S'_{k-1}) = \text{span}(S_k)$. Let $v_k = w_k - \sum_{i=1}^{k-1} \frac{\langle w_k, v_i \rangle}{\|v_i\|^2} v_i$. Then we have the following properties:
 - We know $v_k \neq 0$ because otherwise, $w_k - \sum_{i=1}^{k-1} \frac{\langle w_k, v_i \rangle}{\|v_i\|^2} v_i = 0$ and $v_k \in \text{span}(S'_{k-1}) = \text{span}(S_{k-1})$, but S is linearly independent.
 - For $j \leq k-1$,

$$\begin{aligned}
 \langle v_k, v_j \rangle &= \langle w_k, v_j \rangle - \sum_{i=1}^{k-1} \frac{\langle w_k, v_i \rangle}{\|v_i\|^2} \langle v_i, v_j \rangle && \text{(Note, } \langle v_i, v_j \rangle = 0 \text{ if } i \neq j) \\
 &= \langle w_k, v_j \rangle - \frac{\langle w_k, v_j \rangle}{\|v_j\|^2} \langle v_j, v_j \rangle && \text{(By inductive hypothesis)} \\
 &= 0.
 \end{aligned}$$

Thus, $S_k = \{v_1, \dots, v_k\}$ is orthogonal. From $v_k = w_k - \sum_{i=1}^{k-1} \frac{\langle w_k, v_i \rangle}{\|v_i\|^2} v_i$, we have $v_k \in \text{span}(w_k, v_1, \dots, v_{k-1}) \in \text{span}(w_1, \dots, w_k) = \text{span}(S_k)$. Thus, $\text{span}(S'_k) \subseteq \text{span}(S_k)$. Also, $w_k \in \text{span}(v_1, \dots, v_k)$ and so $\text{span}(S_k) \subseteq \text{span}(S'_k)$.

□

Example: (Legendre Polynomials of the form $\frac{v_k(x)}{v_k(1)}$)

Let $P_2(\mathbb{R})$ with the standard basis $\beta = \{1, x, x^2\}$. Consider $\langle f, g \rangle = \int_{-1}^1 f(x)g(x)dx$. Replace β with an orthogonal basis.

Proof. Let $\beta = \{w_1, w_2, w_3\}$. Then

- $v_1 = w_1 = 1, \|v_1\|^2 = \langle v_1, v_1 \rangle = 2$
- $v_2 = w_2 - \frac{\langle w_2, v_1 \rangle}{\|v_1\|^2} v_1 = x$
- $v_3 = w_3 - \frac{\langle w_3, v_2 \rangle}{\|v_2\|^2} v_2 - \frac{\langle w_3, v_1 \rangle}{\|v_1\|^2} v_1 = x^2 - \frac{1}{3}$.

Thus, $\alpha = \{2, x, x^2 - \frac{1}{3}\}$ is an orthonormal basis for $P_2(\mathbb{R})$.

□

Theorem 6.5. Let V be a finite-dimensional inner product space and let $\beta = \{v_1, \dots, v_n\}$ be an orthonormal basis for V . Then for every $x \in V$, $x = \sum_{i=1}^n \langle x, v_i \rangle v_i$.

Corollary 7 Let $\beta = \{v_1, \dots, v_n\}$ be orthonormal, let $T : V \rightarrow V$ be linear, and let $A = [T]_\beta$. Then $A_{ij} = \langle T(v_j), v_i \rangle$.

Fourier coefficient of x relative to β is $\langle x, y \rangle$ where $y \in \beta$.

Example: Let $S = \{e^{int} : n \in \mathbb{Z}\}$ in H with $\langle f, g \rangle = \frac{1}{2\pi} \int_0^{2\pi} f(t) \overline{g(t)} dt$. Find the Fourier coefficients of $f(t) = 1$.

Proof.

- If $n = 0$, $\langle f, 1 \rangle = \pi$
- If $n \neq 0$, $\langle f, f_n \rangle = \frac{1}{2\pi} \int_0^{2\pi} t e^{-int} dt = -\frac{1}{in}$

□

Definition 6 Let S be a non-empty subset of V . Define $S^\perp = \{x \in V : \langle x, y \rangle = 0 \text{ for every } y \in S\}$.

Note: S^\perp is a subspace of V .

Theorem 6.6. Let W be a finite-dimensional subspace of an inner product space V and let $y \in V$. Then there exist unique $u \in W$ and $z \in W^\perp$ such that $y = u + z$. Furthermore, if $\{v_1, \dots, v_k\}$ is an orthonormal basis for W , then $u = \sum \langle y, v_i \rangle v_i$.

Proof. (Outline)

- Let $u = \sum \langle y, v_i \rangle v_i$ and $z = y - u$. Check that $z \in W^\perp$.
- For the uniqueness $u - u' \in W$ and $z' - z \in W^\perp$.

Proof. Let $u = \sum \langle y, v_i \rangle v_i$ and $z = y - u$. Then

$$\begin{aligned} \langle z, v_j \rangle &= \langle y - u, v_j \rangle \\ &= \langle y, v_j \rangle - \sum_{i=1}^k \langle y, v_i \rangle \langle v_i, v_j \rangle \\ &= \langle y, v_j \rangle. \end{aligned}$$

Then, for uniqueness, $u' + z' = y = u + z$ and $u' + u = z - z' \in W \cap W^\perp = \{0\}$. So, $u' = u$ and $z' = z$. □

Corollary 9 For any $x \in W$, $\|y - x\| \geq \|y - u\|$ and if $\|y - x\| = \|y - u\|$ then $x = u$.

Proof.

$$\begin{aligned}
 \|y - x\|^2 &= \|u + z - x\|^2 \\
 &= \|u - x + z\|^2 \\
 &= \langle u - x + z, u - x + z \rangle \\
 &= \|u - x\|^2 + \|z\|^2 \\
 &\geq \|z\|^2 \\
 &= \|y - u\|^2
 \end{aligned}$$

□

Theorem 6.7. Suppose $S = \{v_1, \dots, v_k\}$ is an orthonormal set in an n -dimensional inner product space V . Then

- (a) S can be extended to an orthonormal basis $\{v_1, \dots, v_n\}$ for V .
- (b) $\{v_{k+1}, \dots, v_n\}$ is an orthonormal basis for $(\text{span}(S))^\perp$.
- (c) If W is a subspace of V , then $\dim(W) + \dim(W^\perp) = \dim(V)$.

6.3 The adjoint of a linear operator

Theorem 6.8. Let V be a finite-dimensional inner product space over \mathbb{F} , and let $g : V \rightarrow \mathbb{F}$ be a linear transformation. Then there exists a unique vector $y \in V$ such that $g(x) = \langle x, y \rangle$ for every $x \in V$.

Proof. (Outline) Take an orthonormal basis $\beta = \{v_1, \dots, v_n\}$ and define $y = \sum \overline{g(v_i)} v_i$.

Proof. Let $\{v_1, \dots, v_n\}$ be an orthonormal basis for V . Let $y = \sum_{i=1}^n \overline{g(v_i)} v_i \in V$. Let $h : V \rightarrow \mathbb{F}$ be defined by $h(x) = \langle x, y \rangle$. Then h is a linear transformation. In addition,

$$\begin{aligned}
 h(v_j) &= \langle v_j, y \rangle \\
 &= \left\langle v_j, \sum_{i=1}^n \overline{g(v_i)} v_i \right\rangle \\
 &= \sum_{i=1}^n g(v_i) \langle v_j, v_i \rangle \\
 &= g(v_j).
 \end{aligned}$$

Thus, $h(v_j) = g(v_j)$ for all $j = 1, \dots, n$. Thus, $h = g$. Suppose $\langle x, y \rangle = \langle x, y' \rangle$ for all $x \in V$. Then $y = y'$. □

Theorem 6.9. Let V be a finite-dimensional inner product space and let $T : V \rightarrow V$ be linear. Then there exists a unique function $T^* : V \rightarrow V$ such that $\langle T(x), y \rangle = \langle x, T^*(y) \rangle$ for all x, y . Furthermore, T^* is linear.

Proof. (Outline)

- Fix y . Check that $g(x) = \langle T(x), y \rangle$ is linear.
- Theorem 6.8 gives unique y' such that $g(x) = \langle x, y' \rangle$. Define $T^*(y) = y'$.
- Note that T^* is a function and check that it is linear.

Proof. Fix $y \in V$. Consider $g(x) = \langle T(x), y \rangle$. Then $g : V \rightarrow \mathbb{F}$ is linear. By Theorem 6.8, there exists a unique y' such that $g(x) = \langle x, y' \rangle$. Let $T^*(y) = y'$. Then $T^* : V \rightarrow V$ and $\langle T(x), y \rangle = g(x) = \langle x, T^*(y) \rangle$. In addition, T^* is linear. Let $y_1, y_2 \in V$ and $c \in \mathbb{F}$. Then

$$\begin{aligned} \langle x, T^*(cy_1 + y_2) \rangle &= \langle T(x), cy_1 + y_2 \rangle \\ &= c\langle T(x), y_1 \rangle + \langle T(x), y_2 \rangle \\ &= \langle x, cT^*(y_1) \rangle + \langle x, T^*(y_2) \rangle \\ &= \langle x, cT^*(y_1) + T^*(y_2) \rangle. \end{aligned}$$

Thus, $T^*(cy_1 + y_2) = cT^*(y_1) + T^*(y_2)$. Suppose $\langle T(x), y \rangle = \langle x, U(y) \rangle$ for all $x, y \in V$. Then, $\langle T(x), y \rangle = \langle x, T^*(y) \rangle$, which implies $T^* = U$. \square

Theorem 6.10. Let V be a finite-dimensional inner product space and let β be an orthonormal ordered basis for V . If T is a linear operator on V , then $[T^*]_\beta = [T]_\beta^*$.

Proof. Let $A = [T^*]_\beta$ and $B = [T]_\beta$ where $\beta = \{v_1, \dots, v_n\}$. Note, $B_{ij} = \langle T(v_j), v_i \rangle$. Then,

$$\begin{aligned} A_{ij} &= \langle T^*(v_j), v_i \rangle \\ &= \overline{\langle v_i, T^*(v_j) \rangle} \\ &= \overline{\langle T(v_i), v_j \rangle} \\ &= \overline{B_{ji}}. \end{aligned}$$

\square

Theorem 6.11. Let V be an inner-product space, and let T, U be linear operators on V . Then

(a) $(T + U)^* = T^* + U^*$

$$(b) \quad (cT)^* = \bar{c}T^*$$

$$(c) \quad (TU)^* = U^*T^*$$

Proof.

$$\begin{aligned} \langle x, (TU)^*y \rangle &= \langle (TU)x, y \rangle \\ &= \langle T(Ux), y \rangle \\ &= \langle Ux, T^*y \rangle \\ &= \langle x, U^*T^*y \rangle \end{aligned}$$

□

$$(d) \quad T^{**} = T$$

$$(e) \quad I^* = I$$

6.4 Normal and self-adjoint operators

Lemma 15. Let T be a linear operator on a finite-dimensional inner product space V . If T has an eigenvector, then so does T^* .

Theorem 6.14 (Schur). Let T be a linear operator on a finite-dimensional inner product space V and suppose that the characteristic polynomial of T splits. Then there exists an orthonormal basis β for V such that $[T]_\beta$ is upper triangular.

Proof. (Outline)

- Show that W^\perp is T -invariant and $\dim(W^\perp) = n - 1$.
- The characteristic polynomial of T_{W^\perp} divides $f_T(t)$ (so it splits) and by **IH** for some γ , $[T_{W^\perp}]_\gamma$ is upper-triangular.
- Let $\beta = \gamma \cup \{z\}$ Then $[T]_\beta$ is upper-triangular.

Proof. Let $n = \dim(V)$.

- Base step: Let $n = 1$. Then clearly the statement holds.
- Induction Step: Let the statement be true for some $n - 1 \geq 1$. Then we can assume T^* has an eigenvector z such that $\|z\| = 1$ and $T^*(z) = \lambda z$ by Lemma 15. Let

$W = \text{span}(z)$. We claim W^\perp is T -invariant. Let $y \in W^\perp$. Then $\langle y, z \rangle = 0$. Now, for $w \in W$,

$$\begin{aligned} \langle T(y), w \rangle &= \langle y, T^*(w) \rangle \\ &= \langle y, cT^*(z) \rangle && \text{(For some } c \in \mathbb{F}) \\ &= \langle y, c\lambda z \rangle \\ &= \bar{c}\bar{\lambda} \langle y, z \rangle \\ &= 0. \end{aligned}$$

Thus, $T(y) \in W^\perp$. In addition, $\dim(W^\perp) = n - \dim(W) = n - 1$ and $f|_{T_{W^\perp}}$ divides $f|_T$ and so it splits. By inductive hypothesis, there exists an orthonormal basis γ such that $[T_{W^\perp}]_\gamma$ is upper-triangular. Let $\beta = \gamma \cup \{z\}$. Then $[T]_\beta$ is upper-triangular because $T(w) \in W^\perp \in \text{span}(\gamma)$.

□

Note: If V has an orthonormal basis of eigenvectors of T , then $TT^* = T^*T$.

Definition 7 $T : V \rightarrow V$ is *normal* if $TT^* = T^*T$. $A \in M_{n \times n}(\mathbb{F})$ is *normal* if $AA^* = A^*A$.

Note: If there exists an orthonormal basis β consisting of eigenvectors of T , then T is normal.

Proof. Suppose there exists a basis β consisting of eigenvectors of T . Then $[T]_\beta$ is diagonal and $[T^*]_\beta = ([T]_\beta)^*$ is diagonal. Therefore,

$$[TT^*]_\beta = [T]_\beta [T^*]_\beta = [T^*]_\beta [T]_\beta = [T^*T]_\beta.$$

Thus, $TT^* = T^*T$.

□

Examples:

- If $A \in M_{n \times n}(\mathbb{R})$ is symmetric, then A is normal.
- If $A \in M_{n \times n}(\mathbb{R})$ is asymmetric, $A^T = -A$, and so $AA^* = -A^2 = A^*A$.

Theorem 6.15. Let T be a normal operator on V . Then

(a) $\|T(x)\| = \|T^*(x)\|$

Proof.

$$\begin{aligned} \|Tx\|^2 &= \langle T(x), T(x) \rangle \\ &= \langle T^*Tx, x \rangle \\ &= \langle TT^*x, x \rangle \\ &= \langle T^*x, T^*x \rangle \\ &= \|T^*x\|^2 \end{aligned}$$

□

(b) $T - cI$ is normal for every $c \in \mathbb{F}$.

Proof.

$$\begin{aligned}
 (T - cI)(T - cI)^* &= (T - cI)(T^* - \bar{c}I) \\
 &= TT^* - cT^* - \bar{c}T - c\bar{c}I \\
 &= T^*T - cT^* - \bar{c}T - c\bar{c}I \\
 &= (T - cI)^*(T - cI)
 \end{aligned}$$

□

(c) If x is an eigenvector of T , then x is an eigenvector of T^* .

Proof. We can show

$$\begin{aligned}
 0 &= \langle (T - \lambda I)x, (T - \lambda I)x \rangle \\
 &= \|(T - \lambda I)x\|^2 \\
 &= \|(T - \lambda I)^*x\|^2 && \text{(From (a) and (b))} \\
 &= \langle (T^* - \bar{\lambda}I)x, (T^* - \bar{\lambda}I)x \rangle.
 \end{aligned}$$

Thus, $T^*x = \bar{\lambda}x$.

□

(d) If λ_1, λ_2 are distinct eigenvalues of T with eigenvectors x_1, x_2 , then $\langle x_1, x_2 \rangle = 0$.

Proof. We can show

$$\begin{aligned}
 \lambda_1 \langle x_1, x_2 \rangle &= \langle \lambda_1 x_1, x_2 \rangle \\
 &= \langle T(x_1), x_2 \rangle \\
 &= \langle x_1, T^*(x_2) \rangle \\
 &= \langle x_1, \bar{\lambda}_2 x_2 \rangle && \text{(From (c))} \\
 &= \lambda_2 \langle x_1, x_2 \rangle,
 \end{aligned}$$

which implies $(\lambda_1 - \lambda_2)\langle x_1, x_2 \rangle = 0$. Since $\lambda_1 \neq \lambda_2$, then $\langle x_1, x_2 \rangle = 0$.

□

Theorem 6.16. Let T be a linear operator on a finite-dimensional complex inner-product space V . Then T is normal if and only if there exists an orthonormal basis for V consisting of eigenvectors of T .

Proof. (Outline) Suppose T is normal.

- T splits over \mathbb{C} and so apply Schur's lemma to get an orthonormal basis $\beta = \{v_1, \dots, v_n\}$.
- $A := [T]_\beta$ is upper-triangular and so $T(v_1) = A_{11}v_1$.

- Show that e_k is an eigenvector by induction on k using the fact that $A_{jk} = \langle T(v_k), v_j \rangle$.

Proof. Let T be a linear operator on a finite-dimensional complex inner-product space V .

\Rightarrow Suppose T is normal. Since we are over \mathbb{C} , the characteristic polynomial of T splits. By Schur's Theorem, there exists an orthonormal basis $\beta = \{v_1, \dots, v_n\}$ such that $A = [T]_\beta$ is upper-triangular. We will show that v_1, \dots, v_n are in addition eigenvectors of T . We will use induction on n .

- Base Case: Let $n = 1$. We have $T(v_1) = A_{11}v_1$ and so v_1 is an eigenvector.
- Induction Step: Suppose v_1, \dots, v_{k-1} are eigenvectors of T with eigenvalues $\lambda_1, \dots, \lambda_{k-1}$. Then $T(v_k) = \sum_{i=1}^k A_{ik}v_i$ and in addition, $A_{ik} = \langle T(v_k), v_i \rangle = \langle v_k, T^*(v_i) \rangle$. For $i < k$, we have

$$\begin{aligned} A_{ik} &= \langle v_k, \overline{\lambda_i} v_i \rangle \\ &= \lambda_i \langle v_k, v_i \rangle \\ &= 0. \end{aligned}$$

Thus, $T(v_k) = A_{kk}v_k$. So, v_k is an eigenvector of T .

\Leftarrow Suppose β is an orthonormal basis consisting of eigenvectors of T . Then $[T]_\beta$ is diagonal, and so $[T^*]_\beta = [T]_\beta^*$ is diagonal. Thus,

$$\begin{aligned} [TT^*]_\beta &= [T]_\beta [T^*]_\beta \\ &= [T^*]_\beta [T]_\beta \\ &= [T^*T]_\beta. \end{aligned}$$

Thus, $TT^* = T^*T$.

□

Definition 8 $T : V \rightarrow V$ is *self-adjoint* (Hermitian) if $T = T^*$. $A \in M_{n \times n}(\mathbb{F})$ is *Hermitian* if $A = A^*$.

Lemma 19. Let T be a Hermitian operator on a finite-dimensional inner product space V . Then,

- (a) All eigenvalues of T are real.

Proof. Let λ be an eigenvalue correspond with the eigenvector v . Then $\lambda v = T(v) = T^*(v) = \overline{\lambda}v$ and so $\lambda = \overline{\lambda}$. Thus, $\lambda \in \mathbb{R}$. □

(b) If V is a real inner product space, then the characteristic polynomial splits.

Proof. Suppose T is self-adjoint. Let β be an orthonormal basis for T . Then $A = [T]_\beta$ is self-adjoint. Let $L_A : \mathbb{C}^n \rightarrow \mathbb{C}^n$ where $n = \dim(V)$ defined by $L_A(x) = Ax$. Then $[L_A]_\gamma = A$ where γ is the standard basis. Thus, L_A is self-adjoint. Thus, f_{L_A} splits over \mathbb{C} . Then $f_{L_A}(t) = (-1)^n(t - \lambda_1) \dots (t - \lambda_n)$ where λ_i is an eigenvalue of L_A . By (a), $\lambda_i \in \mathbb{R}$. Thus, f_{L_A} splits over \mathbb{R} . Therefore, $f_T = f_A = f_{L_A}$ splits over \mathbb{R} . \square

Theorem 6.17. Let T be a linear operator on a finite-dimensional real inner-product space V . Then T is Hermitian if and only if there exists an orthonormal basis for V consisting of eigenvectors of T .

Proof. (Outline)

The characteristic polynomial splits and so we may apply Schur's lemma. $A := [T]_\beta$ is upper-triangular and so A^* . Thus it must be a diagonal matrix.

Proof. Let T be a linear operator on a finite-dimensional real inner-product space V .

\implies Suppose T is self-adjoint. By Lemma 19, f_T splits. By Schur's Theorem, there exists an orthonormal basis β such that $A = [T]_\beta$ is an upper-triangular. Then, $A = [T]_\beta = [T^*]_\beta$. Thus, A and A^* are upper-triangular. Thus, A is diagonal and so β contains eigenvectors of T .

\impliedby Suppose there is an orthonormal basis β consisting of eigenvectors of T . Then $[T]_\beta$ is diagonal with entries from \mathbb{R} . So, $[T^*]_\beta = [T]_\beta^* = [T]_\beta$. Thus, $T^* = T$. Therefore, T is self-adjoint. \square

6.5 Unitary and orthogonal operators and their matrices

Definition 9 Let V be a finite-dimensional inner-product space over F and let $T : V \rightarrow V$ be linear. If $\|T(x)\| = \|x\|$ for every $x \in V$, then T is called *unitary* if $F = \mathbb{C}$ and *orthogonal* if $F = \mathbb{R}$.

Theorem 6.18. Let T be a linear operator on a finite-dimensional inner-product space V . Then the following statements are equivalent.

- (a) $TT^* = T^*T = I$
- (b) $\langle T(x), T(y) \rangle = \langle x, y \rangle$ for all $x, y \in V$
- (c) If β is an orthonormal basis, then so is $T(\beta)$.

(d) There exists an orthonormal basis β such that $T(\beta)$ is orthonormal.

(e) $\|T(x)\| = \|x\|$ for every x .

Proof. (Outline)

- (d) \implies (e) Let $\beta = \{v_1, \dots, v_n\}$ be orthonormal such that $T(\beta)$ is orthonormal. Take $x \in V$. Then $x = \sum a_i v_i$ and

$$\|x\|^2 = \sum |a_i|^2 = \sum \|T(x)\|^2.$$

- (e) \implies (a) $\langle x, x \rangle = \langle x, T^*T(x) \rangle$ by (e). Thus $\langle x, (I - T^*T)(x) \rangle = 0$ for every x . Set $U := I - T^*T$. Then U is self-adjoint and so there is an orthonormal basis consisting of eigenvectors of U . Check that $U(x) = \lambda x$ implies $\lambda = 0$; $U = T_0$; $T^*T = I$; $TT^* = I$ as well because $[T]_\beta$ is a square matrix.

Definition 10 A square matrix A is called *orthogonal* if $A^t A = A A^t = I$ and *unitary* if $A^* A = A A^* = I$.

Note:

- $AA^* = I$ if and only if A are orthonormal.
- $A^*A = I$ if and only if columns of A are orthonormal.
- If λ is an eigenvalue of a unitary (orthogonal) matrix, then $|\lambda| = 1$.

Definition 11 $A, B \in M_{n \times n}(\mathbb{C})$ ($A, B \in M_{n \times n}(\mathbb{R})$) are *unitarily (orthogonally) equivalent* if there exists a unitary (orthogonal) matrix P such that $A = P^{-1}BP$.

Theorem 6.19. Let $A \in M_{n \times n}(\mathbb{C})$. Then A is normal if and only if A is unitarily equivalent to a diagonal matrix.

Theorem 6.20. Let $A \in M_{n \times n}(\mathbb{R})$. Then A is symmetric if and only if A is orthogonally equivalent to a diagonal matrix.

Theorem 6.21 (Schur). Let $A \in M_{n \times n}(\mathbb{F})$ and suppose $f_A(t)$ splits over \mathbb{F} .

- (a) If $\mathbb{F} = \mathbb{C}$, then A is unitarily equivalent to a complex upper-triangular matrix.
- (b) If $\mathbb{F} = \mathbb{R}$, then A is orthogonally equivalent to a real upper-triangular matrix.