

MAT 473: Intermediate Real Analysis II

Trey Manuszak
Arizona State University

April 24, 2020

Problem 49. Prove Theorem 22.5 parts (iii) and (iv): Let $f, g : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ be integrable functions.

(iii) Suppose that $f \leq g$. Prove that $\int f \leq \int g$.

Proof. Let $f, g : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ be integrable functions and $f \leq g$. Then, by linearity,

$$0 \leq \int (g - f) = \int g - \int f,$$

which implies $\int f \leq \int g$. □

(iv) Prove that $|\int f| \leq \int |f|$.

Proof. Let $f : \mathbb{R} \rightarrow \overline{\mathbb{R}}$ be integrable. Since $|f|$ is measurable and bounded and $-|f| \leq f \leq |f|$, by linearity and monotonicity, we get

$$-\int |f| \leq \int f \leq \int |f|.$$

□

Problem 50. Compute the value of the limit

$$\lim_{n \rightarrow \infty} \int_0^\infty \left(1 + \frac{x}{n}\right)^{-n} \cos \frac{x}{n} dx.$$

Justify every step of your argument. (Hint: use the monotone convergence theorem, and the theorem on equality of the Riemann and Lebesgue integrals when both apply, to show that e^{-x} is integrable on $[0, \infty]$. Then use the dominated convergence theorem.)