

# Uniqueness of fit-width solution

---

Let  $(a, b)$  be a minimal fit-width point. **Prove** there is no  $b'$  such that  $(a, b')$  is a fit-width point when  $r \geq 1$  and  $(a, b)$  corresponds to the best solution, meaning that  $s > s_h$  for  $(a_h, b_h)$  minimal fit-width point.

If you didn't read the script, here's a self-contained problem statement:

Let:

$$r \in [1, \infty),$$

$$n \in \mathbb{N} \text{ (for this task } \mathbb{N} \text{ does not include } 0),$$

$$W = \{ (a, b) \in \mathbb{N}^2 \mid n \leq ab \wedge a \geq rb \} \text{ (the set of "fit-width" points),}$$

$$H = \{ (a, b) \in \mathbb{N}^2 \mid n \leq ab \wedge rb \geq a \} \text{ (the set of "fit-height" points),}$$

$$(a_w, b_w) = \arg \min_{(a,b) \in W} a,$$

$$(a_h, b_h) = \arg \min_{(a,b) \in H} b.$$


$$\text{Assume } a_w < rb_h.$$

Prove there exists no other  $b \neq b_w$  such that  $(a_w, b) \in W$ .

Try to prove this yourself after reading the main part of the script, **without** reading the appendix. Three hints are provided in case you get stuck; attempt to solve without them first, then try again one hint at a time. Don't jump to the end directly.

## Hint 1

Recall graphs A ( $n = 8, r = 5$ ) and C ( $n = 23, r = 3.9$ ) from the script:



How do the fit-width and fit-height solutions compare in each of them?

## Hint 2

---

By hint 1 a viable proof path is to show that if such a  $b'$  exists then one can find a point  $(a', b')$  which is fit-height.

Try to prove that if  $b'$  exists then specifically  $(a - 1, b')$  is fit-height.

## Hint 3

---

Attached are proof attempts from LLMs. They present various useful approaches but have subtle mistakes. Find these mistakes and use the proofs as inspiration for your own!

### Proof 1

Courtesy of DeepSeek v4.

---

#### Setup

Let the fit-width solution be  $(a, b)$  where

$b = \lceil n/a \rceil$  and  $a$  is the smallest integer satisfying  $a \geq r \cdot b$ .

Its square side is  $s_{\text{fw}} = w/a$ .

A **valid fit-width solution** with the same  $a$  and side length  $s_{\text{fw}}$  uses the same  $a$  and any row count  $b'$  with

$$\lceil n/a \rceil \leq b' \leq \lfloor a/r \rfloor.$$

The algorithm picks the minimal  $b' = b$ .

---

#### Claim

If  $s_{\text{fw}} > s_{\text{fh}}$  for **every** fit-height solution, then there is no  $b' > b$  with  $b' \leq a/r$ .

In other words, the feasible interval for  $b'$  collapses to a single value  $b = \lfloor a/r \rfloor = \lceil n/a \rceil$ .

---

#### Proof

Assume, for contradiction, that such a  $b'$  exists. Then  $b + 1 \leq \lfloor a/r \rfloor$ , so

$$a \geq r(b + 1). \tag{1}$$

Let  $k = \lfloor a/r \rfloor$ . Clearly  $k \geq b + 1$ .

Now examine the fit-height condition for row count  $k$ :

$$\lceil n/k \rceil \leq rk.$$

We know  $n \leq ab$  (from  $b = \lceil n/a \rceil$ ). Thus

$$\frac{n}{k} \leq \frac{ab}{k} \leq \frac{ab}{b+1} = a - \frac{a}{b+1}.$$

From (1),  $\frac{a}{b+1} \geq r$ , so

$$\frac{n}{k} \leq a - r.$$

Since  $rk > a - r$  (because  $k = \lfloor a/r \rfloor > a/r - 1$ ), we obtain

$$\frac{n}{k} < rk.$$

Hence  $\lceil n/k \rceil \leq rk$ , i.e.,  $k$  satisfies the fit-height condition.

The fit-height algorithm will therefore find an optimal row count  $b_{\text{fh}} \leq k \leq a/r$ .

Then

$$s_{\text{fh}} = \frac{h}{b_{\text{fh}}} \geq \frac{h}{a/r} = \frac{rh}{a} = \frac{w}{a} = s_{\text{fw}},$$

contradicting the strict optimality of the fit-width solution.

Therefore, no such  $b'$  can exist when the fit-width solution is strictly better than all fit-height solutions. The pair  $(a, b)$  is the unique grid achieving the maximal square side length.

## Proof 2

Courtesy of [QED-Nano](#), a tiny 4B open-source model developed exclusively for math proofs.

---

### Proof.

Define for each  $b \in \mathbb{N}$

$$f(b) = \max \left\{ \lceil rb \rceil, \left\lceil \frac{n}{b} \right\rceil \right\}.$$

For a pair  $(a, b) \in \mathbb{N}^2$  we have

$$(a, b) \in W \iff a \geq rb \text{ and } ab \geq n.$$

Because  $a, b$  are integers,

$$a \geq rb \iff a \geq \lceil rb \rceil, \quad ab \geq n \iff a \geq \left\lceil \frac{n}{b} \right\rceil.$$

Hence  $(a, b) \in W$  iff  $a \geq f(b)$ .

By definition  $a_w = \min_{(a,b) \in W} a$ ; therefore

$$a_w = \min_{b \in N} f(b).$$

Consequently for the fixed  $a_w$ ,

$$(a_w, b) \in W \iff f(b) = a_w.$$

Thus the set we are interested in is exactly the set of minimizers of  $f$ .

---

## 1. A useful auxiliary integer

Set

$$b_0 = \lceil n/a_w \rceil.$$

Then  $b_0 \geq n/a_w$ , so  $a_w b_0 \geq n$ .

Because  $b_w \in W$  we have  $b_w \geq n/a_w$ ; therefore

$$b_0 \leq b_w. \tag{1}$$

From  $a_w \geq r b_w$  we obtain  $b_w \leq a_w/r$ , hence

$$b_0 \leq \left\lfloor \frac{a_w}{r} \right\rfloor. \tag{2}$$

Together (1) and (2) give

$$b_0 \leq \left\lfloor \frac{a_w}{r} \right\rfloor. \tag{3}$$

The hypothesis  $a_w < r b_h$  yields

$$\left\lfloor \frac{a_w}{r} \right\rfloor \leq b_h - 1. \tag{4}$$

## 2. Lower bound for $b_0$

We prove that  $b_0 \geq b_h - 1$ .

Assume, to the contrary, that  $b_0 \leq b_h - 2$ .

Then  $b_0 + 1 \leq b_h - 1 < b_h$ .

Because  $a_w$  is the minimum of  $f$ , we have  $f(b_0) \geq a_w$ .

Now  $f(b_0) = \max \{ \lceil rb_0 \rceil, \lceil n/b_0 \rceil \}$ .

Since  $n/b_0 \leq a_w$ , we get  $\lceil n/b_0 \rceil \leq a_w$ ; therefore the inequality  $f(b_0) \geq a_w$  forces

$$\lceil rb_0 \rceil \geq a_w.$$

The last inequality implies  $rb_0 > a_w - 1$ . Adding  $r \geq 1$  gives

$$r(b_0 + 1) = rb_0 + r > a_w - 1 + r \geq a_w.$$

Let  $k = \lfloor r(b_0 + 1) \rfloor$ . Because  $r(b_0 + 1) > a_w$ , the integer  $a_w$  (which is  $< r(b_0 + 1)$ ) cannot exceed the greatest integer  $\leq r(b_0 + 1)$ ; hence  $a_w \leq k$ . Consequently

$$a_w \leq r(b_0 + 1).$$

Moreover  $a_w(b_0 + 1) = a_w b_0 + a_w \geq n + a_w > n$ , so  $a_w(b_0 + 1) \geq n$ .

Thus  $(a_w, b_0 + 1)$  satisfies both conditions of  $H$ ; i.e.  $(a_w, b_0 + 1) \in H$ .

But  $b_0 + 1 \leq b_h - 1 < b_h$ , contradicting the definition of  $b_h$  as the smallest second coordinate occurring in  $H$ .

Hence our assumption was false, and we must have

$$b_0 \geq b_h - 1. \tag{5}$$

### 3. Determining $b_0$ and $b_w$

From (3) and (4) we obtain  $b_0 \leq b_h - 1$ ; together with (5) this yields  $b_0 = b_h - 1$ . Therefore

$$\left\lceil \frac{n}{a_w} \right\rceil = b_h - 1. \tag{6}$$

Because of (1) and (6) we have  $b_w \geq b_h - 1$ ; together with (2) and (4) we get  $b_w \leq b_h - 1$ . Hence

$$b_w = b_h - 1. \tag{7}$$

Finally (7) and (4) give  $b_h - 1 \leq \lfloor a_w/r \rfloor$ , while (4) itself gives  $\lfloor a_w/r \rfloor \leq b_h - 1$ ; thus

$$\left\lfloor \frac{a_w}{r} \right\rfloor = b_h - 1. \tag{8}$$

### 4. Conclusion

For an integer  $b$  we have  $(a_w, b) \in W$  exactly when

$$\left\lceil \frac{n}{a_w} \right\rceil \leq b \leq \left\lfloor \frac{a_w}{r} \right\rfloor.$$

By (6) and (8) the left-hand side equals the right-hand side and both equal  $b_h - 1$ .

Because of (7) this common value is precisely  $b_w$ . Consequently no other integer  $b \neq b_w$  can satisfy the inequalities, i.e.

$$\left\lceil \frac{n}{a_w} \right\rceil = \lfloor r a_w \rfloor = b_w.$$

Hence there is no  $b \neq b_w$  with  $(a_w, b) \in W$ . ■

### Proof 3

Another attempt from DeepSeek.

#### Proof

Assume the fit-width solution  $(a, b)$  is strictly optimal, meaning its square side  $s_{fw} = w/a$  is larger than the side of every fit-height solution.

- Strict optimality forces  $r = w/h > 1$  (for  $r \leq 1$  the fit-height solution is at least as good).
- The algorithm finds the smallest integer  $a$  satisfying  $a \geq r \lceil n/a \rceil$ . Hence

$$a - 1 < r \lceil n/(a - 1) \rceil \quad \text{and} \quad a \geq r b, \quad b = \lceil n/a \rceil.$$

Because  $a$  is the *smallest* such integer, the gap  $a - r b$  is strictly less than 1.

- For  $r > 1$ , we have

$$\frac{a}{r} - b = \frac{a - r b}{r} < \frac{1}{r} < 1 \quad \implies \quad \frac{a}{r} < b + 1.$$

At the same time  $a/r \geq b$ . Thus  $\lfloor a/r \rfloor = b$ .

- A different row count  $b'$  would still be a valid fit-width solution with the same  $a$  only if  $b' \in [\lceil n/a \rceil, \lfloor a/r \rfloor] = [b, b]$ . Hence no  $b' > b$  exists.

#### Why the converse (alternative grids) cannot happen under strict optimality

If a  $b' > b$  existed with  $b' \leq a/r$ , then  $b + 1 \leq \lfloor a/r \rfloor$ . One can show this forces the existence of a fit-height solution with side  $\geq w/a$ , contradicting strict optimality. In practice, however, the simpler box argument above already forbids the existence of any such  $b'$  when  $r > 1$ .



# Thoughts on the LLMs' performance

---

Using the prompt in a fresh conversation:

```
let r in [1, inf).
let n natural non-zero.
let W = { (a,b) in N^2 | n <= ab and a >= rb }.
let H = { (a,b) in N^2 | n <= ab and rb >= a }.
let (a_w, b_w) = \argmin_{(a,b) in W} a.
let (a_h, b_h) = \argmin_{(a,b) in H} b.
assume a_w < rb_h.
prove that there exists no other b ≠ b_w such that (a_w, b) is in W.
```

All current frontier models managed to prove the statement correctly (don't click on links if you didn't solve yet):

- [GPT 5.5 Pro](#) had a correct, concise, minimal proof, light on algebra and easy to follow;
- [Claude Opus 4.7](#) proved the same result as GPT 5.5 Pro but by more complex, slightly less readable means; a nice touch was providing the geometrical intuition for the proof, just as it is described in the main document;
- [Gemini 3.1 Pro](#) had an unintuitive roundabout excessively long but correct proof;
- [DeepSeek v4](#) had a different yet equally elegant approach as GPT & Claude but the writing itself was hard to follow and full of redundant statements.

DeepSeek's two failed proofs are part of a much greater conversation and used a more ambiguous prompt. It had a lot more context: the algorithm's code, its complexity analysis, the graph representation, the terminology. This seems to have hurt its performance.

The prompt itself was built such that QED-Nano managed to output a response. If any ambiguities were present it would "overthink" them and wouldn't make meaningful progress beyond continuously reanalyzing the problem statement. Running QED-Nano with smaller context (8k is all I could run locally), resulted in it forgetting about the initial problem and proving unrelated statements (e.g. that there are  $a, b, c \in \mathbb{N}$  such that  $a^3 + b^3 = c^3$ , lol). I paid [vast.ai](#) 5\$ to run it using the biggest supported context, which required ~46GB of VRAM.

That being said, [QED-Nano did prove](#) the statement with a very minor push in a second prompt. The approach is equivalent to DeepSeek's, with even more verbose writing. The cause for verbosity is not redundancy but derivations of intermediary results through alternative more inefficient methods.

I've also tried ChatGPT 5.5 Thinking Mini with the same prompt. After a failed proof which didn't use the  $a_w < rb_h$  assumption and some subsequent corrective prompts, it concluded that:

As it stands, I cannot honestly provide a correct proof because the crucial implication needed for the contradiction is not derivable from the assumptions you gave.

Such hubris!