



CSCE 222

Discrete Structures

Functions

Dr. Tim McGuire

Grateful acknowledgement to Professor Bart Selman, Cornell University, and Prof. Johnnie Baker, Kent State, for some of the material upon which these notes are adapted.

Based on Chapter 2 of Rosen
Discrete Mathematics and its Applications

Functions

A **function** f from a set A to a set B is an **assignment** of **exactly one** element of B to **each** element of A .

We write

$$f(a) = b$$

if b is the unique element of B assigned by the function f to the element a of A .

If f is a function from A to B , we write

$$f: A \rightarrow B$$

(note: Here, “ \rightarrow ” has nothing to do with if... then)

Note: Functions are also called **mappings** or **transformations**.

Functions

If $f:A \rightarrow B$, we say that A is the **domain** of f and B is the **codomain** of f .


If $f(a) = b$, we say that b is the **image** of a and a is the **pre-image** of b .

The **range** of $f:A \rightarrow B$ is the set of all images of **all** elements of A .

We say that $f:A \rightarrow B$ **maps** A to B .

Functions

More generally:



A - Domain of f

B- Co-Domain of f

For example: $f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = -(1/2)x - 1/2$

domain

co-domain



Functions

a collection of points!

More formally: a function $f: A \rightarrow B$ is a subset of $A \times B$ where
 $\forall a \in A, \exists! b \in B$ and $\langle a, b \rangle \in f$.

a point!

Remember, this means “there exists a unique ...”




Why not?

Functions

$A = \{\text{Michael, Toby, John, Chris, Brad}\}$

$B = \{\text{Kathy, Carla, Mary}\}$


Let $f: A \rightarrow B$ be defined as $f(a) = \text{mother}(a)$.



Every $b \in B$ has
at most 1
preimage.

Functions - injection

A function $f: A \rightarrow B$ is **one-to-one** (**injective, an injection**) if
 $\forall a, b, c, (f(a) = b \wedge f(c) = b) \rightarrow a = c$




In other words: f is **one-to-one** if and only if it does not map two distinct elements of A onto the same element of B .

Every $b \in B$ has
at least 1
preimage.

Functions - surjection


A function $f: A \rightarrow B$ is **onto** (**surjective, a surjection**) if $\forall b \in B, \exists a \in A f(a) = b$

In other words, f is **onto** if and only if its range is its entire codomain.



Functions – one-to-one-correspondence or bijection


A function $f: A \rightarrow B$ is **bijective** if it is **one-to-one** and **onto**.



Functions: inverse function

Definition:

Given f , a one-to-one correspondence from set A to set B , the **inverse function of f** is the function that assigns to an element b belonging to B the unique element a in A such that $f(a)=b$. The inverse function is denoted f^{-1} . $f^{-1}(b)=a$, when $f(a)=b$.




Functions - examples

Suppose $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $f(x) = x^2$.

Is f one-to-one? **yes**

Is f onto? **yes**

Is f bijective? **yes**



This function is invertible.

Functions - examples

Suppose $f: \mathbb{R} \rightarrow \mathbb{R}^+$, $f(x) = x^2$.

Is f one-to-one?


no

Is f onto?

yes


Is f bijective?

no




This function is not invertible.


(a) One-to-one,
not onto




(b) Onto,
not one-to-one




(c) One-to-one,
and onto



(d) Neither one-to-one
nor onto




(e) Not a function



Functions - composition

Let $f: A \rightarrow B$, and $g: B \rightarrow C$ be functions. Then the **composition** of f and g is:

$$(f \circ g)(x) = f(g(x))$$



Note: $(f \circ g)$ cannot be defined unless the range of g is a subset of the domain of f .

Example:

Let $f(x) = 2x + 3$; $g(x) = 3x + 2$;

$$(f \circ g)(x) = f(3x + 2) = 2(3x + 2) + 3 = 6x + 7.$$

$$(g \circ f)(x) = g(2x + 3) = 3(2x + 3) + 2 = 6x + 11.$$

As this example shows, $(f \circ g)$ and $(g \circ f)$ are not necessarily equal – i.e, the **composition of functions is not commutative**.

Composition

Composition of a function and its inverse:

$$(f^{-1} \circ f)(a) = f^{-1}(f(a)) = f^{-1}(b) = a.$$

$$(f \circ f^{-1})(b) = f(f^{-1}(b)) = f(a) = b$$

$$(f^{-1} \circ f)(x) = f^{-1}(f(x)) = x$$

The composition of a function and its inverse is the **identity function** $I(x) = x$.

Graphs

The **graph** of a function $f:A\rightarrow B$ is the set of ordered pairs $\{(a, b) \mid a\in A \text{ and } f(a) = b\}$.

The graph is a subset of $A\times B$ that can be used to visualize f in a two-dimensional coordinate system.

Some important functions

Absolute value:

Domain \mathbb{R} ; Co-Domain = $\{0\} \cup \mathbb{R}^+$

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Ex: $|-3| = 3$; $|3| = 3$

Floor function (or greatest integer function):

Domain = \mathbb{R} ; Co-Domain = \mathbb{Z}

$\lfloor x \rfloor$ = largest integer not greater than x

Ex: $\lfloor 3.2 \rfloor = 3$; $\lfloor -2.5 \rfloor = -3$

Some important functions

Ceiling function:

Domain = \mathbb{R} ;

Co-Domain = \mathbb{Z}

$\lceil x \rceil$ = smallest integer greater than or equal to x

Ex: $\lceil 3.2 \rceil = 4$; $\lceil -2.5 \rceil = -2$

Table 1, Section 2-3

TABLE 1 Useful Properties of the Floor and Ceiling Functions.

(n is an integer)

$$(1a) \quad \lfloor x \rfloor = n \text{ if and only if } n \leq x < n + 1$$

$$(1b) \quad \lceil x \rceil = n \text{ if and only if } n - 1 < x \leq n$$

$$(1c) \quad \lfloor x \rfloor = n \text{ if and only if } x - 1 < n \leq x$$

$$(1d) \quad \lceil x \rceil = n \text{ if and only if } x \leq n < x + 1$$

$$(2) \quad x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$$

$$(3a) \quad \lfloor -x \rfloor = -\lceil x \rceil$$

$$(3b) \quad \lceil -x \rceil = -\lfloor x \rfloor$$

$$(4a) \quad \lfloor x + n \rfloor = \lfloor x \rfloor + n$$

$$(4b) \quad \lceil x + n \rceil = \lceil x \rceil + n$$

Some important functions

Factorial function: Domain = \mathbf{N} , Range = \mathbf{N}^+

$$n! = n (n-1)(n-2) \dots, 3 \times 2 \times 1$$

$$\text{Ex: } 5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$$

Note: $0! = 1$ by convention.

Some important functions

Mod (or remainder):

Domain = $\mathbf{N} \times \mathbf{N}^+ = \{(m,n) \mid m \in \mathbf{N}, n \in \mathbf{N}^+ \}$

Co-domain Range = \mathbf{N}

$$m \bmod n = m - \lfloor m/n \rfloor n$$

Ex: $8 \bmod 3 = 8 - \lfloor 8/3 \rfloor 3 = 2$

$$57 \bmod 12 = 9;$$

In C++ this is computed using the % operator. That is, to compute $m \bmod n$, we write **`m % n`**

Note: This function computes the remainder when m is divided by n .

The name of this function is an abbreviation of m modulo n , where modulus means with respect to a modulus (size) of n , which is defined to be the remainder when m is divided by n . Note also that this function is an example in which the domain of the function is a 2-tuple.

Some important functions:

Exponential Function

Exponential function:

Domain = $\mathbb{R}^+ \times \mathbb{R} = \{(a, x) \mid a \in \mathbb{R}^+, x \in \mathbb{R}\}$

Co-domain Range = \mathbb{R}^+

$$f(x) = a^x$$

Note: a is a **positive** constant; x varies.

Ex: $f(n) = a^n = a \times a \times \dots \times a$ (n times)

How do we define $f(x)$ if x is not a positive integer?

Some important functions:

Exponential function

Exponential function:

How do we define $f(x)$ if x is not a positive integer?

Important properties of exponential functions:

$$(1) a^{(x+y)} = a^x a^y; \quad (2) a^1 = a; \quad (3) a^0 = 1$$

See:

$$a^2 = a^{1+1} = a^1 a^1 = a \times a;$$
$$a^3 = a^{2+1} = a^2 a^1 = a \times a \times a;$$

...

$$a^n = a \times \cdots \times a \quad (n \text{ times})$$

We get: $a = a^1 = a^{1+0} = a \times a^0$ therefore $a^0 = 1$

$$1 = a^0 = a^{b+(-b)} = a^b \times a^{-b} \text{ therefore } a^{-b} = 1/a^b$$

$$a = a^1 = a^{\frac{1}{2}+\frac{1}{2}} = a^{\frac{1}{2}} \times a^{\frac{1}{2}} = (a^{\frac{1}{2}})^2 \text{ therefore } a^{\frac{1}{2}} = \sqrt{a}$$

By similar arguments:

$$a^{\frac{1}{k}} = \sqrt[k]{a}$$

$$a^{mx} = a^x \times \cdots a^x \text{ (} m \text{ times)} = (a^x)^m, \text{ therefore } a^{\frac{m}{n}} = (a^{\frac{1}{n}})^m = (\sqrt[n]{a})^m$$

Note: This determines a^x for all x rational. x is irrational by continuity (we'll skip “details”).

Some important functions:

Logarithm Function

$$\log_a x = y \text{ means } a^y = x$$

A one-word definition for
logarithm is *exponent*

Logarithm base a:

$$\text{Domain} = \mathbb{R}^+ \times \mathbb{R} = \{(a, x) \mid a \in \mathbb{R}^+, a > 1, x \in \mathbb{R}\}$$

$$\text{Co-domain Range} = \mathbb{R}$$

$$y = \log_a(x) \Leftrightarrow a^y = x$$

$$\text{Ex: } \log_2(8) = 3; \quad \log_2(16) = 4; \quad 3 < \log_2(15) < 4.$$

Some important functions:

Logarithm Function

$$\log_a x = y \text{ means } a^y = x$$

A one-word definition for *logarithm* is *exponent*

Key properties of the log function (they follow from those for exponential):

1. $\log_a (1)=0$ (because $a^0 =1$)
2. $\log_a (a)=1$ (because $a^1 =a$)
3. $\log_a (xy) = \log_a (x) + \log_a (y)$ (similar arguments)
4. $\log_a (x^r) = r \log_a (x)$
5. $\log_a (1/x) = - \log_a (x)$ (note $1/x = x^{-1}$)
6. $\log_b (x) = \log_a (x) / \log_a (b)$

Logarithm Functions

Examples:

$$\log_2 (1/4) = -\log_2 (4) = -2.$$


$$\log_2 (-4) \text{ undefined}$$

$$\begin{aligned}\log_2 (2^{10} 3^5) &= \log_2 (2^{10}) + \log_2 (3^5) = 10 \log_2 (2) + 5 \log_2 (3) \\ &= 10 + 5 \log_2 (3)\end{aligned}$$

Limit Properties of Log Function

$$\lim_{x \rightarrow \infty} \log(x) = \infty$$

$$\lim_{x \rightarrow \infty} \frac{\log(x)}{x} = 0$$



As x gets large, $\log(x)$ grows without bound.

But x grows **MUCH** faster than $\log(x)$...more soon on growth rates.

Some important functions: Polynomials

Polynomial function:

Domain = usually \mathbb{R}

Co-domain Range = usually \mathbb{R}

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x^1 + a_0$$

n , a nonnegative integer is the degree of the polynomial; $a_n \neq 0$ (so that the term $a_n x^n$ actually appears)

$(a_n, a_{n-1}, \dots, a_1, a_0)$ are the coefficients of the polynomial.

Ex:

$$y = P_1(x) = a_1 x^1 + a_0 \text{ linear function}$$

$$y = P_2(x) = a_2 x^2 + a_1 x^1 + a_0 \text{ quadratic polynomial or function}$$

Exponentials grow MUCH faster than polynomials:

$$\lim_{x \rightarrow \infty} \frac{a_0 + \cdots + a_k x^k}{b^x} = 0 \text{ if } b > 1$$

We'll talk more about growth in Chapter 3