

Van der Waerden's Theorem — The Root of Arithmetic Ramsey Theory

Where Ramsey finds cliques in graphs and Erdős–Szekeres finds convex polygons in point sets, van der Waerden finds regular patterns in colorings of numbers. Oldest Ramsey-type theorem 1927, archetypal statement that complete disorder impossible.

Van der Waerden 1927: For any finite r colors and desired length k , exists minimum $W(r, k)$ such that for any $N \geq W(r, k)$, every r -coloring of $\{1, \dots, N\}$ contains monochromatic arithmetic progression length k .

Paint integers red/blue however cleverly to break equally-spaced patterns. Paint long enough, you lose — long one-color progression forced.

Key Concepts — Van der Waerden

Coloring: $c : \{1, \dots, N\} \rightarrow \{1, \dots, r\}$.

Function assigning each integer a color label. For $r = 2$, Red/Blue.

Think painting houses $1..N$ red/blue. No rule, any assignment allowed — you're adversary trying to avoid pattern.

AP: Arithmetic progression $a, a + d, \dots, a + (k - 1)d$.

- a start, $d > 0$ common difference, k length.
- $3, 6, 9 = a = 3, d = 3, k = 3$.
- $4, 11, 18, 25 = a = 4, d = 7, k = 4$.
- Difference d not required to be small — can be large jump.

Equal spacing is order: points line up arithmetic, not geometric.

Monochromatic: All k terms same color.

$$c(a) = c(a + d) = \dots = c(a + (k - 1)d).$$

You tried to color to break APs, but this k -set slipped through one color.

Van der Waerden number $W(r, k)$: Threshold where avoidance ends.

Least N such that every r -coloring of $\{1, \dots, N\}$ contains monochromatic k -AP.

Two sides:

- $W - 1$ avoidable: exists at least one coloring of $\{1, \dots, W - 1\}$ with *no* monochromatic k -AP — maximal disorder example. For $W(2, 3) = 9$, that's $RRBBRRBB$ on $1..8$.
- W unavoidable: no coloring of $\{1, \dots, W\}$ avoids. Every attempt forces k -AP. For $W(2, 3) = 9$, every red/blue of $1..9$ yields red $a, a + d, a + 2d$ or blue.

Party version: $W(r, k)$ = number of houses on street you must paint with r colors before k equally-spaced houses of one color become unavoidable, no matter how clever you paint.

It's Ramsey number for integer line — host structure is $\{1, \dots, N\}$ with AP hyperedges, colors try to avoid monochromatic hyperedge, W is when host large enough that avoidance impossible.

Like $R(3, 3) = 6$ is threshold for monochromatic triangle in complete graph, $W(2, 3) = 9$ is threshold for monochromatic 3-AP on line, $N(4) = 5$ threshold for convex quadrilateral in point set, $HJ(n, c)$ threshold for combinatorial line in cube.

All measure same thing: disorder budget.

Why $W(2, 3) = 9$ — Only Case You Can See

This is van der Waerden's $R(3, 3) = 6$ — only number you can prove by hand, and it contains whole general proof idea: local constraints propagate until collision.

Goal: 2 colors, avoid monochromatic 3-term $a, a + d, a + 2d$.

Forcing rule: If a, b same color, then $2b - a$ must be opposite color, else $a, b, 2b - a$ is monochromatic AP with difference $d = b - a$.

Part A: $W(2, 3) > 8$ — 8 Can Avoid

$$1 : R, 2 : R, 3 : B, 4 : B, 5 : R, 6 : R, 7 : B, 8 : B = RRBBRRBB$$

Check reds $\{1, 2, 5, 6\}$:

- $d = 1$: 1, 2, 3 needs 3 — 3 is B, safe. 5, 6, 7 needs 7 — B safe.
- $d = 2$: 1, 3, 5 needs 3 — B safe. 2, 4, 6 needs 4 — B safe.
- $d = 3$: 2, 5, 8 needs 8 — B safe. 1, 4, 7 needs 7 but 4 is B.
- $d = 4$: 1, 5, 9 out of range.

No red $a, a + d, a + 2d$ fully red.

Blues $\{3, 4, 7, 8\}$ symmetric: 3, 5, 7 needs 5 R safe, 4, 6, 8 needs 6 R safe.

So 8 avoids. Pattern = blocks of 2: $RRBBRRBB$ — you avoid 3-AP by never making run of 3, and offsetting blocks so $a, a + d, a + 2d$ always hits opposite block.

Thus $W(2, 3)$ at least 9.

Part B: $W(2, 3) \leq 9$ — 9 Forces

We need show *every* 2-coloring of 1..9 forces monochromatic 3-AP, not just $RRBBRRBB$ extended.

Brute casework but organized by forcing:

Assume coloring of 1..8 avoids 3-AP. WLOG $1 = R$ by symmetry.

Consider middle point 5 — it participates in many APs: 1, 5, 9 ; 2, 5, 8 ; 3, 5, 7 ; 4, 5, 6.

If we try to 2-color 1..8 without 3-AP, constraints propagate:

- $1 : R$ and $2 : R \rightarrow 3$ must be B , else 1, 2, 3 red.
- $3 : B$ and 4? If $4 = R$, then 2, 3, 4? 2 R, 3 B no. Let's systematic search — classic proof enumerates 8 avoiding colorings of length 8 — there are essentially 2 types up to swap colors and reflection: $RRBBRRBB$ and $RBRBR...$ variations?

Actually complete enumeration shows only 4 colorings of 1..8 avoid 3-AP up to symmetry, all end with ...BB.

Simplified argument for $RRBBRRBB$ case you wrote:

Extend to 9:

- If $9 = R$: then 1, 5, 9 with $d = 4$ is 1 : R , 5 : R , 9 : R — red AP.
- If $9 = B$: then 7, 8, 9 with $d = 1$ is 7 : B , 8 : B , 9 : B — blue AP.

So this maximal coloring dies at 9.

Why do all other 8-colorings die? Same collision: any avoiding coloring of 1..8 must have last two same color — say 7 : B , 8 : B — otherwise you'd have created AP earlier. Then $9 = B$ completes 7, 8, 9 blue. If you try $9 = R$ to avoid that, you complete long difference AP 1, 5, 9 or 3, 6, 9 or 2, 5, 8 etc red.

You run out of room to satisfy all $2b - a \neq \text{color}(a)$ constraints simultaneously.

Formal: 8 points impose constraints like $c(9) \neq c(1)$ if $c(1) = c(5)$, and $c(9) \neq c(7)$ if $c(7) = c(8)$, etc. At $N = 9$, constraint graph has odd cycle — unsatisfiable.

Hence no coloring of 1..9 avoids monochromatic 3-AP. $W(2, 3) \leq 9$.

Together > 8 and $\leq 9 \Rightarrow W(2, 3) = 9$.

Try to avoid red $a, a + d, a + 2d$ and blue $a, a + d, a + 2d$ by alternating blocks $RRBB$. You can juggle 8 houses. 9th house forces you to close a monochromatic equally-spaced triple — either you close 7, 8, 9 same block, or you close long jump 1, 5, 9.

That's local-to-global forcing — same as $N(4) = 5$ hull case analysis and $R(3, 3) = 6$ degree case analysis: enumerate small types of disorder, show each forces order at next size.

How Fast Do They Grow? Faster Than Ramsey Graph Numbers

$W(r, k)$ exists — van der Waerden proved finiteness — but values explode much faster than $R(n, m)$.

W	VALUE	MEANING
$W(2, 3)$	9	1..8 avoids 3-AP, 1..9 cannot. Hand.
$W(3, 3)$	27	3 colors, 3-AP. First non-trivial with >2 colors
$W(2, 4)$	35	Chvátal 1970 backtracking search — 2^{34} naive
$W(2, 5)$	178	2^{177} colorings ruled out conceptually
$W(2, 6)$	1,132	Kouril & Paul 2008 SAT solvers months CPU. You <i>can</i> 2-color 1..1131 with no 6-AP, but 1132 forces
$W(2, 7)$	3,703	Current SAT best 2008-2017
$W(2, 10)$	Unknown	lower bound >10k, upper bound tower

Why Explosive

$R(4, 4) = 18$ means among 18 people, monochromatic K_4 forced. Host graph has $\binom{18}{2} = 153$ edges, 2^{153} colorings — huge but search feasible with symmetry.

$W(2, 6) = 1,132$ means among 1,132 integers, monochromatic 6-AP forced. Host has 2^{1132} colorings — astronomically larger search space, cannot brute force. Plus constraints long-range: AP with $d = 500$ connects distant positions, constraint graph not local.

Proofs for $W(2, 6)$ used DPLL SAT solvers with clause learning:

- Encode each $a \in [1, N]$ boolean variable $x_a = \text{Red/Blue}$.
- For each 6-AP $a, a + d, \dots, a + 5d$ in $[[1, N]]$, add clauses $\neg(x_a \wedge \dots \wedge x_{a+5d})$ and $\neg(\neg x_a \wedge \dots)$ — forbid all-red and all-blue.
- SAT solver tries to satisfy. If UNSAT, $N \geq W$. If SAT, coloring found, $N < W$.

For $N = 1,131$, SAT finds model — explicit coloring avoiding 6-AP, highly non-repetitive, not simple $RRBB$ pattern, looks pseudo-random.

For $N = 1,132$, solver proves UNSAT after exploring search tree with backtracking, symmetry breaking ($1 = R$ wlog), and learning.

Months CPU 2008, today hours with modern CDCL solvers — but still non-trivial.

Comparison to Ramsey Numbers

- $R(3, 3) = 6$, $R(4, 4) = 18$, $R(5, 5)$ unknown 43-48.
- $W(2, 3) = 9$, $W(2, 4) = 35$, $W(2, 5) = 178$, $W(2, 6) = 1132$.

Growth of $W(2, k)$ already exponential 2^k lower bound (Berlekamp $p \cdot 2^p$ algebraic using finite fields), upper bound tower of exponentials — gap huge.

$W(3, 3) = 27$ vs $R(3, 3) = 6$ shows extra color costs less than longer progression.

$W(2, 6) = 1132$ instructive because it shows disorder budget: you can be clever and avoid 6 equally spaced same color for 1,131 houses, using complicated pattern that is not simple periodic — looks random — but at 1,132 cleverness exhausted, arithmetic order forced.

Same as Erdős–Szekeres: you can avoid convex 6-gon with 16 points using flat double construction, but 17 forces it.

Difference: $N(6) = 17$ small, $W(2, 6) = 1132$ large — integer line needs much larger host to force AP than plane needs to force convex polygon, because AP is more rigid condition than convexity.

That's why van der Waerden numbers are among largest small Ramsey-type numbers known exactly — they measure how long you can delay arithmetic order.

History and Bounds: From Ackermann to Fields Medal

Van der Waerden proved 1927 answering conjecture of Baudet and Schur: does $W(r, k)$ finite? Yes.

His original proof double induction on r and k :

- Induct on k , with inner induction building blocks of size $W(r', k - 1)$ etc.
- Each step replaces N by r^N — exponentiation.

- After k steps, tower of height k — Ackermann-type, not primitive recursive.

For 60 years bound stayed tower of exponentials whose height grows with k — essentially uncomputable for large k . Upper bound far from true value.

Modern View 1: Hales-Jewett $\Rightarrow$ van der Waerden

Today we don't teach original proof. We derive van der Waerden from Hales-Jewett — shift from numbers to words.

Coding:

Alphabet $A = \{0, \dots, k-1\}$. Word $w \in A^H$ length H .

Pick large base $M > kH$ to avoid carries. Map word to number:

$$N(w) = \sum_{i=0}^{H-1} w_i M^i$$

This is injective — different words different numbers because no carry.

Variable word $v(x)$ — word with x placeholders, e.g. $v(x) = 1x0x2$ with x at positions $i = 1, 3$.

Then $N(v(a)) = C + a \cdot d$ where $C = \sum_{\text{const positions}} c_i M^i$, $d = \sum_{\text{x positions}} M^i$ fixed independent of a .

As a runs $0, \dots, k-1$, $N(v(a))$ runs k -term AP: $C, C + d, C + 2d, \dots, C + (k-1)d$.

HJ $\Rightarrow$ vdW: Hales-Jewett says for $H = HJ(k, c)$ large enough, any c -coloring of A^H yields monochromatic combinatorial line — set $\{v(0), \dots, v(k-1)\}$ for some variable word v all same color.

Color integers by coloring words via N : color of $N(w) = \text{color of } w$. Then monochromatic line maps to monochromatic k -AP $C + ad$.

So

Order in words forces order in numbers.

Van der Waerden becomes coding of HJ — purely combinatorial operation substitution, no arithmetic needed. Shows AP forcing is special case of variable substitution forcing.

This also gives new upper bound $W(r, k) \leq M^{HJ(k, r)}$ — still tower, because HJ itself tower, but conceptually clean.

Modern View 2: Gowers Quantitative Bound

1987 Shelah gave primitive recursive bound for HJ, thus for vdW — first tower not Ackermann.

2001 Gowers gave first *reasonable* bound using Fourier analysis and Szemerédi theorem:

$$W(2, k) \leq 2^{2^{2^{k+9}}}$$

Tower of 5 exponentials.

Still astronomically huge — for $k = 6$, bound is $2^{2^{2^{15}}}$ with $2^{15} = 32768$, tower $\approx 10^{10^{9860}}$ vs true $W(2, 6) = 1132$. But it was first bound that is elementary recursive — fixed height tower independent of k growth inside top.

Idea: Gowers uniformity norms U^d — measure how "pseudo-random" set is vs structured. Decompose dense set $A \subset [1, N]$ into structured + small + uniform. If uniform, many k -APs by counting lemma; if structured, correlate with polynomial phase, iterate.

For this work connecting Fourier analysis to Ramsey theory, Gowers received Fields Medal 1998.

Lower bounds: Exponential, not tower.

Berlekamp 1968 algebraic construction: For prime p , using quadratic residues in $\mathbb{F}_p$, construct coloring of $p \cdot 2^p$ with no $(p + 1)$ -AP? More precisely $W(2, p + 1) > p \cdot 2^p$.

Take finite field $\mathbb{F}_{2^n}$, trace map, color by quadratic character — gives long AP-free coloring length $\approx k \cdot 2^k$.

So $p2^p \leq W(2, p + 1) \leq \text{tower}(k)$.

True growth between exponential and tower — one of largest gaps in combinatorics. We don't know if closer to bottom or top — most conjecture closer to exponential, but no proof.

Timeline

- 1927 van der Waerden double induction — Ackermann tower bound.
- 1968 Berlekamp exponential lower bound.
- 1963 Hales-Jewett — new abstraction, implies vdW.
- 1987 Shelah primitive recursive bound for HJ.
- 2001 Gowers 5-exponential bound — Fields Medal work.
- 2008 $W(2, 6) = 1132$ SAT.
- 2017-2024 Polymath improvements lower order exponents to $O(k^2)$ in top.

Status: $W(r, k)$ finite, exact values up to $k = 7$, growth exponential $\leq W \leq \text{tower}$ — same pattern as $N(n) = 2^{n+o(n)}$ vs $2^{n-2} + 1$ and HJ tower vs primitive recursive — order forced, but how soon remains mystery.

Why It Matters — Partition Regularity

Van der Waerden first formalized what all Ramsey-type theorems formalize:

Partition regularity: You cannot destroy all arithmetic structure by finitely partitioning $\mathbb{N}$. One cell of the partition still contains arbitrarily long APs.

Try to paint integers red/blue to break equally-spaced patterns — you can delay, up to $W(r, k) - 1$, but not forever. Some color retains structure.

This idea launched three generations — each stronger, each asking "how sparse can host be and still force AP?"

1. Szemerédi 1975 — From Partition to Density

van der Waerden: you color *all* of $\mathbb{N}$ with r colors — whole line partitioned — one color forces k -AP.

Szemerédi: you don't need to color everything. If $A \subset \mathbb{N}$ has positive upper density

$$\bar{d}(A) = \limsup \frac{|A \cap [1, N]|}{N} \geq \delta > 0$$

— e.g., contains at least 1% of numbers up to large N — then A alone contains arbitrarily long APs.

No partition, no other colors. Thickness alone forces.

This implies van der Waerden: r -color $\mathbb{N}$, one color has density $\geq 1/r > 0$, apply Szemerédi, get k -AP.

Strictly stronger — Szemerédi applies to sets that are not complements of partition, like set of numbers with first digit 1 — density ≈ 0.1 but not from finite coloring.

Proof created regularity lemma, launched additive combinatorics.

2. Green-Tao 2004 — Density Zero Still Forces

Primes $P = \{2, 3, 5, 7, 11, \dots\}$ have density 0 — $|P \cap [1, N]| \sim N / \log N$, so $|P|/N \rightarrow 0$. Szemerédi does NOT apply.

Green-Tao: primes contain arbitrarily long APs anyway.

Why? Primes are not arbitrary sparse set — they are *dense inside pseudorandom host*.

Take almost-primes — numbers with no small prime factors, called W -tricked primes / Selberg sieve majorant ν . ν is pseudorandom — Fourier uniform — and has positive density. Primes sit inside ν with relative density > 0 — weighted primes behave like dense subset of random-like set.

Relative Szemerédi: dense subset of pseudorandom set forces k -AP.

Thus k -AP of primes exists for every k . Longest known as of 2024: 27-term AP of primes, start $224584605939537911 + 81292139 \cdot 23\# \cdot n$ — huge.

Primes have 26-term APs known explicitly, but Green-Tao says arbitrarily long exist, not constructed.

Ramsey Language Unification

- $R(3, 3) = 6$: host = complete graph K_6 , coloring edges 2 colors, forced structure = monochromatic triangle.
- $N(4) = 5$: host = 5 points in plane, forced = convex quadrilateral.
- $N(n) = 2^{n-2} + 1$: host = $2^{n-2} + 1$ points, forced = convex n -gon.
- $HJ(n, c)$: host = n -dimensional cube $[n]^H$, coloring words c colors, forced = monochromatic combinatorial line.
- $W(r, k)$: host = interval $\mathbb{Z}$, coloring integers r colors, forced = monochromatic k -AP. [1][W]
- Szemerédi: host = dense set $A \subset [1, N]$, forced = k -AP inside A .
- Green-Tao: host = primes weighted by sieve, forced = k -AP of primes.

Same philosophy:

For any finite coloring / dense subset of large enough structure, monochromatic / internal order unavoidable.

Disorder Budget View

Each theorem defines budget parameter and threshold where budget exhausted:

- Gomory: budget $W - B$ color imbalance — if $\neq 0$, no tiling.
- Egregium: budget K curvature difference — if $\neq 0$, no isometry.

- Ramsey R : budget $N = R - 1$ vertices — can avoid monochrome clique, at R cannot.
- Erdős–Szekeres $N(n)$: budget 2^{n-2} points — can avoid convex n -gon, at $2^{n-2} + 1$ cannot — conjectured.
- Hales-Jewett HJ : budget $H = HJ - 1$ dimension — can avoid monochromatic line, at HJ cannot.
- van der Waerden $W(r, k)$: budget $W - 1$ length — can avoid k -AP, at W cannot.

All say: **Local disorder can exist for a while, but global order forced by size alone in right parameter.**

And that's why van der Waerden matters — first time arithmetic progression shown partition regular, seed for 100 years of additive combinatorics, from Szemerédi to Green-Tao to 2023 breakthrough on Roth's theorem. Order hides in numbers, not just graphs and point sets.

Szemerédi's Theorem — Density Forces Progression

Szemerédi 1975, resolving Erdős–Turán 1936: Any subset of $\mathbb{N}$ with positive upper density contains arbitrarily long arithmetic progressions.

This is van der Waerden on steroids.

Core Concepts — Szemerédi's Theorem

Density: $A \subset \mathbb{N}$ has positive upper density if

$$\bar{d}(A) = \limsup_{N \rightarrow \infty} \frac{|A \cap \{1, \dots, N\}|}{N} > 0$$

Meaning A contains fixed percentage of numbers up to large N infinitely often.

- Evens: $|E \cap [1, N]| \approx N/2 \rightarrow$ density $1/2$.
- Multiples of 3: $N/3 \rightarrow 1/3$.
- Squares: $|Squares \cap [1, N]| \approx \sqrt{N} \rightarrow \sqrt{N}/N = 1/\sqrt{N} \rightarrow 0 \rightarrow$ density 0.
- Primes: $N/\log N \rightarrow$ density 0.

$\limsup$ not $\lim$ — set may not have limiting frequency, but if it infinitely often reaches δ fraction, positive upper density.

Positive density = not too sparse. You keep δ fraction infinitely often.

AP: $a, a + d, a + 2d, \dots, a + (k - 1)d$ constant step $d > 0$.

k -term arithmetic progression — k equally spaced numbers.

- 3, 5, 7 is $a = 3, d = 2, k = 3$.
- 10, 20, 30, 40 is $a = 10, d = 10, k = 4$.
- d can be huge — not required small.

Structure to avoid if you want disorder.

Arbitrarily long: For every k , exists k -term AP inside A . Not same AP for all k .

For $k = 3$ some triple $a, a + d, a + 2d \in A$ — maybe $a = 100, d = 7$.

For $k = 4$ maybe different quadruple $a' = 1000, d' = 50$.

For $k = 100$ maybe a'' enormous.

Theorem does NOT say infinite AP inside A — set A finite blocks? Actually infinite AP would be $a, a + d, a + 2d, \dots$ forever — density would be > 0 but infinite AP would imply A contains infinite arithmetic progression, stronger than Szemerédi claims. Szemerédi claims for each finite k finite progression, not one infinite progression.

If set is not too sparse — contains $\delta > 0$ fraction of integers infinitely often — you cannot avoid k -term progression, no matter how you try to hide gaps.

Try to construct A density $1/2$ with no 3-AP: choose numbers avoiding $a, a + d, a + 2d$. Greedy fails — eventually pigeonhole forces. Szemerédi says impossible for any k .

This is why stronger than van der Waerden: van der Waerden needs coloring of all $\mathbb{N}$ — whole line partitioned — pigeonhole guarantees one color has density $\geq 1/r$. Szemerédi needs only one set thick enough — no partition of complement needed.

And why Green-Tao needed new idea: primes density 0, so Szemerédi does NOT force AP in primes — primes too sparse. Need relative density inside pseudorandom host.

Budget view: Density δ is disorder budget. If $\delta > 0$, you can avoid k -AP for a while up to some $N(\delta, k)$, but eventually AP forced. Threshold $N(\delta, k)$ grows as $\delta \rightarrow 0$ — sparser you are, longer you can avoid.

Exactly same as $W(r, k)$ budget $W - 1$, $N(n)$ budget 2^{n-2} , HJ budget dimension — local disorder exists, global order forced by size alone.

Significance — Why Szemerédi Matters

Foundational pillar of additive combinatorics

Before 1975, additive number theory was collection of tricks — van der Waerden, Roth $k = 3$, etc. No systematic theory of *structure* inside dense sets.

Szemerédi proved: density alone forces arbitrary additive structure — k -AP for every k .

This made additive combinatorics field: study of what additive patterns dense sets must contain, and what tools detect them — Fourier analysis, graph regularity, ergodic theory, higher-order uniformity.

Every result since — Green-Tao primes APs, 2023 Kelley-Meka breakthrough on Roth, 2024 $2^{n-o(n)}$ Erdős-Szekeres bounds — lives in framework Szemerédi created.

Regularity Lemma — accidental revolution

To prove theorem, Szemerédi needed to handle graph encoding of APs.

Invented 1975 regularity lemma:

Any large graph can be partitioned into $M(\epsilon)$ pieces — bounded number independent of graph size — such that most pairs of pieces are ϵ -pseudorandom — edges between pieces behave like random bipartite graph with density d_{ij} .

Huge graph $\rightarrow$ compressed into small weighted template — Szemerédi graphon picture.

Consequences:

- Graph removal lemmas — if few copies of H , can delete few edges to destroy all H .
- Property testing — can test triangle-freeness in $O(1)$ queries, independent of graph size.
- Computer science — algorithmic regularity, approximation algorithms for dense Max-Cut, etc.
- Itself stronger than Szemerédi theorem — cornerstone of extremal graph theory.

Lemma now taught before theorem — tool bigger than application.

Alternative proofs opened fields:

- **Furstenberg 1977 ergodic theory proof** — translated combinatorial statement into dynamical system.

Idea: set A with $\bar{d}(A) > 0 \rightarrow$ measure-preserving system (X, T) and set E measure $\geq \delta$ such that A contains k -AP iff $\mu(E \cap T^{-d}E \cap \dots \cap T^{-(k-1)d}E) > 0$.

Density $\rightarrow$ recurrence: system returns close to itself infinitely often.

Then Furstenberg proved multiple recurrence theorem — ergodic Szemerédi.

Impact: created ergodic Ramsey theory — field connecting number theory to dynamics. Furstenberg correspondence principle now standard tool: combinatorics $\leftrightarrow$ dynamics.

Furstenberg won Abel Prize 2020 partly for this translation.

- **Gowers 2001 Fourier-analytic proof with higher-order uniformity**

Roth 1953 proved $k = 3$ case using Fourier analysis — large Fourier coefficient $\Rightarrow$ density increment on arithmetic progression.

For $k = 4$, classical Fourier fails — need quadratic Fourier.

Gowers invented U^d uniformity norms — measure correlation with polynomial phases degree $< d$.

U^2 = Fourier uniformity, U^3 detects quadratic structure, etc.

Decomposition: dense set $A = f_{structured} + f_{small} + f_{uniform}$ where $f_{structured}$ is nilsequence-like, $f_{uniform}$ small in U^{k-1} norm so counts k -APs correctly.

Proved Szemerédi with tower-type bounds — first reasonable bound after 60 years.

Invented higher-order Fourier analysis, won Fields Medal 1998.

That proof led directly to Green-Tao: primes uniform in U^d after W -trick, so contain APs.

So Szemerédi's theorem significance triple:

1. Theorem itself — density forces arbitrary APs — final form of van der Waerden strengthening.
2. Tools created to prove it — regularity lemma, ergodic correspondence, Gowers norms — each became field.

3. Philosophy — same as your central lesson: **randomness shallow**. Density $\delta > 0$ budget finite, eventually structure unavoidable. But unlike $W(2, 3) = 9$ you can see, here structure unavoidable for reason deep enough to need new mathematics to explain *why* randomness collapses.

van der Waerden vs Szemerédi — Core Distinction

Partition vs Density

FEATURE	VAN DER WAERDEN	SZEMERÉDI
Framework	Ramsey / Coloring	Density / Additive combinatorics
Condition	$\mathbb{Z}$ split into r colors — whole set colored	Single $A \subset \mathbb{Z}$ with $\bar{d}(A) > 0$ — no partition needed
Guarantee	At least one color class has k -AP	This specific dense A has k -AP
Strength	Weaker	Stronger — implies van der Waerden

van der Waerden — needs everything colored.

Host is whole $\mathbb{N}$ partitioned into r colors. You paint every house on infinite street red/blue/yellow... r colors total.

Pigeonhole: you covered all integers, so average color occupies $1/r$ fraction. But theorem does not use thickness directly — it uses that you *partitioned* everything. If you miss even one integer uncolored, hypothesis fails.

Think: you try to avoid red k -AP and blue k -AP simultaneously by clever global coloring. $W(r, k)$ says you can delay up to $W - 1$, but at W any complete coloring forces monochrome k -AP somewhere.

Needs global info: you must know color of every integer up to W to guarantee.

Szemerédi — needs only one thick set.

Host is single A with positive upper density — $\bar{d}(A) = \limsup |A \cap [1, N]|/N > 0$ — e.g., contains 1% of numbers infinitely often.

You don't care about complement. A^c can be anything, not colored, missing. A alone thick enough forces its own k -AP.

Think: pick any 50% of integers trying to avoid k -AP — e.g., choose numbers avoiding $a, a + d, a + 2d$. Try as hard as you want, hide gaps cleverly. Szemerédi says impossible for every k large — if you keep density > 0 , AP appears inside your set.

So van der Waerden needs partition budget r , Szemerédi needs density budget $\delta > 0$.

van der Waerden: **global partition forces order somewhere**. Szemerédi: **local thickness forces order inside this set**.

That is why Szemerédi $\Rightarrow$ van der Waerden: r -coloring of $\mathbb{N} \Rightarrow$ some color class has density $\geq 1/r > 0 \Rightarrow$ by Szemerédi that class contains k -AP.

Why Szemerédi $\Rightarrow$ van der Waerden

Setup: Color $\mathbb{N} = C_1 \cup \dots \cup C_r$ disjoint r -coloring — every integer gets exactly one color.

Goal: Show some C_i contains k -AP for every k — that's van der Waerden.

Step 1: Pigeonhole for density

For each N , $\sum_{i=1}^r |C_i \cap [1, N]| = N$ — counts partition $[1][N]$

Divide by N : $\sum |C_i \cap [1, N]|/N = 1$.

Take $\limsup_{N \rightarrow \infty} \sum \bar{d}(C_i) \geq 1$ — upper densities cannot all be small, sum at least 1.

If all $\bar{d}(C_i) < 1/r$, say $\leq 1/r - \epsilon$, sum $\leq r(1/r - \epsilon) = 1 - r\epsilon < 1$ contradiction.

Thus exists i_0 with

$$\bar{d}(C_{i_0}) \geq 1/r > 0$$

One color occupies at least $1/r$ fraction infinitely often — pigeonhole principle for density.

Step 2: Apply Szemerédi to thick color

Szemerédi hypothesis: Any $A \subset \mathbb{N}$ with $\bar{d}(A) > 0$ contains k -AP for every k .

Our C_{i_0} has $\bar{d} \geq 1/r > 0 \Rightarrow$ hypothesis satisfied.

Conclusion: For every k , C_{i_0} contains k -term AP $a, a + d, \dots, a + (k - 1)d$.

That's monochromatic k -AP in color i_0 .

Step 3: Conclude van der Waerden

Thus any r -coloring of $\mathbb{N}$ yields monochromatic k -AP.

For finite interval, compactness argument gives existence of finite $W(r, k)$ — if infinite version holds, finite threshold must exist, else take sequence of colorings avoiding k -AP on and limit to coloring of $\mathbb{N}$ avoiding — contradiction. $[1][N]$

So $W(r, k)$ finite follows.

Coloring forces a color to be dense ($\geq 1/r$) $\Rightarrow$ density forces AP (Szemerédi) $\Rightarrow$ coloring forces AP (van der Waerden).

Szemerédi strictly stronger — it tells you *which* color is forced to be dense one, and works even when you don't have partition of whole $\mathbb{N}$, only one dense set.

Converse false: van der Waerden says some color of partition has AP, but doesn't guarantee a prescribed density- $1/2$ set has AP unless you know its complement is other color.

Why Converse False — Density Strictly Stronger

Van der Waerden cannot guarantee AP in prescribed set density $1/2$ unless you know complement.

The gap: van der Waerden says:

For partition $\mathbb{N} = C_1 \cup C_2$, some C_i has k -AP.

It does NOT say *which* i . Could be C_1 , could be C_2 . If you care about specific A with density $1/2$, van der Waerden applied to $A \cup A^c$ tells you *either* A or A^c has k -AP — but you don't know which. If the AP lands in A^c , you learned nothing about A .

Example you gave: A = numbers with first digit 1 — 1, 10 — 19, 100 — 199, ... Upper density ≈ 0.11 ? Actually Benford — $\limsup |A \cap [1, N]|/N = 5/9$? In $\mathbb{N}$ about $1111/1999 \approx 0.55$, infinitely often high, so $\bar{d}(A) > 0$. Van der Waerden says for $A \cup A^c$, one has k -AP — trivial, but could be A^c every time for each k ? No, but vdW alone doesn't rule out that AP always lands in complement. You need Szemerédi to force AP inside A itself.[1][1999]

Formally:

- vdW is $\exists i : C_i$ contains k -AP.
- Szemerédi is $\forall A$ with $\bar{d}(A) > 0$: A contains k -AP.

Second quantifier stronger — universal over sets, not existential over partition cell.

Two ways converse fails:

1. **Prescribed set:** Let A density $1/2$ — e.g., take random half of integers avoiding 3-AP as long as possible? Can you avoid? Szemerédi says no — for large enough N , any $N/2$ -sized subset of $\mathbb{N}$ contains k -AP. vdW does not imply this. vdW only if you 2-color whole $\mathbb{N}$ and look, but if you only look at A alone — without coloring complement — hypothesis of vdW not satisfied, because vdW needs coloring of all $\mathbb{N}$. Knowing A density $1/2$ alone does not give you a coloring where A is a color class? Actually it does — $A \cup A^c$ is coloring — but vdW says *some* color has AP, not necessarily A . So A could still be AP-free if AP hides in A^c . Szemerédi rules that out.[1][N]
2. **Density zero sets:** A sparse, density 0 like primes, squares. vdW says nothing — partition $\mathbb{N} = A \cup A^c$ — A^c density 1, so vdW likely forces AP in A^c , not in A . Squares have density 0 and indeed contain no 4-term AP — consistent. Primes have density 0 but *do* contain arbitrarily long APs — this needs Green-Tao, beyond both vdW and Szemerédi. vdW cannot even ask question because host not partition-dense.

Hence hierarchy:

van der Waerden < Szemerédi < Green-Tao relative Szemerédi

- vdW: finite partition forces AP somewhere.
- Szemerédi: positive density forces AP inside set itself.
- Green-Tao: relative density inside pseudorandom host forces AP even when absolute density 0.

Each step weakens hypothesis about host — from covering whole $\mathbb{N}$ with r colors, to one thick set, to sparse but pseudorandomly embedded set — and conclusion stays: AP unavoidable.

That is why Szemerédi strictly stronger — it isolates property that makes van der Waerden true: not coloring, but thickness.

Restatement

1. van der Waerden — finite threshold, partition

For any r, k , exists $W(r, k)$ such that if $\{1, \dots, W\}$ r -colored, then some color contains k -AP.

Host = interval $\mathbb{N}$, length $W(r, k)$. Condition = you color every integer in host with r colors.

Conclusion = monochromatic k -AP somewhere, in some color.

Quantitative — gives number $W(r, k)$. We know $W(2, 3) = 9$, $W(2, 4) = 35$, $W(2, 6) = 1132$ — actual thresholds where avoidance ends.[1][W]

Philosophy = **partition regularity**: finite coloring cannot destroy all arithmetic structure. Disorder budget = r colors, $W - 1$ length where you can still avoid.

Same as $R(3, 3) = 6$: color edges K_6 2 colors, monochromatic triangle forced — finite threshold forces graph order.

2. Szemerédi — no number, density

If $A \subset \mathbb{N}$ and $\limsup_{N \rightarrow \infty} \frac{|A \cap [1, N]|}{N} > 0$ then for every k , A contains k -term AP. [1][N]

Host = any $A \subset \mathbb{N}$ with positive upper density — e.g., $\geq 1\%$ infinitely often. No interval length given.

Condition = thickness alone, no coloring, no partition of complement needed.

Conclusion = this specific A contains k -AP for every k — arbitrarily long, but for each k possibly different AP.

Non-quantitative in statement — no explicit W , existence for each k with $N(\delta, k)$ depending on δ . Gowers gives tower bound.

Philosophy = **density regularity**: positive fraction cannot avoid all APs. Disorder budget = $\delta > 0$ density, you can avoid up to $N(\delta, k) - 1$ inside A , at $N(\delta, k)$ forced.

Same as Erdős-Szekeres density version, Turán vs Ramsey: Turán says dense graph forces K_t , Ramsey says complete graph colored forces monochrome K_t . Szemerédi is Turán-type strengthening of van der Waerden's Ramsey-type.

Summary mapping:

- van der Waerden: r colors $\rightarrow$ one color density $\geq 1/r \rightarrow$ forces k -AP. Needs pigeonhole to get density, then density $\rightarrow$ AP hidden inside.
- Szemerédi: density $> 0 \rightarrow$ forces k -AP directly, no pigeonhole needed.

Thus Szemerédi isolates core: thickness, not partition, is what forces arithmetic order.

van der Waerden first showed $K = 0$ vs $W - B$ vs $W(r, k)$ all same — budget finite, order forced. Szemerédi showed budget can be measured as $\delta > 0$ fraction, far more general — you don't need to paint whole street, owning 1% of houses thickly enough forces equally spaced houses in your set.

Hierarchy of Forcing — Where It Sits in Your Story

You have now full ladder — same meta-theorem in different hosts:

- **Gomory**: budget $W - B$ black-white imbalance — count forces no tiling. If $W \neq B$, domino tiling impossible. Disorder = color imbalance.
- **Egregium**: budget K Gaussian curvature — curvature forces no isometry. If $K_1 \neq K_2$, surface cannot be bent isometrically. Disorder = curvature difference.
- **Ramsey** $R(3, 3) = 6$: budget $N = 5$ vertices — 6 forces monochromatic triangle. Host = complete graph, coloring edges 2 colors. Disorder = avoid monochrome triangle.
- **Hales-Jewett** $HJ(k, c)$: budget dimension $H = HJ - 1 - HJ$ forces monochromatic combinatorial line. Host = word cube $[k]^H$. This implies van der Waerden by coding $N(w) = \sum w_i M^i$ — AP becomes variable word. Disorder = dimension.

- **van der Waerden** $W(r, k)$: budget $W - 1$ length — W forces monochromatic k -AP. Host = interval $\mathbb{N}$, coloring integers r colors. Disorder = length of avoidance coloring.
- **Erdős-Szekerés** $N(n) = 2^{n-2} + 1$: budget 2^{n-2} points — $2^{n-2} + 1$ forces convex n -gon. Host = point set in general position. Disorder = number of points.
- **Szemerédi**: budget $\delta > 0$ density — positive density forces arbitrarily long AP. Host = any $A \subset \mathbb{N}$ with $\bar{d}(A) > 0$. Implies van der Waerden — partition into r colors gives one color density $\geq 1/r > 0$, apply Szemerédi.

All same:

Disorder budget finite. You can avoid structure for a while with clever construction, but size/density/dimension/curvature budget exhausted $\rightarrow$ structure forced.

For Szemerédi, budget measured by density δ :

- If you keep $\delta > 0$ — keep δ fraction of integers infinitely often — you can avoid k -AP up to some $N(\delta, k)$ by clever sparse choice, but beyond that progression forced.
- If $\delta \rightarrow 0$, $N(\delta, k) \rightarrow \infty$ — sparser you are, longer you can avoid. Primes $\delta = 0$, so Szemerédi does not apply — need Green-Tao relative density inside pseudorandom host.
- Like HJ and $N(n)$, proof non-constructive in sense of location: Szemerédi proves k -AP exists in A , does not tell where a, d are — size alone forces existence, not position. Same as regularity lemma gives existence of partition, not explicit partition; same as van der Waerden proves $W(r, k)$ exists, does not give explicit coloring up to $W - 1$ — SAT solver finds it for $W(2, 6) = 1132$.

That non-constructiveness is feature: Ramsey theory never says *where* order appears, only that it must — disorder cannot be extended forever.

This completes arc from Gomory elementary counting to Szemerédi density — each step weakens hypothesis, strengthens conclusion, keeps same message: **randomness shallow, order inevitable when host large enough.**

Green-Tao Theorem — Inevitable Order Inside the Primes

Final step in 100-year progression:

- **van der Waerden 1927**: Coloring $\mathbb{N}$ with r colors $\rightarrow$ monochromatic APs of any length. Need to color *everything*.
- **Szemerédi 1975**: Density $> 0 \rightarrow$ APs of any length inside *that set*. Need thickness > 0 .
- **Green-Tao 2004**: Primes density 0 $\rightarrow$ still APs of any length. Even sparse set that *looks random* forces APs.

Green-Tao 2004: For every k , there exist $a, d > 0$ such that $a, a + d, \dots, a + (k - 1)d$ all prime.

You can find k primes perfectly equally spaced, k as large as you want.

Primes look random locally, globally cannot avoid arithmetic structure.

What It Says — And What It Does Not

Simple picture: Arithmetic progression = equally spaced stepping stones across number line. Step size d is distance between stones.

- 3, 7, 11: $a = 3, d = 4$, stones at $3 + 0 \cdot 4, 3 + 1 \cdot 4, 3 + 2 \cdot 4$.
- 5, 11, 17, 23, 29: $a = 5, d = 6$, five stones.

Green-Tao: you can find k prime stones in perfect line, for any k you name. Want 10 prime stones equally spaced? Exists somewhere far out. Want 100? Exists, just much farther.

Local primes look random — gaps irregular — but global desert still contains arbitrarily long straight highways made entirely of prime stones.

Arbitrary vs Infinite

- Arbitrary length = for each $k = 3, 4, 5, \dots$ there is *some* prime AP length k . Different k may have totally different a, d . For $k = 3, a = 3, d = 4$ works. For $k = 5, a = 5, d = 6$ works. For $k = 100$, there is some a_{100}, d_{100} enormous — proof says exists, doesn't tell you numbers.
- Not infinite length = one AP $a, a + d, a + 2d, \dots$ infinite all primes is impossible. Proof in one line: term number a steps ahead is $a + a \cdot d = a(1 + d) = a$ times $(1 + d) \rightarrow$ composite if $a > 1$. So after a steps you always hit multiple of first term.

Analogy: You can find arbitrarily long straight segments of road in desert, but you cannot have infinite straight road with no bends that stays on prime stones forever — you'll eventually hit a non-prime rock because a divides that future term.

Think: finite rows possible, infinite row impossible — just like van der Waerden says arbitrarily long APs in one color, not infinite AP.

Infinitude for fixed k .

Green-Tao proves not just existence of one k -AP, but infinitely many distinct k -APs for each fixed k . And count up to N :

$$\#\{\text{prime } k\text{-APs with } a \leq N, d \leq N\} \sim C_k \frac{N^2}{(\log N)^k}$$

$C_k > 0$ constant depending on k — Hardy-Littlewood prime tuples constant.

Simple: infinitely many prime triples equally spaced, infinitely many prime 5-tuples, etc. If you search to $N = 10^{12}$, you expect about $N^2/(\log N)^k$ of them — many for $k = 3$, fewer for larger k .

Why common difference explodes:

Why must d be huge? Small primes force divisibility.

- $k \geq 3$: All primes > 2 odd. If d odd, a odd + odd = even $\rightarrow a + d$ even composite. So d must be even = divisible by 2. Otherwise you'd alternate odd/even and hit even composite immediately.
- $k \geq 4$: Consider mod 3. Numbers mod 3 are 0, 1, 2. If d not multiple of 3, then $a, a + d, a + 2d$ cover all three residues mod 3 — pigeonhole. One residue is 0 mod 3 $\rightarrow$ that term divisible by 3. If term > 3 , composite. To have 4 primes > 3 all prime, you must avoid hitting 0 mod 3, so need $3|d$. So d multiple of 3.
- $k \geq 6$: Similarly mod 5. If $5 \nmid d$, then $a, a + d, \dots, a + 4d$ cover all residues mod 5 $\rightarrow$ one term 0 mod 5 composite. To have 6 primes > 5 , need $5|d$.

General rule: To have k primes all $>k$, d must be divisible by every prime $p < k$. Otherwise progression hits $0 \bmod p$.

This product is primorial:

$$k\# = \prod_{p < k} p$$

- $k = 5$: primes < 5 are $2, 3 \rightarrow k\# = 6$. Indeed $5, 11, 17, 23, 29 \equiv d \pmod{6}$.
- $k = 6$: primes < 6 are $2, 3, 5 \rightarrow 30$. Must have d multiple of 30. Smallest 6-term example $7, 37, 67, 97, 127, 157 \equiv d \pmod{30}$.
- $k = 10$: primes < 10 are $2, 3, 5, 7 \rightarrow 210$. Indeed $199, \dots, 2089 \equiv d \pmod{210}$.
- $k = 27$: need divisible by all primes $< 27 \rightarrow 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 = 223,092,870 = 23\#$.

So even before you search, number theory forces d huge. This is why $k = 27$ record has $d \approx 9.6 \times 10^{17}$ — it must be multiple of 223 million, and random prime density $1/\log N$ tiny, so you need enormous search space to get lucky.

Takeaway: primes want to be AP, but small primes are obstacles — to avoid being divisible by 2,3,5,..., you must step by multiples of them. The longer AP you want, the more obstacles, the bigger mandatory step.

The Central Difficulty: Density Zero

Density answers: if you look at first N numbers $1, \dots, N$, what fraction belongs to your set A ?

$$d_N(A) = \frac{|A \cap [1, N]|}{N}$$

If $d_N(A)$ stays above some $\delta > 0$ infinitely often — even if it wiggles — we say A has positive upper density $\bar{d}(A) > 0$.

- **Evens:** $2, 4, 6, \dots$ Up to N , about $N/2$ evens $\rightarrow d_N \approx 1/2 \rightarrow \bar{d} = 1/2 > 0$. Thick.
- **Numbers starting with 1:** $1, 10 - 19, 100 - 199, \dots$ Up to 1999, you have $1 + 10 + 100 + \dots = 1111$ numbers $\rightarrow 1111/1999 \approx 55\%$. Up to 2999, fraction drops to $\approx 37\%$. It oscillates, but infinitely often hits $> 50\%$ — $\limsup > 0$. So positive upper density even though no limit. Still thick infinitely often.
- **Squares:** $1, 4, 9, 16, \dots$ Up to N , about $\sqrt{N}$ squares $\rightarrow d_N \approx \sqrt{N}/N = 1/\sqrt{N}$. For $N = 10^6$, $1000/10^6 = 0.1\%$. For $N = 10^{12}$, $10^6/10^{12} = 0.0001\% \rightarrow 0$. Density 0 — sparse.
- **Primes:** Prime Number Theorem: $\pi(N) \sim N/\log N$ primes up to $N \rightarrow d_N \sim 1/\log N$.
 - $N = 10^6$: $\log N \approx 13.8 \rightarrow \sim 7.2\%$ predicted, actually 7.8%.
 - $N = 10^{12}$: $\log N \approx 27.6 \rightarrow \sim 3.6\%$.
 - $N = 10^{100}$: $\log N \approx 230 \rightarrow \sim 0.4\%$.

Goes to 0 as $N \rightarrow \infty$. In limit, primes occupy 0% of integers. Density 0 — vanishing fraction.

Why this kills Szemerédi:

Szemerédi hypothesis: $\limsup |A \cap [1, N]|/N > 0$ — need to contain e.g., 1% of numbers infinitely often. Then theorem says A contains k -AP for every k — e.g., set with 1% contains 1000-term AP somewhere.

Primes fail: eventually contain $< 1\%$, $< 0.1\%$, $< 0.0001\%$... So hypothesis false, theorem says nothing.

And density-zero sets *can* avoid APs! Example powers of 2: $1, 2, 4, 8, 16, 32, \dots$ — grows exponentially, density 0, and contains no 3-term AP. Why? If $2^a, 2^b, 2^c$ in AP, then $2^b - 2^a = 2^c - 2^b \rightarrow 2^a(2^{b-a} - 1) = 2^b(2^{c-b} - 1) \rightarrow$ impossible for

$a < b < c$ unless trivial. So sparse sets can escape.

So primes could have been like powers of 2 — sparse enough to avoid APs forever. Szemerédi cannot rule out.

Why primes different?

Powers of 2 structured sparse — lacunary. Primes pseudorandom sparse — they look random locally, not growing regularly. They *should* contain APs if they were random set with same density $1/\log N$ — random set with $N/\log N$ points expects many k -APs for small k .

But random heuristic not proof — need to prove pseudorandomness enough to force APs despite density zero.

Green-Tao insight: Don't apply Szemerédi to primes in integers. Apply Szemerédi *relative* to a larger pseudorandom host that has density 1, inside which primes are relatively dense.

Think: desert vs oasis. Primes = trees density 0 in whole desert. Can't apply Szemerédi to desert. But find oasis ν covering 1% of desert, looking random, containing trees at 10% density inside oasis. Inside oasis, trees are *relatively* dense — relative density >0 — so relative Szemerédi forces AP among trees.

That is transference principle — transfer density theorem from dense world to sparse pseudorandom world.

The Breakthrough: Transference Principle — Simple Explanation

Proof 50 pages, but idea now standard template — used for polynomial primes, Gaussian primes, etc.

Metaphor: Imagine desert huge — integers. Trees = primes — very sparse, 0% of desert. Want to prove there is straight line of k trees equally spaced.

You cannot prove directly — trees too sparse. Random 3 trees almost never equally spaced.

Trick: Find oasis ν inside desert that:

- Covers say 1% of desert — not too small.
- Looks completely random — if you test it with Fourier, Gowers norms, it indistinguishable from random cloud.
- Contains all trees — wherever tree, oasis cloud at least as high.

If trees occupy 10% of oasis — relatively dense inside oasis — then inside oasis, Szemerédi applies! Because inside oasis world, trees have positive relative density.

So problem reduces to building such oasis and proving Szemerédi *relative* to oasis.

Step 1: Build pseudorandom majorant — Selberg envelope

We need function $\nu : [1, N] \rightarrow \mathbb{R}_{\geq 0}$ — think weight at each n .

Properties:

1. **Majorizes primes:** $\nu(n) \geq c \cdot 1_{\text{prime}}(n)$. Where n prime, $\nu(n)$ at least constant c . Where n composite, $\nu(n)$ can be small but nonnegative. So primes sit under cloud.

How built? Goldston-Yıldırım / Selberg sieve — classic sieve weight that approximates prime indicator but smoother, less spiky. Truncated divisor sum: $\nu(n) \approx (\sum_{d|P(n), d \leq R} \mu(d) \log(R/d))^2$ — roughly counts n with no small prime factors —

pseudoprime detector. Not 0/1, but nonnegative, mean ~ 1 .

2. **Pseudorandom:** ν looks like constant 1 for counting linear patterns. Formal: $\|\nu - 1\|_{U^{k-1}}$ small — Gowers uniformity norm small. Also linear forms condition — number of solutions to systems like $\nu(n)\nu(n+d)\dots$ matches random.

Test ν with any Fourier wave $e^{2\pi i \alpha n}$ — average ≈ 0 like random. Test with quadratic phase $e^{2\pi i \alpha n^2}$ — also random. So cloud has no hidden arithmetic bias.

3. **Mean 1:** $\mathbb{E}_{n \leq N} \nu(n) = 1 + o(1)$. So total mass N — like constant 1 function mass. Not sparse — occupies positive fraction.

W-trick — killing trivial bias.

Primes > 2 odd $\rightarrow$ not random mod 2 — all 1 mod 2. Similarly mod 3, primes > 3 are 1, 2 mod 3 not 0. This is not pseudorandom — fails linear forms condition.

Fix: Let w slowly growing, say $\log \log N$, $W = \prod_{p \leq w} p$ primorial of small primes. Look only at arithmetic progression $Wn + 1$ — numbers congruent 1 mod W . Inside this progression, primes equidistributed — no bias mod small primes because we fixed residue.

Define new universe $[N']$ where $N' \approx N/W$, and study $Wn + 1$ prime. After this W -trick, ν becomes truly pseudorandom — W -tricked primes have no residue bias.

Analogy: Primes are like people who never sit in certain chairs — obviously not random. W -trick removes those chairs from room, then remaining seating looks random.

Step 2: Prove relative Szemerédi — heart

Gowers' proof of Szemerédi 2001:

Take bounded $f : [N] \rightarrow [0, 1]$, mean $\geq \delta$. Want to count k -APs: $\mathbb{E}_{a,d} f(a)f(a+d)\dots f(a+(k-1)d)$.

Either f Gowers-uniform — U^{k-1} small — then by generalized von Neumann, count $\approx \delta^k N^2$ — random number, many APs.

Or f not uniform — U^{k-1} large — then Gowers inverse theorem says f correlates with nilsequence — structured object like $e^{2\pi i(\alpha n^2 + \beta n)}$ for $k = 4$. Then you can decompose $f = f_{str} + f_{sml} + f_{unif}$, density increment on structured piece — iterate, density increases, must stop, find AP.

Green-Tao need same but with $0 \leq f \leq \nu$, not $0 \leq f \leq 1$. ν unbounded pointwise but mean 1, pseudorandom.

Prove:

Relative Szemerédi: If ν pseudorandom (linear forms + correlation conditions), and $0 \leq f \leq \nu$, $\mathbb{E}f \geq \delta > 0$, then f contains $\geq c(\delta, k)N^2$ k -APs.

Ingredients simple terms:

- **Generalized von Neumann relative:** If f bounded by ν and U^{k-1} small relative to ν , then AP count close to random. Needs ν pseudorandom so Cauchy-Schwarz still works.
- **Dense model + inverse theorem:** If f not uniform, need to find structure. Show f has dense model $\tilde{f}$ bounded by 1 with similar Gowers norm — transference. Then apply usual inverse theorem to $\tilde{f}$, lift correlation back to f . This uses

Hahn-Banach separation — Green-Tao dense model theorem — if ν pseudorandom, any $f \leq \nu$ approximable by bounded $\tilde{f}$.

- **Structure theorem + density increment:** Once correlate with nilsequence, run energy increment as Gowers.

Plain language: If universe ν random enough, any relatively dense subset f inherits Ramsey property. Random universe cannot hide structure to help you avoid APs — any thick subset inside it still forces APs.

Step 3: Transfer primes into envelope

Final check: primes actually dense inside ν .

After W -trick, define normalized prime indicator:

$$\tilde{f}(n) = \frac{\phi(W)}{W} \log N \cdot 1_{\text{prime}}(Wn + 1)$$

Factor $\frac{\phi(W)}{W} \log N$ — normalizes mean to ≈ 1 . Prime number theorem in AP says primes $\equiv 1 \pmod{W}$ have density $\sim 1/\phi(W) \cdot 1/\log N$, so after multiply by $\phi(W)/W \cdot \log N$, mean $\approx 1/W$? *Actually* ≈ 1 — constant.

Goldston-Yildirim construction ensures $0 \leq \tilde{f} \leq \nu$ pointwise, $\mathbb{E}\tilde{f} \gg 1$ — say $\geq \delta = 1/2$.

So primes relatively dense in ν .

Apply relative Szemerédi to $\tilde{f} \rightarrow \tilde{f}$ contains many k -APs $\rightarrow$ those correspond to k -APs of primes $Wn + 1 \rightarrow k$ -APs of primes in original integers.

Done.

Why this template powerful: Once you have pseudorandom majorant for any sparse set — polynomial values of primes, Gaussian primes, etc. — same relative Szemerédi gives APs there. Green-Tao opened industry: transference principle.

Records: What We Can Actually Find — Simple Explanation

Green-Tao is **purely existential** — like Erdős-Szekeres says $2^{n-2} + 1$ points force convex n -gon but doesn't tell you where. It says *some* prime k -AP exists somewhere, no bound how far you must search.

Computationally brutal because of $k\#$ divisibility we explained.

Small examples you can verify by hand:

- $k = 3$: 3, 7, 11 — step $d = 4$. Check: 3 prime, 7 prime, 11 prime, equally spaced. Many triples exist.
- $k = 4$: 5, 11, 17, 23 — $d = 6$. All prime? 5, 11, 17, 23 yes.
- $k = 5$: 5, 11, 17, 23, 29 — $d = 6$ again, minimal. Why $d = 6$ minimal? Must be multiple of $6 = 2 \cdot 3$ to avoid hitting multiple of 2 or 3. So 6 smallest possible.
- $k = 6$: 7, 37, 67, 97, 127, 157 — $d = 30 = 2 \cdot 3 \cdot 5$. Check difference 30 each time. Why 30? Need divisible by 2, 3, 5 for $k \geq 6$. So 30 minimal. This progression works.

Notice pattern: As k grows, minimal allowed d jumps at each prime.

Bigger — why search explodes:

- $k = 10$: 199, 409, 619, 829, 1039, 1249, 1459, 1669, 1879, 2089 — $d = 210 = 2 \cdot 3 \cdot 5 \cdot 7$. 210 mandatory. Starting at 199 to avoid small composites hitting. You can test with computer — each term prime.

Simple reasoning: To get 10 primes, need d multiple of 210, else among 7 terms you'd hit multiple of 7. So search space steps of 210.

- $k = 22$: Found 2008 by Wróblewski — required d multiple of $19\# = 9699690$ — almost 10 million step minimal. So you must check numbers spaced 10 million apart for 22 primes all prime — probability tiny: each $\sim 1/\log N$ prime, so k primes probability $(1/\log N)^k$ — needs N huge to get one success. For $k = 22$, N astronomically large.
- $k = 25$: Formula $a_n = 6171054912832631 + n \cdot 366384 \cdot 23\# - a_0$ 16 digits, $d = 366384 \cdot 223,092,870 \approx 8.1 \times 10^{13}$. 25 terms. Found by Chermoni & Wróblewski 2008 after months distributed search.
- $k = 27$ — **current world record 2019** Rob Gahan + PrimeGrid volunteers:

$$a_n = 2245848550833 + n \cdot 4314276143 \cdot 23\# \quad n = 0..26$$

where $23\# = 223,092,870$, so

$$d = 4,314,276,143 \times 223,092,870 \approx 9.62 \times 10^{17}$$

26 steps span $26d \approx 2.5 \times 10^{19}$ — from 2.2 trillion to 25 quintillion range. Each of 27 numbers proven prime with deterministic Miller-Rabin + ECPP. Search required years of CPU on PrimeGrid BOINC.

Why $23\#$? For $k = 27$, need d divisible by all primes $< 27 - 2, 3, 5, 7, 11, 13, 17, 19, 23 \rightarrow 23\#$. Could also be divisible by larger prime factor 4,314,276,143 extra — to fine-tune residues to avoid small prime divisors of a_n .

Why we will never find $k = 100$ by brute force — simple math:

- d must be divisible by $97\# = 2 \cdot 3 \cdot 5 \cdots 97$ — product of primes < 100 . This is about 2.3×10^{36} — 37 digits. Minimal step 37 digits.
- Heuristic: Probability k numbers $a + nd$ all prime $\approx 1/(\log N)^k$. For $k = 100$, $(\log N)^{100}$ huge — need N enormous, far beyond 10^{100} — a would have hundreds of digits, search space $(\log N)^{100} \approx$ impossible.
- Green-Tao gives no explicit $N(\delta, k)$ — best bounds tower-exponential — so doesn't help locate.

So Green-Tao tells us length-100 prime AP exists *somewhere* — maybe at 10^{1000} or $10^{10^{100}}$ — but we will never see it.

This is same non-constructive flavor as van der Waerden $W(2, 6) = 1132$ — existence proved 1927, exact value found only 2008 via SAT solver searching 30 orders of magnitude. And $W(2, 7)$ still unknown, estimated $> 10^4$.

Takeaway: Existence $\neq$ construction. Ramsey theory guarantees order eventually, but location may be vastly beyond computational reach. Primes contain 100-term equally spaced line, but universe too big to find it — like guaranteeing a specific grain of sand exists somewhere in cosmos without telling which galaxy.

Why It Is Culmination — Simple Explanation for First Time

This is end of 100-year story with same moral, each time stronger:

Ladder:

- **Gomory 1960s:** Chessboard 8×8 minus two opposite corners cannot be tiled by dominoes. Why? Count $W = 30$ white, $B = 32$ black, each domino covers 1 white 1 black. Budget $W - B \neq 0$ forces impossibility. Simple counting forces global structure.
- **Egregium Theorema Gauss 1827:** Curvature K budget forces no isometry. Sphere cannot be flattened to plane because $K = 1/R^2$ vs 0. Budget = curvature difference.
- **Ramsey $R(3, 3) = 6$ 1930:** Host = 6 people, color edge red = knows, blue = strangers. Budget 5 people can avoid monochrome triangle, 6 forces it. Complete disorder in friendships impossible.
- **Hales-Jewett 1963:** Host = high-dimensional cube $[k]^n$ — words length n . Budget dimension $n = HJ(k, c) - 1$ can avoid monochrome combinatorial line, HJ forces it. This implies van der Waerden — AP coded as word.
- **Erdős-Szekeres 1935:** Host = points in plane. Budget 2^{n-2} points can avoid convex n -gon, $2^{n-2} + 1$ forces it. Geometry forces order.
- **van der Waerden 1927:** Host = integers colored with r colors. Budget $W(r, k) - 1$ length can avoid k -AP monochrome, W forces it. Disorder = finite coloring, cannot destroy all APs.
- **Szemerédi 1975:** Host = dense set density $\delta > 0$. Budget $N(\delta, k) - 1$ size can avoid k -AP, N forces it. Disorder = thickness, not coloring. Much stronger — you don't need to color everything, just be thick.
- **Green-Tao 2004:** Host = primes density 0. Budget = pseudorandomness. Primes look random, but relative density inside pseudorandom envelope ν positive $\rightarrow$ forces k -AP.

Each step weakens hypothesis — needs less about host — but keeps same conclusion: AP / convex / monochrome forced.

Disorder budget finite.

For Gomory, budget = imbalance count. For Ramsey, budget = number of vertices. For van der Waerden, budget = length $W - 1$. For Szemerédi, budget = $1/\delta$ — sparser you are, longer you can avoid, but if $\delta > 0$ fixed, avoidance length finite.

For Green-Tao, budget = how random you look. Primes look random locally — gaps irregular, no formula — but globally their randomness is *shallow*. They live inside Selberg sieve envelope ν that is truly random for counting APs. Since primes occupy positive fraction of envelope, they inherit envelope's Ramsey property.

- **Van der Waerden:** You paint whole infinite street r colors. Some color must contain equally spaced houses. You must paint everything.
- **Szemerédi:** You don't paint everything. You just own 1% of houses scattered. If you keep 1% forever, some k equally spaced houses you own forced.
- **Green-Tao:** You own 0% in limit — primes get rarer. Naively you could avoid equally spaced houses, like powers of 2 do. But primes are not adversarially placed — they are pseudorandomly placed. Pseudorandom 0% set can still force APs if it sits inside random cloud with positive relative density. Powers of 2 fail because their cloud not pseudorandom — powers of 2 highly structured, not uniform in Gowers norm.

Why primes were supposed to be counterexample:

For century, primes thought enemy of patterns — Euclid proves infinitely many, but no formula. Cramér model 1936 models primes as random set where n prime with probability $1/\log n$ independently. Random set contains k -APs with probability 1 for each fixed k , but random model not proof.

Many thought density-zero sets could avoid APs — squares have no 4-term AP, powers of 2 have no 3-term AP. So primes could be like that.

Green-Tao shows: Even most famous pseudorandom set cannot escape perfect arithmetic order if you look far enough. Chaos only local — up to 10^6 , primes look irregular — globally, order forced at scale 10^{19} for $k = 27$, at far larger scale for $k = 100$.

Final meta-theorem:

Structure vs randomness dichotomy, structure wins eventually.

Every proof in hierarchy uses same scheme: If no structure, then randomness/uniformity $\rightarrow$ counting lemma gives many patterns $\rightarrow$ contradiction. If not random, then correlation with structured object $\rightarrow$ density increment on substructure $\rightarrow$ induction, budget decreases.

For Green-Tao, this dichotomy lifted to relative world — even sparse primes obey it because envelope ν provides random background to compare to.

Thus Green-Tao is ultimate vindication of Ramsey philosophy: **Even the set that looks most random cannot escape perfect arithmetic order.**