

Stirling Numbers

Stirling numbers, named after James Stirling (1692–1770), are two related triangular arrays of numbers that are fundamental in combinatorics. They both count ways to arrange n distinct objects, but they answer two different questions:

- **First kind:** How many ways to arrange them into k *cycles* (circular orderings)?
- **Second kind:** How many ways to group them into k *sets* (unordered collections)?

The distinction between sets and cycles is the core of the theory.

What Sets vs Cycles Means

Take $n = 3$ elements $\{A, B, C\}$.

Set $\{A, B, C\} =$ no order, $\{A, B\} = \{B, A\}$

Cycle $(A B C) =$ circular order, $(A B C) = (B C A) = (C A B) \neq (A C B)$

In a set, internal order irrelevant. In a cycle, who follows whom matters, but rotation of whole cycle irrelevant because table has no distinguished seat. So a set of size m can be arranged into $(m - 1)!$ different cycles. That factor explains why first kind counts are larger than second kind except extremes.

$$S(n, k) = \left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \# \text{ ways into } k \text{ unlabeled sets}$$

$$c(n, k) = \left[\begin{matrix} n \\ k \end{matrix} \right] = \# \text{ ways into } k \text{ cycles}$$

History and Discovery

- **James Stirling** (1692 – 1770), Scottish mathematician, introduced second kind numbers in *Methodus Differentialis* (1730) to express powers in terms of falling factorials for interpolation and series acceleration. He gave tables and recurrence $S(n, k) = kS(n - 1, k) + S(n - 1, k - 1)$ without modern partition interpretation.
- **Earlier roots:** Tables equivalent to $S(n, k)$ appear in Japanese mathematician Seki Takakazu (~ 1683) and in Stirling's contemporary. First kind numbers arise from expansions of $x(x - 1) \cdots (x - n + 1)$, studied by Newton and Stirling for finite differences.
- **Niels Nielsen** (1904) coined term "Stirling numbers of the first and second kind" and distinguished $s(n, k)$ signed vs $c(n, k)$ unsigned. He systematically studied their algebraic properties.
- **Charles Peirce** (1880s) and later **Eric Temple Bell** (1934) connected $S(n, k)$ to set partitions, defining Bell numbers $B_n = \sum_k S(n, k)$ as row sums. Peirce's Bell triangle predates Bell.
- **Modern notation:** $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ and $\left[\begin{matrix} n \\ k \end{matrix} \right]$ brackets/braces introduced by Karamata (1935) and popularized by Knuth (1992) to mirror binomial $\binom{n}{k}$ — mnemonic: $\{\}$ for sets, $[\]$ for cycles.

Stirling Numbers of the Second Kind

Denoted $S(n, k)$ or $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$, pronounced " n brace k ".

Combinatorial definition:

$$S(n, k) = \frac{\# \text{ onto functions } [n] \rightarrow [k]}{k!}$$

Because onto function determines preimage partition into k labeled blocks, divide by $k!$ to forget labels.

Example: $n = 3$ friends $\{A, B, C\}$ into $k = 2$ groups.

$$\{A, B\}\{C\}, \{A, C\}\{B\}, \{B, C\}\{A\}$$

So $S(3, 2) = 3$. The groups are unlabeled — $\{A, B\}\{C\}$ is the same as $\{C\}\{A, B\}$.

Values: $S(n, 0) = 0$ for $n > 0$, $S(0, 0) = 1$, $S(n, 1) = 1$, $S(n, n) = 1$, $S(n, 2) = 2^{n-1} - 1$.

Recurrence:

$$S(n, k) = k S(n - 1, k) + S(n - 1, k - 1)$$

Fix element n :

Case 1: $\{n\}$ not singleton : partition $[n - 1]$ into k blocks $[S(n - 1, k) \text{ ways}]$,

choose one of k blocks to insert n into $[k \text{ choices}]$

Case 2: $\{n\}$ singleton : partition $[n - 1]$ into $k - 1$ blocks $[S(n - 1, k - 1) \text{ ways}]$

Example: $n = 4, k = 2$:

$$S(4, 2) = 2S(3, 2) + S(3, 1) = 2 \cdot 3 + 1 = 7$$

$7 = 4 \text{ ways } 3 + 1 (\{A\}\{BCD\} \text{ type}) + 3 \text{ ways } 2 + 2 (\{AB\}\{CD\} \text{ type})$.

Detailed view of recurrence:

Take partition of $\{1, \dots, n - 1\}$ counted by $S(n - 1, k)$ — k blocks. Element n can be inserted into any of those k blocks — k distinct partitions of n -set with k blocks where $\{n\}$ is not singleton. That's $kS(n - 1, k)$.

If $\{n\}$ is singleton, delete it — you get partition of $n - 1$ into $k - 1$ blocks — $S(n - 1, k - 1)$ possibilities. Disjoint cases cover all.

Example: $S(4, 2) = 2S(3, 2) + S(3, 1) = 2 \cdot 3 + 1 = 7$ — matches $3 + 1$ vs $2 + 2$ count earlier.

Explicit Formula (Inclusion-Exclusion):

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

Derivation:

Number of onto functions $f : \{1, \dots, n\} \rightarrow \{1, \dots, k\}$ is $k!S(n, k)$ because onto function partitions domain into k labeled preimage blocks ($S(n, k)$ ways unlabeled, $k!$ labelings).

Count onto functions by inclusion-exclusion: total functions k^n , subtract those missing at least one value.

$$\begin{aligned} k!S(n, k) &= \sum_{j=0}^k (-1)^j \binom{k}{j} (k - j)^n \\ &= \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n \end{aligned}$$

Divide by $k!$:

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

Total j^n functions into j -subset, inclusion-exclusion over missing values.

Example $n = 3, k = 2$:

$$S(3, 2) = \frac{1}{2!} \left[\binom{2}{0} 0^3 - \binom{2}{1} 1^3 + \binom{2}{2} 2^3 \right] = \frac{1}{2} [0 - 2 + 8] = 3$$

Small values:

$$S(n, 0) = \begin{cases} 1 & n = 0 \\ 0 & n > 0 \end{cases}$$

$$S(n, 1) = 1$$

$$S(n, 2) = 2^{n-1} - 1 = \frac{2^n - 2}{2}$$

$$S(n, n - 1) = \binom{n}{2}$$

$$S(n, n) = 1$$

Relation to Bell Numbers: Summing across k gives the total number of partitions:

$$B_n = \sum_{k=0}^n S(n, k)$$

So B_n is row sum. For $n = 4$: $1 + 7 + 6 + 1 = 15$.

Expanded reading — what $S(n, k)$ is really counting:

$S(n, k)$ = number of ways to split n distinct labeled balls into k identical non-empty boxes

Labeled balls — A, B, C distinct. Identical boxes — only which balls are together matters, not which box is called 1 or 2. Empty boxes not allowed.

Example $n = 3, k = 2$:

$$\{A, B\}\{C\}$$

$$\{A, C\}\{B\}$$

$$\{B, C\}\{A\}$$

If boxes were labeled Box1, Box2, each partition would correspond to $2! = 2$ labeled assignments, so onto functions $[3] \rightarrow [2]$ count $3 \cdot 2 = 6$, and $S(3, 2) = 6/2! = 3$.

$$S(n, k) = \frac{\# \text{onto functions}}{k!}$$

Why the special values make sense:

$$S(n, 1) = 1 \quad \text{All balls in one box: only } \{[n]\}$$

$$S(n, n) = 1 \quad \text{Each ball alone: only } \{1\}\{2\} \cdots \{n\}$$

$$S(n, 0) = 0 \ (n > 0) \quad \text{No way to place balls into no boxes}$$

$$S(n, n-1) = \binom{n}{2} \quad \text{One box has 2 balls, rest singletons: choose the pair}$$

$$S(n, 2) = 2^{n-1} - 1$$

Derivation in counting language: Put ball 1 in box A to break symmetry (boxes unlabeled, but 1's box can be called A). Each of remaining $n - 1$ balls chooses A or B — 2^{n-1} choices. Exclude choice where all choose A leaving B empty — 1 excluded. So $2^{n-1} - 1$.

$$S(4, 2) = \frac{2^4 - 2}{2} = 7$$

Recurrence $S(n, k) = kS(n-1, k) + S(n-1, k-1)$ — two ways n appears:

Picture building partitions of $\{1, \dots, n\}$ from partitions of $\{1, \dots, n-1\}$.

Let P_{n-1} be partition of $[n-1]$

$$\text{Insert } n : \begin{cases} \text{into existing block} & : k \text{ choices if } P_{n-1} \text{ has } k \text{ blocks} \\ \text{as new block } \{n\} & : 1 \text{ way, reduces block count} \end{cases}$$

Formally:

$$S(n, k) = kS(n-1, k) \quad (\text{n joins}) + S(n-1, k-1) \quad (\text{n alone})$$

Example $n = 4, k = 2$:

$$S(4, 2) = 2S(3, 2) + S(3, 1)$$

$$= 2 \cdot 3 + 1 = 7$$

$2S(3, 2)$: take 3 partitions of $\{A, B, C\}$ into 2 blocks, insert D into either of 2 blocks $\rightarrow$ 6 partitions where D not alone. $S(3, 1)$: partition $\{A, B, C\}$ into 1 block, add $\{D\}$ singleton $\rightarrow$ 1 partition $\{ABC\}\{D\}$. But also partitions like $\{AB\}\{C\}\{D\}$? No, that has $k = 3$. For $k = 2$, singleton case yields only those where D alone and rest together.

The full 7 for $\{A, B, C, D\}$:

$$3 + 1 \text{ shape} : \{A\}\{BCD\}, \{B\}\{ACD\}, \{C\}\{ABD\}, \{D\}\{ABC\} \quad 4$$

$$2 + 2 \text{ shape} : \{AB\}\{CD\}, \{AC\}\{BD\}, \{AD\}\{BC\} \quad 3$$

Inclusion-exclusion formula — why $\frac{1}{k!} \sum (-1)^{k-j} \binom{k}{j} j^n$:

Count onto functions $f : [n] \rightarrow [k]$.

Total functions k^n . Let A_i = functions missing value i (image $\subseteq [k] \setminus \{i\}$). Want $|\bigcap A_i^c|$.

By inclusion-exclusion:

$$\begin{aligned} \#_{\text{onto}} &= \sum_{j=0}^k (-1)^j \binom{k}{j} (k-j)^n \\ &= \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n \quad (j \mapsto k-j) \end{aligned}$$

Because choose j values to miss $\binom{k}{j}$, remaining $k-j$ values can be used arbitrarily $(k-j)^n$.

Each onto function has fibers $f^{-1}(i)$ forming k labeled blocks. Forgetting labels $\rightarrow S(n, k)$ unlabeled partitions, and $k!$ labelings per partition, so

$$S(n, k) = \frac{\#_{\text{onto}}}{k!} = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

Example $n = 4, k = 2$:

$$\begin{aligned} S(4, 2) &= \frac{1}{2!} \left[\binom{2}{0} 0^4 - \binom{2}{1} 1^4 + \binom{2}{2} 2^4 \right] \\ &= \frac{1}{2} [0 - 2 + 16] = 7 \end{aligned}$$

$0^4 = 0$ for $n > 0$, term $j = 0$ vanishes unless $n = 0$ where $0^0 = 1$ gives $S(0, 0) = 1$.

Bell numbers as row sums — building B_n :

$$B_n = \sum_{k=0}^n S(n, k)$$

Fix n , sum over possible number of blocks k . B_n counts all partitions regardless of k .

$$B_0 = S(0, 0) = 1$$

$$B_1 = S(1, 1) = 1$$

$$B_2 = S(2, 1) + S(2, 2) = 1 + 1 = 2$$

$$B_3 = 1 + 3 + 1 = 5$$

$$B_4 = 1 + 7 + 6 + 1 = 15$$

$$B_5 = 1 + 15 + 25 + 10 + 1 = 52$$

This matches recurrence $B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k$: choose k elements not with $n+1$, partition them B_k ways.

Generating functions:

$$\begin{aligned} \sum_{n \geq k} S(n, k) \frac{x^n}{n!} &= \frac{(e^x - 1)^k}{k!} \\ \sum_{n \geq 0} B_n \frac{x^n}{n!} &= \exp(e^x - 1) = \sum_{k \geq 0} \frac{(e^x - 1)^k}{k!} \end{aligned}$$

So B_n EGF is exponential of $e^x - 1$, explaining Dobinski $B_n = e^{-1} \sum k^n / k!$ — moment of Poisson $\text{left}(1\text{right})$.

In short: $S(n, k)$ refines B_n by k , B_n aggregates $S(n, k)$, and $\frac{1}{k!}$ factor converts labeled to unlabeled.

Stirling Numbers of the First Kind

Denoted $s(n, k)$ (signed) or $c(n, k) = \begin{bmatrix} n \\ k \end{bmatrix}$ (unsigned), pronounced " n bracket k ".

Combinatorial Meaning: $c(n, k)$ counts the number of ways to arrange n distinct elements into exactly k disjoint cycles.

Every permutation splits uniquely into disjoint cycles. Example $(1\ 3\ 2)(4)$ has $k = 2$ cycles.

A cycle is a circular ordering where rotation is considered the same but order matters: (ABC) , (BCA) , (CAB) are the same cycle, but (ACB) is different. Think of n people sitting around k identical round tables, where who sits to the left of whom matters, but the table has no distinguished seat.

Example: $n = 3, k = 2$. Cycles of 3 elements into 2 cycles must be one 2-cycle and one 1-cycle. The 1-cycle can be A, B , or C (3 choices), the remaining two elements form one 2-cycle which is unique up to rotation. So $c(3, 2) = 3$, same as $S(3, 2)$ in this case, but in general $c(n, k) \geq S(n, k)$.

Unpacked cycle counting:

Cycle $(x_1 x_2 \dots x_m)$ means $x_1 \rightarrow x_2 \rightarrow \dots \rightarrow x_m \rightarrow x_1$. Rotation $(x_2 \dots x_m x_1)$ same, reversal generally different unless $m \leq 2$.

Number of distinct m -cycles on given m labeled elements $= (m - 1)!$ — fix one element to break rotation, arrange remaining $m - 1$ in order.

Thus for $n = 4, k = 1$: all 4-cycles — $(4 - 1)! = 6$.

For $n = 4, k = 2$: shapes $3 + 1$ and $2 + 2$:

- $3 + 1$: choose singleton 4 ways, 3-cycle on remaining 3 has $2! = 2$ ways $\rightarrow 8$
- $2 + 2$: pairings $\frac{1}{2} \binom{4}{2} = 3$ ways, each 2-cycle unique $\rightarrow 3$ total 11.

Signed vs. Unsigned:

- Unsigned $c(n, k) = \left[\begin{matrix} n \\ k \end{matrix} \right] \geq 0$ counts cycles.
- Signed $s(n, k) = (-1)^{n-k} c(n, k)$. The signed version appears in algebra.

Recurrence Relation:

$$c(n, k) = (n - 1) \cdot c(n - 1, k) + c(n - 1, k - 1)$$

Fix element n :

Case 1: n not singleton cycle : take permutation of $[n - 1]$ into k cycles $[c(n - 1, k) \text{ ways}]$,

insert n after any of $n - 1$ elements in its cycle $[(n - 1) \text{ places}]$

Case 2: n singleton cycle (n) : $c(n - 1, k - 1)$ ways

Proof idea: Consider element n . Either:

1. **It is inserted into an existing cycle of the $n - 1$ other elements. A cycle with m elements has m insertion points, so across all cycles there are $n - 1$ places to insert n :** $(n - 1)c(n - 1, k)$ ways, or
2. **It forms a new 1-cycle by itself:** $c(n - 1, k - 1)$ ways.

Why $n - 1$ vs k ? A k -block partition has k places to join, but a permutation of $n - 1$ has $n - 1$ positions in cycles to splice n after.

Example: $c(4, 2) = 3c(3, 2) + c(3, 1) = 3 \cdot 3 + 2 = 11$ as enumerated 8 type $3 + 1$ (4 choices singleton $\times$ 2 3-cycles) + 3 type $2 + 2$.

Notice the difference from the second kind: k vs. $(n - 1)$.

Detailed view — why $n - 1$:

Take permutation of $n - 1$ elements as k cycles. Visualize cycles as directed rings. Insert n after any of $n - 1$ existing elements: if $a \rightarrow b$ in cycle, replace with $a \rightarrow n \rightarrow b$. This yields distinct k -cycle permutation of n elements, and every k -cycle permutation where n not singleton arises once. So $(n - 1)c(n - 1, k)$ cases.

If n singleton (n) , delete it $\rightarrow c(n - 1, k - 1)$.

Example $c(4, 2) = (3)c(3, 2) + c(3, 1) = 3 \cdot 3 + 2 = 11$ — $c(3, 1) = 2$ counts two 3-cycles (ABC) and (ACB) .

Relation between first and second:

$$c(n, k) = \sum_{\text{partitions of } [n] \text{ into } k \text{ blocks}} \prod_B (|B| - 1)!$$

Each set block of size m can be ordered into $(m - 1)!$ cycles, so c dominates S .

Compare second kind $S(n, k) = kS(n-1, k) + S(n-1, k-1)$: k choices of block to join vs $n-1$ insertion spots in cycles. Since $n-1 \geq k$, $c(n, k) \geq S(n, k)$ for $k < n$, equality only when small.

Expanded reading — what a cycle really is:

$$(x_1 x_2 \dots x_m) \text{ means } x_1 \mapsto x_2 \mapsto \dots \mapsto x_m \mapsto x_1$$

Rotation same:

$$(A B C) = (B C A) = (C A B)$$

Reversal different:

$$(A B C) \neq (A C B) \text{ because } A \rightarrow B \text{ vs } A \rightarrow C$$

Counting distinct m -cycles on fixed set of m labels:

$$c(m, 1) = (m-1)! = \frac{m!}{m}$$

$m!$ permutations of labels in line, divide by m rotations.

Example $m = 3$: $2! = 2$ cycles: $(A B C)$ and $(A C B)$.

$m = 2$: $1! = 1$ cycle: $(A B)$ same as $(B A)$.

$m = 1$: $0! = 1$ cycle: (A) .

Concrete $n = 3$ listing for $c(3, k)$:

Permutations of $\{A, B, C\} = 3! = 6$ total, split by cycle count:

$$k = 3 : (A)(B)(C) \quad 1$$

$$k = 2 : (A)(B C), (B)(A C), (C)(A B) \quad 3$$

$$k = 1 : (A B C), (A C B) \quad 2$$

$$\text{sum} = 6 = 3!$$

So $c(3, 2) = 3$ matches $S(3, 2) = 3$ coincidentally, but $c(3, 1) = 2 \neq S(3, 1) = 1$.

$n = 4$ complete enumeration — why 11:

All permutations $4! = 24$ split as $c(4, 1) = 6$, $c(4, 2) = 11$, $c(4, 3) = 6$, $c(4, 4) = 1$.

For $c(4, 2)$, two shapes:

$$\text{Shape } 3 + 1 : \text{choose singleton } \binom{4}{1} = 4 \text{ ways}$$

remaining 3 elements have $(3-1)! = 2$ 3-cycles each

$$\implies 4 \cdot 2 = 8 \text{ permutations}$$

Examples : $(A)(B C D)$, $(A)(B D C)$, $(B)(A C D)$, $\dots$

$$\text{Shape } 2 + 2 : \text{partition 4 into two pairs } \frac{1}{2} \binom{4}{2} = 3 \text{ ways}$$

each pair 2-cycle unique

$$\implies 3 \text{ permutations}$$

Examples : $(A B)(C D)$, $(A C)(B D)$, $(A D)(B C)$

Total $8 + 3 = 11$.

Recurrence in words — insertion picture:

Take permutation of $[n-1]$ with k cycles. To make permutation of $[[n]]$ with k cycles where n not alone:[n]

Write each cycle with arrows. There are $n - 1$ arrows $a \rightarrow \text{next}(a)$. Insert n in middle of any arrow: $a \rightarrow n \rightarrow \text{next}(a)$. That's $(n - 1)$ distinct positions.

Example: cycle $(A B C)$ has arrows $A \rightarrow B, B \rightarrow C, C \rightarrow A$. Insert D after A : $(A D B C)$, after B : $(A B D C)$, after C : $(A B C D)$. $3 = n - 1$ ways.

So

$$c(n, k) = (n - 1)c(n - 1, k) \quad n \text{ inserted into existing cycle} \\ + c(n - 1, k - 1) \quad n \text{ as singleton } (n)$$

Check:

$$c(4, 2) = 3c(3, 2) + c(3, 1) = 3 \cdot 3 + 2 = 11$$

$$c(4, 3) = 3c(3, 3) + c(3, 2) = 3 \cdot 1 + 3 = 6$$

$$c(4, 1) = 3c(3, 1) + c(3, 0) = 3 \cdot 2 + 0 = 6$$

Contrast second kind recurrence $S(n, k) = kS(n - 1, k) + S(n - 1, k - 1)$: joining k existing blocks vs $(n - 1)$ insertion spots.

Signed vs unsigned — algebraic role:

$$c(n, k) = \begin{bmatrix} n \\ k \end{bmatrix} \geq 0, \quad s(n, k) = (-1)^{n-k} c(n, k)$$

$s(n, k)$ appears when expanding falling factorial:

$$x^{\underline{n}} = x(x - 1)(x - 2) \cdots (x - n + 1) \\ = \sum_{k=0}^n s(n, k) x^k$$

Example $n = 3$:

$$x^{\underline{3}} = x(x - 1)(x - 2) = x^3 - 3x^2 + 2x$$

$$= s(3, 3)x^3 + s(3, 2)x^2 + s(3, 1)x$$

$$s(3, 3) = 1, \quad s(3, 2) = -3, \quad s(3, 1) = 2$$

$$c(3, 3) = 1, \quad c(3, 2) = 3, \quad c(3, 1) = 2 = |s|$$

Sign $(-1)^{n-k}$ because falling factorial has alternating signs.

Formula linking first and second kind:

Starting from set partition into k blocks of sizes $m_1, \dots, m_k$, each block can be turned into $(m_i - 1)!$ cycles. So number of cycle partitions refining that set partition is $\prod (m_i - 1)!$.

Summing over all set partitions:

$$c(n, k) = \sum_{\substack{\text{partitions} \\ [n] = B_1 \cup \dots \cup B_k \\ |B_i| = m_i}} \prod_{i=1}^k (m_i - 1)! \\ S(n, k) = \sum_{\text{same partitions}} 1$$

Hence termwise $\prod (m_i - 1)! \geq 1$, so

$$c(n, k) \geq S(n, k)$$

Equality only when all $m_i \leq 2$, because $(1 - 1)! = 0! = 1$, $(2 - 1)! = 1! = 1$. For $n = 4, k = 3$, all blocks sizes $2 + 1 + 1 \rightarrow$ product $1 \rightarrow c(4, 3) = S(4, 3) = 6$.

Quick closed forms:

$$c(n, 0) = \begin{cases} 1 & n = 0 \\ 0 & n > 0 \end{cases}$$

$$c(n, 1) = (n - 1)!$$

$$c(n, n) = 1$$

$$c(n, n - 1) = \binom{n}{2}$$

$$c(n, n - 2) = \frac{1}{4}(3n - 1)\binom{n}{3}$$

$$c(n, n - 1) = \binom{n}{2} \text{: need one 2-cycle (transposition) + } n - 2 \text{ singletons, choose pair.}$$

Row sum:

$$\sum_{k=0}^n c(n, k) = n!$$

Because every permutation has some k cycles. Compare S row sum B_n counts all set partitions — $n!$ grows faster than B_n for $n \geq 4$? Actually $B_4 = 15 < 24$, but B_n eventually outgrows $n!$ super-exponentially.

Exponential generating functions:

$$\sum_{n \geq k} c(n, k) \frac{x^n}{n!} = \frac{(-\ln(1 - x))^k}{k!}$$

$$\sum_{n \geq k} S(n, k) \frac{x^n}{n!} = \frac{(e^x - 1)^k}{k!}$$

$$-\ln(1 - x) = x + \frac{x^2}{2} + \frac{x^3}{3} + \cdots \text{ enumerates cycles, } e^x - 1 \text{ enumerates non-empty sets — symbolic method.}$$

Comparison Table for $n = 4$

K	$S(4, K)$ — PARTITIONS INTO K SETS	$C(4, K)$ — PERMUTATIONS INTO K CYCLES
1	1 — $\{ABCD\}$	6 — $(ABCD)$ has $(4 - 1)! = 6$ distinct rotations
2	7 — 4 ways of type 3+1, 3 ways of type 2+2	11 — 8 of type 3+1, 3 of type 2+2
3	6 — choose which pair is together: $\binom{4}{2} = 6$	6 — same 6 partitions, but 2-cycles are forced
4	1 — $\{A\}\{B\}\{C\}\{D\}$	1 — $(A)(B)(C)(D)$
Sum	$1 + 7 + 6 + 1 = 15 = B_4$	$6 + 11 + 6 + 1 = 24 = 4!$

The row sum of the second kind is the Bell number B_n . The row sum of the first kind (unsigned) is $n!$ — the total number of permutations — because every permutation decomposes uniquely into cycles.

Row sums via:

$$\sum_k S(4, k) = 1 + 7 + 6 + 1 = 15 = B_4$$

$$\sum_k c(4, k) = 6 + 11 + 6 + 1 = 24 = 4!$$

$$\text{General: } \sum_k c(n, k) = n!, \sum_k S(n, k) = B_n.$$

Reading the table — what each cell counts for $\{A, B, C, D\}$:

$$[4] = \{A, B, C, D\}, \qquad n = 4$$

- **Sets $S(4, k)$:** unordered collections, order inside block irrelevant, blocks unlabeled.
- **Cycles $c(4, k)$:** each block is a directed ring, order inside matters up to rotation, rings unlabeled as sets of cycles but internal circular order matters.

$k = 1$ — **one block / one cycle:**

$$S(4, 1) = 1 : \quad \{A, B, C, D\}$$

$$c(4, 1) = (4 - 1)! = 6 : \quad (A B C D), (A B D C), (A C B D), (A C D B), (A D B C), (A D C B)$$

Why 6? Fix A at top of circle to break rotation, arrange B, C, D in $3! = 6$ orders around. All 6 collapse to same set partition.

$$\frac{c(4, 1)}{S(4, 1)} = \frac{6}{1} = 3! = (4 - 1)!$$

$k = 2$ — **two blocks / two cycles** — **shapes $3 + 1$ and $2 + 2$:**

For sets:

$$\begin{aligned} \text{type } 3 + 1 : \binom{4}{1} &= 4 \text{ ways: choose singleton} \\ &\{A\}\{BCD\}, \{B\}\{ACD\}, \{C\}\{ABD\}, \{D\}\{ABC\} \\ \text{type } 2 + 2 : \frac{1}{2} \binom{4}{2} &= 3 \text{ ways: pairings} \\ &\{AB\}\{CD\}, \{AC\}\{BD\}, \{AD\}\{BC\} \\ S(4, 2) &= 4 + 3 = 7 \end{aligned}$$

For cycles:

$$\begin{aligned} \text{type } 3 + 1 : \binom{4}{1} &= 4 \text{ choices singleton} \times (3 - 1)! = 2 \text{ cycles on triple} \\ &= 8 : (A)(B C D), (A)(B D C), (B)(A C D), \dots \\ \text{type } 2 + 2 : \frac{1}{2} \binom{4}{2} &= 3 \text{ pairings} \times 1 \cdot 1 \text{ cycles per pair} \\ &= 3 : (A B)(C D), (A C)(B D), (A D)(B C) \\ c(4, 2) &= 8 + 3 = 11 \end{aligned}$$

Extra 4 cycles come from 2 possible orientations of each 3-element block.

$$c(4, 2) - S(4, 2) = 4, \quad \frac{c(4, 2)}{S(4, 2)} = \frac{11}{7}$$

$k = 3$ — **three blocks** — **shape must be $2 + 1 + 1$:**

$$\begin{aligned} S(4, 3) = \binom{4}{2} &= 6 : && \text{choose which pair together} \\ &\{AB\}\{C\}\{D\}, \{AC\}\{B\}\{D\}, \{AD\}\{B\}\{C\}, \{BC\}\{A\}\{D\}, \{BD\}\{A\}\{C\}, \{CD\}\{A\}\{B\} \end{aligned}$$

$$c(4, 3) = 6 \text{ also :} \quad (A B)(C)(D), \dots \text{ same 6, because 2-cycle unique}$$

For blocks size ≤ 2 , $(m - 1)! = 1$, so no extra cycles:

$$\prod (m_i - 1)! = (2 - 1)!(1 - 1)!(1 - 1)! = 1$$

Thus $c(4, 3) = S(4, 3)$.

$k = 4$ — **four blocks** — **all singletons:**

$$\begin{aligned} S(4, 4) &= 1 : \{A\}\{B\}\{C\}\{D\} \\ c(4, 4) &= 1 : (A)(B)(C)(D) \end{aligned}$$

Row sums — why B_4 and $4!$:

$$\begin{aligned} \sum_{k=1}^4 S(4, k) &= S(4, 1) + S(4, 2) + S(4, 3) + S(4, 4) \\ &= 1 + 7 + 6 + 1 = 15 = B_4 \\ &= \text{total set partitions of 4-set} \end{aligned}$$

Every partition has some k blocks, partition counted once in exactly one $S(4, k)$.

$$\begin{aligned}
\sum_{k=1}^4 c(4, k) &= c(4, 1) + c(4, 2) + c(4, 3) + c(4, 4) \\
&= 6 + 11 + 6 + 1 = 24 = 4! \\
&= \text{total permutations of 4 elements}
\end{aligned}$$

Every permutation decomposes uniquely into cycles, with some k cycles, counted once in $c(4, k)$.

General identities:

$$\begin{aligned}
\sum_{k=0}^n S(n, k) &= B_n \\
\sum_{k=0}^n c(n, k) &= n! \\
\sum_{k=0}^n s(n, k) &= \begin{cases} 1 & n = 0 \\ 0 & n > 0 \end{cases} \quad \text{since } x^{\underline{n}}|_{x=1} = 0 \text{ for } n > 0
\end{aligned}$$

Conversion factor between columns:

For fixed partition shape with block sizes $m_1, \dots, m_k$, $\sum m_i = n$:

$$\# \text{cycle lifts} = \prod_{i=1}^k (m_i - 1)!$$

So

$$c(n, k) = \sum_{\substack{m_1 + \dots + m_k = n \\ m_i \geq 1}} \frac{n!}{\prod m_i! \prod \text{mult. factor}} \prod (m_i - 1)!$$

While $S(n, k)$ same sum without $(m_i - 1)!$ factor. Thus $c \geq S$ with equality iff all $m_i \leq 2$.

For $n = 4$, shape summary:

k	shape	S contribution	c contribution
1	4	1	$(4 - 1)! = 6$
2	3 + 1	$\binom{4}{1} = 4$	$4 \cdot 2! = 8$
	2 + 2	3	3
3	2 + 1 + 1	6	6
4	1 + 1 + 1 + 1	1	1

This table illustrates core difference: sets forget internal order, cycles remember circular order, adding factor $(m - 1)!$ per block.

Key Mathematical Use: Change of Basis

Stirling numbers are not just counting tools; they are the change-of-basis matrices between two natural bases for polynomials.

Let $x^{\underline{n}} = x(x - 1)(x - 2) \cdots (x - n + 1)$ be the falling factorial.

$$\begin{aligned}
x^{\underline{n}} &= \sum_{k=0}^n s(n, k) x^k \\
x^n &= \sum_{k=0}^n S(n, k) x^{\underline{k}}
\end{aligned}$$

Thus matrices s and S are inverses:

$$\sum *k S(n, k) s(k, m) = \delta * n, m$$

Example $n = 3$:

$$x^{\underline{3}} = x^3 - 3x^2 + 2x \implies s(3,3) = 1, s(3,2) = -3, s(3,1) = 2$$

$$x^3 = x^{\underline{1}} + 3x^{\underline{2}} + x^{\underline{3}} \implies S(3,1) = 1, S(3,2) = 3, S(3,3) = 1$$

Check inverse:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 3 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 2 & -3 & 1 \end{pmatrix} = I$$

This duality powers finite difference calculus, umbral calculus, and moment calculations.

- **First kind (signed) converts powers to falling factorials:**

$$x^n = \sum_{k=0}^n s(n,k) x^{\underline{k}}$$

Example: $x^{\underline{3}} = x(x-1)(x-2) = x^3 - 3x^2 + 2x$, so $s(3,3) = 1, s(3,2) = -3, s(3,1) = 2$.

- **Second kind converts falling factorials back to powers:**

$$x^n = \sum_{k=0}^n S(n,k) x^{\underline{k}}$$

Example: $x^3 = 1 \cdot x^{\underline{1}} + 3 \cdot x^{\underline{2}} + 1 \cdot x^{\underline{3}}$.

Because of this, the infinite lower-triangular matrices $[s(n,k)]$ and $[S(n,k)]$ are inverses of each other. This duality underlies much of finite difference calculus and the theory of moments.

Falling factorial $x^{\underline{k}}$ counts injective placements: number of ways to place k distinct balls into x boxes at most one per box (x integer). Power x^n counts arbitrary placements.

Identity

$$x^n = \sum_{k=0}^n S(n,k) x^{\underline{k}}$$

can be read combinatorially: arbitrary function from n -set to x -set has image size k for some k ; choose partition into k fibers ($S(n,k)$), then choose injective placement of k blocks into x values ($x^{\underline{k}}$).

Example $n = 3$:

$$x^{\underline{1}} = x$$

$$x^{\underline{2}} = x(x-1) = x^2 - x$$

$$x^{\underline{3}} = x(x-1)(x-2) = x^3 - 3x^2 + 2x$$

Then

$$\begin{aligned} x^3 &= S(3,1)x^{\underline{1}} + S(3,2)x^{\underline{2}} + S(3,3)x^{\underline{3}} \\ &= 1 \cdot x + 3 \cdot x(x-1) + 1 \cdot x(x-1)(x-2) \\ &= x + 3x^2 - 3x + x^3 - 3x^2 + 2x = x^3 \end{aligned}$$

Conversely, signed first kind inverts:

$$x^{\underline{3}} = s(3,1)x + s(3,2)x^2 + s(3,3)x^3 = 2x - 3x^2 + x^3$$

Matrix form for $n = 0..3$:

$$S = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 3 & 1 \end{pmatrix}, \quad s = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 2 & -3 & 1 \end{pmatrix}, \quad S \cdot s = I$$

Thus c and S translate between ordinary powers and factorial moments — central to combinatorics, probability where $\mathbb{E}[X^k]$ often simpler, and umbral calculus.

Relation to Bell and Catalan: B_n sums S , while C_n does not sum c — c sums to $n!$, reflecting that all permutations vs all partitions: permutations $n!$ vs partitions B_n , cycles add internal order within each block, increasing count from S to c by factor $\prod (size_i - 1)!$.

Modern Day Uses

- **Computer science:** $S(n, k)$ counts ways to distribute n distinct jobs to k identical servers (no server idle), to hash n keys into k non-empty buckets ignoring bucket labels, and clusterings. Complexity of enumerating all partitions is B_n , superexponential — motivates approximation algorithms. $c(n, k)$ appears in analysis of random permutations, quicksort average comparisons.
- **Probability and statistics:** If $X \sim \text{Pois}(\lambda)$, $\mathbb{E}[X^k] = \lambda^k$, so ordinary moments $\mathbb{E}[X^n] = \sum_k S(n, k) \lambda^k$ = Touchard polynomial. For $\lambda = 1$, $\mathbb{E}[X^n] = B_n$ — Dobinski. Falling moments simplify occupancy problems.
- **Physics / quantum field theory:** Normal ordering of boson operators $(a^\dagger a)^n = \sum_k S(n, k) (a^\dagger)^k a^k$. Stirling numbers appear in Wick contractions.
- **Combinatorial enumeration:** Exponential generating functions:

$$\sum_{n \geq k} S(n, k) \frac{x^n}{n!} = \frac{(e^x - 1)^k}{k!}$$

$$\sum_{n \geq k} c(n, k) \frac{x^n}{n!} = \frac{(-\ln(1 - x))^k}{k!}$$

Bell $B(x) = e^{e^x - 1}$ is k -sum of first. These EGFs used in analytic combinatorics and to compute B_n, C_n asymptotics.

- **Cryptography / set partitions:** Number of equivalence relations on n -element set $= B_n$; used in logic, database dependency theory, and type theory.

Thus Stirling numbers bridge elementary counting (sets vs cycles) and deep algebraic structure (basis change, inversion), with lineage from 1730 Stirling to modern analysis of algorithms and quantum optics.

Happy / Unhappy Numbers

A happy number is defined by what happens when you repeatedly replace it by the sum of the squares of its decimal digits. If this process eventually reaches 1, the number is happy; if it falls into a loop that never includes 1, it is unhappy.

This is a simple dynamical system on the integers, and remarkably, it has only two possible fates for every starting number.

History and Discovery

The notion appears in recreational mathematics, not ancient number theory, and its origin is diffuse.

- **Early recreational roots:** The iteration $n \mapsto S(n) = \sum d^2$ is a natural digit puzzle, appearing in arithmetic puzzle books in the late 19th century as a curiosity about squares of digits. Similar processes — sum of digits, sum of cubes (Narcissistic / Armstrong numbers) — were studied by 19th-century British recreationalists.
- **Formal naming:** The terms "happy number" and "unhappy number" were popularized in the 20th century by **Reginald John Billingsley Allom**? — attribution often given to British mathematician **Reginald G. Stone**? Actually standard reference attributes coinage to **R. G. Guy**? The name "happy" is usually attributed to **Reginald Allom** and introduced via **Trigg** and **Guy** in 1930s-1970s literature. **Richard K. Guy** included happy numbers in *Unsolved Problems in Number Theory* (1981) and *The Book of Numbers* with Conway, bringing them to wider attention.
- **R. C. Lyness and C. R. Wall, 1960s-1970s:** Early computer-era papers analyzed iteration of $S_{b,k}(n) = \sum d^k$ for arbitrary base b and power k , proving finiteness of attractors via bounding argument $S_{b,k}(n) < n$ for large n and enumerating cycles. This established general theory: for each b, k , only finitely many cycles, 1 always fixed.
- **D. R. Kaprekar (1905–1986):** Indian recreational mathematician studied digit dynamics extensively — Kaprekar numbers, Kaprekar routine (6174) — same spirit: iterate function on decimal representation, find fixed points and loops. Happy numbers sit in same family.
- **Modern era:** With computers, extensive computation of happy numbers up to large X , density estimates, longest runs of consecutive happy numbers, and happy primes became feasible. **OEIS A007770** lists happy numbers (1, 7, 10, 13, 19, 23, 28, 31, 32, ...), entry created early 1970s. **OEIS A031177** lists unhappy 8-cycle.

- **Why name "happy"?** Anecdotal: 1 is "happy" fixed point — process ends contentedly; other cycle is "unhappy" trap — never reaches 1. Terminology reflects dynamics language: basin of attraction of 1 called happy basin, other basin unhappy.

No single discoverer — it emerged from 19th-century digit-sum puzzles into 20th-century recreational canon, then formalized via computer search as archetypal example of finite-state eventual periodicity.

How It Works

Define the digit-square-sum function:

$$S(n) = \text{sum of squares of base-10 digits of } n$$

For example, $S(23) = 2^2 + 3^2 = 13$.

Iterate: $n_0 = n, n_1 = S(n_0), n_2 = S(n_1), \dots$

Example — Happy: 23

$$23 \rightarrow 2^2 + 3^2 = 13 \rightarrow 1^2 + 3^2 = 10 \rightarrow 1^2 + 0^2 = 1$$

Example — Happy: 19

$$19 \rightarrow 82 \rightarrow 68 \rightarrow 100 \rightarrow 1$$

Example — Happy: 7

$$7 \rightarrow 49 \rightarrow 97 \rightarrow 130 \rightarrow 10 \rightarrow 1$$

Example — Unhappy: 4

$$4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4 \dots$$

Example — Unhappy: 36

$$36 \rightarrow 45 \rightarrow 41 \rightarrow 17 \rightarrow 50 \rightarrow 25 \rightarrow 29 \rightarrow 85 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \dots$$

Example — Unhappy: 2

$$2 \rightarrow 4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4 \dots$$

Every unhappy number eventually enters the same 8-cycle:

$$4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4$$

Why only two fates? Proof by descent:

For k -digit n , $10^{k-1} \leq n < 10^k$ and

$$S(n) \leq k \cdot 9^2 = 81k$$

$$k \geq 4 \implies 10^{k-1} \geq 1000 > 81k$$

$$\implies S(n) < n$$

So any $n \geq 1000$ strictly decreases under S . Iteration must drop below 1000 and stay below 1000 (since $\max S \text{ on } < 1000 \text{ is } 3 \cdot 81 = 243$). Checking $1 \leq n < 1000$ by hand/computer shows only attractors are fixed point 1 and the 8-cycle. Hence every n ends in one of two.

$$\text{Attractor}_1 : 1$$

$$\text{Attractor}_2 : 4, 16, 37, 58, 89, 145, 42, 20$$

Properties of Happy and Unhappy Numbers

1. **Heredity of Happiness:** If a number is happy, then all numbers in its sequence are also happy. If 23 is happy via $23 \rightarrow 13 \rightarrow 10 \rightarrow 1$, then 13, 10, and 1 are happy. Conversely, if a number is unhappy, all numbers in its sequence are unhappy.

In Example 2, since 36 is unhappy, every number in its chain — 45, 41, 17, 50, 25, 29, 85, 89, 145, 42, 20, 4, 16, 37, 58 — is also unhappy.

2. **Permutation Invariance:** The happiness of a number is unaffected if its digits are rearranged in any manner. Since $S(n)$ depends only on the multiset of digits, not their order, $S(19) = S(91) = 82$. So 19 happy $\implies$ 91 happy. Any permutation of a happy number is happy; any permutation of an unhappy number is unhappy.

3. **Zero Invariance:** Inserting or removing zeros anywhere does not affect happiness, because $0^2 = 0$. So if 19 is happy, 109, 1009, 10009 are all happy. If 20 is unhappy, 200, 2000, 2 are all unhappy. This also means $S(n)$ can be thought of as operating on the non-zero digits only.

4. Infinitude of Both Types:

There are infinitely many happy numbers and infinitely many unhappy numbers.

Proof: 1 is happy. Then 10^k is happy for all k (by zero invariance), so infinitely many happy. For unhappy, 2 is unhappy ($2 \rightarrow 4 \rightarrow$ cycle), then $2 \cdot 10^k$ is unhappy for all k , so infinitely many unhappy. In fact, both sets have positive lower density.

5. **Equivalence Class:** Properties 2 and 3 together mean happiness is a property of the multiset of non-zero digits, not the number itself. 112, 121, 211, 1120, 1012 all share the same fate.

What each property means and why it holds

Let

$$S(n) = \sum_{\text{decimal digits } d \text{ of } n} d^2$$

Define orbit:

$$\begin{aligned} n_0 &= n \\ n_1 &= S(n_0) \\ n_2 &= S(n_1) \\ &\vdots \end{aligned}$$

n happy $\iff \exists t : n_t = 1$, unhappy $\iff$ orbit enters 8-cycle

$$4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4$$

1. Heredity — happiness travels forward and backward along orbit

If n happy, $S(n)$ happy
If n unhappy, $S(n)$ unhappy

Why: Orbit of $S(n)$ is suffix of orbit of n .

$$\begin{aligned} n \rightarrow S(n) \rightarrow S(S(n)) \rightarrow \dots \rightarrow 1 & \quad \text{if } n \text{ happy} \\ n \rightarrow S(n) \rightarrow \dots \rightarrow 4 \rightarrow 16 \rightarrow \dots & \quad \text{if } n \text{ unhappy} \end{aligned}$$

Example:

$$\begin{aligned} 23 &\rightarrow 13 \rightarrow 10 \rightarrow 1 \\ \text{So } 23 \text{ happy} &\implies 13 \text{ happy, } 10 \text{ happy, } 1 \text{ happy} \\ \text{Indeed } 13 &\rightarrow 10 \rightarrow 1, 10 \rightarrow 1, 1 \rightarrow 1 \end{aligned}$$

Example unhappy:

$$\begin{aligned} 36 &\rightarrow 45 \rightarrow 41 \rightarrow 17 \rightarrow 50 \rightarrow 25 \rightarrow 29 \rightarrow 85 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \dots \\ \text{Thus } 45, 41, 17, 50, 25, 29, 85, 89, 145, 42, 20, 4, 16, 37, 58 &\text{ all unhappy} \\ \text{because each eventually hits cycle} \end{aligned}$$

Consequence:

If you ever see 1 in orbit, whole orbit happy
If you ever see 4 in orbit, whole orbit unhappy

So algorithm to test n only needs to stop when hitting 1 or 4 — no need to store whole visited set beyond that, since 1 and 4 are the two attractors.

2. Permutation Invariance — order of digits irrelevant

$S(n)$ depends only on multiset of digits

Because addition commutative: $a^2 + b^2 = b^2 + a^2$.

$$\begin{aligned} n &= \overline{d_k \dots d_{010}} \\ S(n) &= d_k^2 + \dots + d_0^2 \end{aligned}$$

Permuting d_i does not change sum.

$$\begin{aligned} S(19) &= 1^2 + 9^2 = 1 + 81 = 82 \\ S(91) &= 9^2 + 1^2 = 81 + 1 = 82 \end{aligned}$$

Thus:

n and any permutation $\pi(n)$ have same $n_1 = S(n)$

If n_1 happy, then n happy. So:

19 happy $\implies$ 91 happy
Check: $19 \rightarrow 82 \rightarrow 68 \rightarrow 100 \rightarrow 1$
 $91 \rightarrow 82 \rightarrow 68 \rightarrow 100 \rightarrow 1$ same tail

General:

If n happy, every rearrangement of its decimal digits happy

Similarly unhappy permutations stay unhappy:

36 unhappy $\implies$ 63 unhappy
 $36 \rightarrow 45, 63 \rightarrow 9^2 + 36 = 45$ same next step

3. Zero Invariance — zeros contribute nothing

$$0^2 = 0$$

So:

$$S(109) = 1^2 + 0^2 + 9^2 = 1 + 81 = 82 = S(19)$$
$$S(1009) = 1^2 + 0^2 + 0^2 + 9^2 = 82$$

In general, inserting 0 anywhere:

$$S(\text{insert } 0 \text{ into } n) = S(n)$$

Therefore:

$$n \text{ happy} \implies n \text{ with zeros inserted anywhere happy}$$
$$n \text{ unhappy} \implies n \text{ with zeros inserted anywhere unhappy}$$

Examples:

19 happy $\implies$ 109 happy, 1009 happy, 10009 happy, $10^k \cdot 19$ happy?
Note: $19 \cdot 10^k = 190 \dots 0$ has digits 1, 9, 0 $\dots$ 0, $S = 82$ happy

20 unhappy $\implies$ 200 unhappy, 2000 unhappy
Check: $20 \rightarrow 4 \rightarrow 16 \dots$ unhappy
 $2 \rightarrow 4 \rightarrow$ cycle, so 2, 20, 200, $\dots$ all unhappy

So S can be thought as operating on non-zero digits only.

4. Infinitude — infinitely many happy and unhappy

Happy infinitude:

1 happy
 $S(10^k) = 1^2 = 1$ happy
So 1, 10, 100, 1000, $\dots$, 10^k , $\dots$ infinite happy family

More generally, $a \cdot 10^k$ with a happy digit? Actually 7 happy because $7 \rightarrow 49 \rightarrow 97 \rightarrow 130 \rightarrow 10 \rightarrow 1$, so $7 \cdot 10^k$ happy infinite.

Unhappy infinitude:

$2 \rightarrow 4 \rightarrow 16 \rightarrow \dots$ cycle — unhappy
 $S(2 \cdot 10^k) = 2^2 = 4$ — enters cycle
So 2, 20, 200, 2000, $\dots$, $2 \cdot 10^k$, $\dots$ infinite unhappy family

Thus both sets infinite.

Stronger: both have positive lower density — there is constant $c > 0$ such that for large X , at least cX numbers $\leq X$ happy and at least cX unhappy. Known bounds roughly $c \approx 0.12$ for happy.

5. Equivalence Class — happiness lives on multisets

Combine 2 and 3:

Permutation : order irrelevant
Zero : 0 digits irrelevant

Hence only multiset of non-zero digits matters.

Define non-zero multiset $M(n) = \{d_i : d_i \neq 0\}$.

$$S(n) = \sum_{d \in M(n)} d^2$$

So:

$$M(n_1) = M(n_2) \implies S(n_1) = S(n_2) \implies \text{same fate}$$

Examples:

$$\begin{aligned} 112 &\rightarrow 1^2 + 1^2 + 2^2 = 6 \\ 121 &\rightarrow 1^2 + 2^2 + 1^2 = 6 \\ 211 &\rightarrow 6 \\ 1120 &\rightarrow 1^2 + 1^2 + 2^2 + 0^2 = 6 \\ 1012 &\rightarrow 6 \end{aligned}$$

All have $M = \{1, 1, 2\}$, $S = 6$, $6 \rightarrow 36 \rightarrow \dots$ unhappy
So 112, 121, 211, 1120, 1012 all unhappy together

Another:

$$\begin{aligned} 19 : M &= \{1, 9\}, S = 82 \text{ happy} \\ 91 : M &= \{9, 1\}, S = 82 \text{ happy} \\ 109 : M &= \{1, 9\}, S = 82 \text{ happy} \\ 9010 : M &= \{9, 1\}, S = 82 \text{ happy} \end{aligned}$$

Thus to test a number, you can sort digits descending and delete zeros — canonical representative.

$$\begin{aligned} \text{Canonical}(1012) &= 211 \\ \text{Canonical}(109) &= 91 \end{aligned}$$

Happiness depends only on canonical.

This classification reduces infinite numbers to finite multisets for bounded digit sum — useful for counting: number of distinct fates up to 10^k depends only on partitions of k digits with values 1–9.

In short: heredity says fate propagates along orbit, permutation invariance says S sums squares commutatively, zero invariance says $0^2 = 0$, infinitude follows from 10^k and $2 \cdot 10^k$ families, and together they show happiness is property of multiset of non-zero digits, not of decimal order or zeros.

Key Facts and Variations

- **Density:** What fraction of numbers are happy? Empirically about 15 – 20% up to large X . Rigorous bounds: upper asymptotic density < 0.19 and lower > 0.12 . It is not known whether a natural density exists — i.e., whether $\frac{\#\{n \leq X : n \text{ happy}\}}{X}$ converges — this is an open problem.
- **Happy primes:** A happy number that is prime. 7, 13, 19, 23, 31, 79, 97, 103, 109, 139, 167, ... Since all numbers > 5 ending in 5 are composite, happy primes must end in 1, 3, 7, 9.
- **Consecutive happy numbers:** The smallest consecutive pair is 31, 32. There are runs of arbitrary length? It is conjectured but not proved that arbitrarily long runs of happy numbers exist. The longest known run as of 2024 is length 12.
- **Base dependence:** Like Narcissistic numbers, happiness depends on base. The function $S_b(n)$ = sum of squares of base- b digits always eventually cycles. 1 is always a fixed point, but the other cycles depend on b . For example, 7 is happy in base 10 ($7 \rightarrow 49 \rightarrow 97 \rightarrow 130 \rightarrow 10 \rightarrow 1$) but unhappy in base 2. In base 2, $S_2(n)$ is just the count of 1-bits, so every number eventually reaches 1.
- **Generalization — k -happy:** Replace squares with k -th powers. Standard happy is 2-happy. $k = 3$ gives "cubed happy" numbers, etc. The same bounding argument shows every k has finitely many attractors.

Happy numbers are not deep in algebraic number theory, but they are a perfect classroom example of iteration, attractors, invariants, and proof by descent: to prove an infinite process has only two outcomes, you show it must eventually enter a finite, checkable region.

Think of S as machine:

$$S(n) = \sum_{\text{decimal digits } d \text{ of } n} d^2$$

Example $n = 145$: digits 1, 4, 5 $\rightarrow S(145) = 1^2 + 4^2 + 5^2 = 1 + 16 + 25 = 42$.

Iterating creates orbit:

$$\begin{aligned} n_0 &= n \\ n_1 &= S(n_0) \\ n_2 &= S(n_1) \\ &\vdots \end{aligned}$$

Two worked orbits with:

$$\begin{aligned} 23 &\rightarrow S(23) = 2^2 + 3^2 = 4 + 9 = 13 \\ &\rightarrow S(13) = 1^2 + 3^2 = 1 + 9 = 10 \\ &\rightarrow S(10) = 1^2 + 0^2 = 1 \\ &\rightarrow S(1) = 1^2 = 1 \text{ fixed point — happy} \end{aligned}$$

$$\begin{aligned} 36 &\rightarrow 3^2 + 6^2 = 9 + 36 = 45 \\ &\rightarrow 4^2 + 5^2 = 16 + 25 = 41 \\ &\rightarrow 4^2 + 1^2 = 16 + 1 = 17 \\ &\rightarrow 1^2 + 7^2 = 1 + 49 = 50 \\ &\rightarrow 5^2 + 0^2 = 25 \\ &\rightarrow 2^2 + 5^2 = 4 + 25 = 29 \\ &\rightarrow 2^2 + 9^2 = 4 + 81 = 85 \\ &\rightarrow 8^2 + 5^2 = 64 + 25 = 89 \\ &\rightarrow 8^2 + 9^2 = 64 + 81 = 145 \\ &\rightarrow 1^2 + 4^2 + 5^2 = 42 \\ &\rightarrow 4^2 + 2^2 = 20 \\ &\rightarrow 2^2 + 0^2 = 4 \\ &\rightarrow 4^2 = 16 \\ &\rightarrow 1^2 + 6^2 = 37 \\ &\rightarrow 3^2 + 7^2 = 58 \\ &\rightarrow 5^2 + 8^2 = 89 \text{ loop detected — unhappy} \end{aligned}$$

Once 4 appears, you loop $4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4$.

Why no number escapes to infinity — the descent lemma:

Let n have k digits, so $10^{k-1} \leq n < 10^k$ for $n \geq 1$.

$$S(n) \leq k \cdot 9^2 = 81k$$

Because each digit at most 9.

Compare n vs $81k$:

$$\begin{aligned} k = 1 : n &\leq 9, \quad 81k = 81 \\ k = 2 : n &\leq 99, \quad 81k = 162 \\ k = 3 : n &\leq 999, \quad 81k = 243 \\ k = 4 : n &\geq 1000, \quad 81k = 324 < n \\ k = 5 : n &\geq 10000, \quad 81k = 405 < n \end{aligned}$$

For $k \geq 4$, $10^{k-1} > 81k$, so $S(n) < n$.

Thus any $n \geq 1000$ strictly decreases. Iterate:

$$n_0 > n_1 > n_2 > \dots \text{ until } < 1000$$

Cannot decrease forever (positive integers), must enter

.Onceinside

, $S(n) \leq 3 \cdot 81 = 243$, then $S(n) \leq 2 \cdot 81 + 3^2 = \dots < 1000$, so stays below 1000.[1][999]

Therefore to understand all n , it suffices to compute orbits of 1 to 999 — finite check. That computation reveals exactly two attractors:

$1 \rightarrow 1$ — fixed point

$4 \rightarrow 16 \rightarrow 37 \rightarrow 58 \rightarrow 89 \rightarrow 145 \rightarrow 42 \rightarrow 20 \rightarrow 4$ — 8-cycle

No other loops. So every starting n ends in one of those two.

Properties in formulas:

Permutation invariance:

$$S(\text{permutation of digits of } n) = S(n)$$

Because sum of squares commutative. Example $S(19) = 1^2 + 9^2 = 82 = S(91)$.

If orbit of n hits 1, orbit of any permutation hits same next value $S(n)$, hence also hits 1.

Zero invariance:

$$S(1009) = 1^2 + 0^2 + 9^2 = 1^2 + 9^2 = S(19)$$

$0^2 = 0$ adds nothing. So appending zeros does not change fate.

Heredity:

$$n \text{ happy} \implies S(n) \text{ happy}$$

$$n \text{ unhappy} \implies S(n) \text{ unhappy}$$

Because $S(n)$ is next step in orbit — if orbit of n ends at 1, orbit of $S(n)$ is suffix of same orbit, also ends at 1.

Infinitude:

$$1 \text{ happy} \implies 10^k \text{ happy for all } k \geq 0$$

$$2 \rightarrow 4 \text{ unhappy} \implies 2 \cdot 10^k \text{ unhappy for all } k \geq 0$$

Thus infinitely many each.

Density question:

$$D(X) = \frac{\#\{n \leq X : n \text{ happy}\}}{X}$$

Computations:

$$D(10^3) \approx 0.20$$

$$D(10^6) \approx 0.15$$

$$D(10^9) \approx 0.15\text{-}0.19 \text{ range}$$

Conjectured limit ≈ 0.15 , but existence of $\lim_{X \rightarrow \infty} D(X)$ open.

Consecutive happy numbers:

31 happy ($31 \rightarrow 10 \rightarrow 1$), 32 happy ($32 \rightarrow 13 \rightarrow 10 \rightarrow 1$) — smallest adjacent pair.

Longer runs exist via Chinese remainder / digit construction — e.g., find m such that S values of $m, m+1, \dots$ map into known happy numbers. Existence of arbitrarily long runs conjectured but not proved.

Base dependence and generalization:

Define for base $b \geq 2$:

$$S_{b,k}(n) = \sum_{\text{base-}b \text{ digits } d} d^k$$

Standard happy $b = 10, k = 2$. For each b, k , same descent argument: for k fixed, $S_{b,k}(n) = O(\log_b n)$, so $S_{b,k}(n) < n$ for large n , finite check suffices, finitely many attractors. 1 always fixed because $1^k = 1$.

Example $b = 2, k = 2$: $S_2(n)$ = number of 1-bits in binary (since $0^2 = 0, 1^2 = 1$). Then $S_2(n) \leq \log_2 n + 1$, iteration goes to 1 for all $n \geq 1$ — every number happy in base 2.

For $b = 10, k = 3$ (cubed happy): $S(n) = \sum d^3$, only attractors are 1, $55 \rightarrow 250 \rightarrow 133 \rightarrow 55$ etc.

This makes happy numbers accessible illustration of dynamical systems: map S , phase space $\mathbb{N}$, two basins of attraction, trapping region 55 , invariants under digit permutation and zero insertion.[1][999]

Modern Developments and Variations

- **Density:** Let $D(X) = \frac{\#\{n \leq X : \text{happy}\}}{X}$. Empirically $D(X) \approx 0.15$ - 0.20 . Best known bounds $0.12 \lesssim D(X) \lesssim 0.19$ for large X (Gilmer, 2013; El-Sedy & Siksek, 2000). Existence of natural density $\lim_{X \rightarrow \infty} D(X)$ open — analogous to Collatz density problem.
- **Happy primes:** 7, 13, 19, 23, 31, 79, 97, 103, 109, 139, 167, ... — OEIS A035497. Infinite? Conjectured yes, not proved, but follows from plausible distribution of happy among primes.
- **Consecutive runs:** 31, 32 smallest pair. Construction using $n \rightarrow S(n)$ mapping and Chinese remainder allows building long runs: if a such that S values cover interval of happy numbers, $a \cdot 10^k + \text{offset}$ yields run. Longest recorded run as of 2024 is 12 consecutive numbers. Existence of arbitrarily long runs conjectured, not proved — similar to open problem for prime runs.
- **Base b , power k generalization:** $S_{b,k}(n) = \sum_{\text{base-}b \text{ digits } d} d^k$. For each b, k , same descent: $S_{b,k}(n) = O(\log n) < n$ for large n , so finitely many cycles, checkable. Base 2, $k = 2$: $S_2(n)$ = popcount(n) — all n happy. Base 10, $k = 3$: cycles include 1, $55 \rightarrow 250 \rightarrow 133 \rightarrow 55$, etc. — "cubes happy".
- **Pedagogical use:** Since 1970s, standard example in introductory programming (cycle detection via Floyd, hash set) and discrete dynamical systems — illustrates trapping region, attractor, invariant, and proof that infinite process reduces to finite check — same technique used for Collatz-type problems, Kaprekar routine 6174, and $3n + 1$.

In short: recreational origin late 19th century, name coined mid-20th, formal cycle classification via bounding lemma $81k < 10^{k-1}$, modern interest in density, happy primes, and long runs, with B_n ? — not related — but with same flavor as Stirling/Bell enumeration of partitions: finite description captures infinite behavior.

Harshad / Niven Numbers

A Harshad number — from Sanskrit *harṣa* meaning "great joy" and *da* meaning "to give," literally "joy-giver" — also called a Niven number after Canadian mathematician Ivan M. Niven, who studied them in a 1977 paper, is an integer divisible by the sum of its digits in base 10.

The term Harshad was coined by the Indian recreational mathematician D. R. Kaprekar — the same Kaprekar of Kaprekar numbers and Kaprekar's constant 6174. He loved digit-dependent properties.

Definition: Let $s_{10}(n)$ be the sum of the decimal digits of n . Then n is Harshad (or Niven) if $s_{10}(n) \mid n$, i.e. $n \bmod s_{10}(n) = 0$.

- **Single digits:** Every number from 1 through 9 is trivially Harshad, since $s(d) = d$ and $d \mid d$. By convention we include 10 as Harshad because $1 + 0 = 1$ divides 10.
- **Origin of name:** Harshad = joy-giver, Niven = after Ivan Niven. Both names are used interchangeably in literature; "Harshad" is more common in recreational contexts, "Niven" in formal papers.
- **Universal numbers:** Only 1, 2, 4, 6 are Harshad numbers in *every* number base. These are called all-Harshad or all-Niven numbers.

History and Discovery

Harshad: Kaprekar 1955, Niven: Niven 1977

- **D. R. Kaprekar (1905–1986):** Indian schoolteacher in Devlali, Maharashtra, self-taught recreational mathematician. In 1955 paper "Multidigital Numbers" and later notes in *Scripta Mathematica*, he defined "Harshad" (joy-giver). Kaprekar was fascinated by base-10 digit interactions: Harshad numbers, Kaprekar numbers $\text{left}(n^2)$ split and sum to n , e.g. $45^2 = 2025, 20+25=45$ (right), Demlo numbers, Kaprekar constant 6174 from routine 4-digit sort-desc minus sort-asc $\rightarrow$ 6174 loop. He chose Sanskrit name to celebrate Indian mathematical tradition. His work was largely ignored by formal journals but circulated in recreational circles.
- **Ivan M. Niven (1915–1999):** Canadian-American number theorist at University of Oregon, author of *Irrational Numbers* and co-author of *Introduction to the Theory of Numbers*. In 1977 talk at conference and 1980 paper with H. S. Zuckerman, he independently studied numbers divisible by sum of digits,

calling them perhaps not knowing Kaprekar's term. His framework introduced b -Niven numbers (base b), $s_b(n) \mid n$, and proved basic density results. Western literature adopted "Niven number" after him; OEIS uses both.

- **Grundman 1994 — All-Harshad classification:** H. G. Grundman proved only 1, 2, 4, 6 are Harshad in every base $b \geq 2$. Proof uses representation $n = 11_b$ for $b = n - 1$, etc., forcing evenness and smallness.
- **De Koninck & Doyon, Cooper & Kennedy:** Jean-Marie De Koninck and Nicolas Doyon (2003) proved distribution results: number of b -Niven up to X is $o(X)$, natural density 0, yet there are arbitrarily long gaps of non-Niven. Curtis Cooper and Robert E. Kennedy (1984) proved existence of 20 consecutive Niven numbers in base 10, disproving early conjecture that gap bounded by 20 — showing block structure subtle.
- **Modern extensions:** Multiple Harshad (MFA) numbers studied by Grundman, k -Niven where n divisible by $s_b(n)^k$, strict Harshad where $n/s(n)$ coprime to $s(n)$, and Zuckerman numbers where n divisible by product of digits. All stem from Kaprekar's original joy-giver idea.

How to Check

1. **Sum the decimal digits:** $s(n)$.
2. **Divide:** is $n \bmod s(n) = 0$?

Examples:

- 18: $s = 1 + 8 = 9$, $\frac{18}{9} = 2$ Harshad
- 21: $s = 2 + 1 = 3$, $\frac{21}{3} = 7$ Harshad
- 1729 — **the Hardy-Ramanujan taxicab number:** $s = 1 + 7 + 2 + 9 = 19$, $\frac{1729}{19} = 91$ Harshad
- 19: $s = 1 + 9 = 10$, $19 \bmod 10 = 9$ not Harshad
- 100: $s = 1$, $100 \bmod 1 = 0$ — any power of 10 is Harshad

Sequence (base 10): All 1 – 10 are Harshad, then

12, 18, 20, 21, 24, 27, 30, 36, 40, 42, 45, 48, 50, 54, 60, 63, 70, 72, 80, 81, 84, 100, 102, 108, 110, 111, 112, ...

111 is interesting: $1 + 1 + 1 = 3$, $\frac{111}{3} = 37$.

How Divisibility Works

Let decimal expansion $n = \overline{d_k d_{k-1} \dots d_{010}} = \sum d_i 10^i$, define

$$s(n) = \sum_{i=0}^k d_i$$

Harshad condition:

$$s(n) \mid n \iff \exists q \in \mathbb{N} : n = q \cdot s(n)$$

$$q = \frac{n}{s(n)} \text{ called Harshad quotient}$$

Example $n = 18$: $s = 9$, $q = 18/9 = 2$.

Example $n = 1729$:

$$\begin{aligned} s(1729) &= 1 + 7 + 2 + 9 = 19 \\ \frac{1729}{19} &= 91 \quad \text{since } 19 \cdot 91 = 19 \cdot 90 + 19 = 1710 + 19 = 1729 \end{aligned}$$

So 1729 Harshad.

Why powers of 10 always work:

$$\begin{aligned} n &= 10^k = 1 \underbrace{0 \dots 0}_k \\ s(n) &= 1 \\ \frac{10^k}{1} &= 10^k \in \mathbb{N} \end{aligned}$$

Similarly $a \cdot 10^k$ with $1 \leq a \leq 9$: $s = a$, $a \mid a10^k$ always, so infinite family.

$$a \cdot 10^k \text{ Harshad for all } k \geq 0, 1 \leq a \leq 9$$

Mod 9 connection:

$$10 \equiv 1 \pmod{9} \implies 10^i \equiv 1^i \equiv 1 \pmod{9}$$

Hence

$$n = \sum d_i 10^i \equiv \sum d_i = s(n) \pmod{9}$$

So $n - s(n)$ always divisible by 9. If n Harshad, $n = qs(n)$, then

$$qs(n) \equiv s(n) \pmod{9}$$

$$s(n)(q-1) \equiv 0 \pmod{9}$$

Thus $9 \mid s(n)(q-1)$. This restricts possible q when $3 \nmid s(n)$. Example $n = 21$: $s = 3$, $q = 7$, $s(q-1) = 3 \cdot 6 = 18$ divisible by 9.

This is generalization of divisibility test for 9: $9 \mid n \iff 9 \mid s(n)$.

Key Properties

1. **All single digits are Harshad** — by definition.

2. **Infinitude:** There are infinitely many Harshad numbers. Simple families: 10^k , $a \cdot 10^k$ for $1 \leq a \leq 9$.

$$\{10^k : k \geq 0\} \subset \text{Harshad}, |\text{Harshad}| = \infty$$

3. **Density:** Natural density 0. Let $H(X) = \#\{n \leq X : n \text{ Harshad}\}$. Then

$$\lim_{X \rightarrow \infty} \frac{H(X)}{X} = 0$$

Reason: $s(n) \leq 9 \log_{10} X$ for $n \leq X$, so average size of divisor $s(n)$ much smaller than n , divisibility by small random-ish number $s(n)$ has probability about $1/s(n)$ on average, sum diverges slowly, proportion tends to 0. Empirically:

$$\begin{aligned} \frac{H(1000)}{1000} &\approx 0.30 \\ \frac{H(100000)}{100000} &\approx 0.14 \\ \frac{H(10^9)}{10^9} &\approx 0.05 \end{aligned}$$

Yet gaps: De Koninck-Doyon proved arbitrarily long runs of consecutive non-Harshad numbers exist. Cooper-Kennedy proved 20 consecutive Harshad exist in base 10 (e.g. starting at $10^{\text{something}}$), and Grundman showed any base b has runs of $2b$ consecutive b -Niven.

4. **Multiple Harshad / MFA Numbers:** A Multiple Harshad Number is one where repeatedly dividing by digit sum stays Harshad.

Let $n_0 = n$, $n_{i+1} = n_i / s(n_i)$ if divisible.

If this can be done k times and each n_i is Harshad, n has MFA-degree k .

Example: 2016

$$s(2016) = 2 + 0 + 1 + 6 = 9$$

$$\frac{2016}{9} = 224$$

$$s(224) = 8, 224/8 = 28$$

$$s(28) = 10, 28 \bmod 10 \neq 0 \text{ stop}$$

So 2016 degree 1.

Stronger: 378

$$s(378) = 18, 378/18 = 21$$

$$s(21) = 3, 21/3 = 7$$

$$s(7) = 7, 7/7 = 1$$

So chain $378 \rightarrow 21 \rightarrow 7 \rightarrow 1$, each Harshad, degree 3.

5. **All-Harshad Numbers (AHN):** Numbers Harshad in every base $b \geq 2$. Grundman 1994 proved only 1, 2, 4, 6.

Sketch: For $b = n - 1 \geq 2$, $n = 11_b$ (since $b + 1 = n$), so $s_b(n) = 2$, need $2 \mid n \rightarrow n$ even. For $b = n - 2$, $n = 12_b = b + 2$, $s = 3$, need $3 \mid n$ unless n small, etc., forcing $n \in \{1, 2, 4, 6\}$.

6. **Harshad Primes:** Prime Harshad can only be 2, 3, 5, 7.

Proof:

$$n \geq 11 \implies 2 \leq s(n) < n$$

$$s(n) \mid n \text{ and } 1 < s(n) < n \implies n \text{ composite}$$

If $s(n) = 1$, $n = 10^k$, composite for $k \geq 1$. So no multi-digit Harshad prime.

Generalizations

- **b -Niven:** $s_b(n) \mid n$, s_b sum base- b digits. Theory changes with b .
- **Strict Harshad:** $\frac{n}{s(n)}$ Harshad and $\gcd\left(s(n), \frac{n}{s(n)}\right) = 1$.
- **Zuckerman / s -divisible:** Product of digits divides n .
- **Connection to 9:** $n \equiv s(n) \pmod{9}$ as above, underlying all base-10 digit sum divisibility.

Thus Harshad numbers link Kaprekar's recreational joy (1955) to Niven's formal study (1977) to modern questions about distribution of $s_b(n) \pmod{m}$ and long runs — simple definition, deep irregular behavior because $s(n)$ fluctuates while divisibility is rigid.

Wieferich Primes

They come from asking what happens when you strengthen Fermat's Little Theorem by one power of p .

Fermat's Little Theorem: For prime p not dividing a ,

$$a^{p-1} \equiv 1 \pmod{p}$$

So p always divides $a^{p-1} - 1$. Arthur Wieferich asked in 1909: When does p^2 divide $a^{p-1} - 1$? When does the congruence hold $\pmod{p^2}$ instead of just $\pmod{p}$?

Definition: A prime p is a Wieferich prime base a if

$$a^{p-1} \equiv 1 \pmod{p^2}$$

In other words, $p^2 \mid a^{p-1} - 1$. When no base is mentioned, base 2 is the default: $2^{p-1} \equiv 1 \pmod{p^2}$.

For base 2, this is p^2 divides the Mersenne number $M_{p-1} = 2^{p-1} - 1$.

Example: $p = 1093$ is Wieferich base 2:

$$2^{1092} = 1 + k \cdot 1093^2$$

with $k = 364245589\dots$ large. The key point is the remainder upon division by $1093^2 = 1,194,649$ is 1. So 1093 divides $2^{1092} - 1$ twice.

For contrast, take $p = 5$ non-Wieferich: $2^4 = 16$, $16 - 1 = 15$, divisible by 5 but not by 25. So 5 is not Wieferich base 2.

History and Discovery

Wieferich 1909 — Meissner 1913 — Beeger 1922 — Modern sieve to 10^{17}

- **Arthur Wieferich (1884–1954), German mathematician, 1909:** In *Journal für die reine und angewandte Mathematik* paper "Zum letzten Fermatschen Theorem", he proved:

If $x^p + y^p = z^p$ with $p \nmid xyz$ (FLT first case) has integer solution, then $2^{p-1} \equiv 1 \pmod{p^2}$.

This was first restriction linking FLT failure to special primes. At the time FLT was great open problem, so any necessary condition for counterexample was intensely pursued. Wieferich's theorem instantly made primes satisfying $2^{p-1} \equiv 1 \pmod{p^2}$ important.

- **Mirimanoff 1910:** Russian mathematician Dmitry Mirimanoff extended Wieferich: if first case fails for p , then also $3^{p-1} \equiv 1 \pmod{p^2}$. So FLT counterexample needs p Wieferich for both bases 2 and 3 simultaneously. This made search for Wieferich primes base 2 and 3 a route to prove FLT first case for many p .
- **Meissner 1913 — first Wieferich found:** Using desk calculations, German mathematician Walther Meissner found $p = 1093$ satisfies $2^{1092} \equiv 1 \pmod{1093^2}$. Computation:

$$1093^2 = 1194649$$

$$2^{1092} \bmod 1194649 = 1$$

He verified by repeated squaring mod p^2 — heroic for 1913.

- **Beeger 1922 — second:** Dutch mathematician N. G. W. H. Beeger found $p = 3511$ satisfies same. No other below 10^9 at that time.
- **1909–1995 FLT hunt era:** Before Wiles proof of FLT in 1995, Wieferich criterion used to prove FLT first case for all p up to large bounds: if no Wieferich primes exist up to X , then FLT first case true for $p \leq X$. Computations of $q_p(2) = \frac{2^{p-1} - 1}{p} \bmod p$ driven by FLT. Work by Vandiver, Lehmer, Emma Lehmer (1930s – 1960s) expanded search to 10^6 , then 10^9 with early computers.
- **After FLT proved:** Interest shifted from FLT to intrinsic number theory: distribution of Fermat quotients, p -adic logarithms, abc conjecture, cryptography.

Key Facts and Properties

1. Known primes — Extreme Rarity: Only two base-2 Wieferich primes are known:

$$1093 \quad \text{and} \quad 3511$$

- 1093 was found by Meissner in 1913.
- 3511 was found by Beeger in 1922.

Despite enormous effort, no third is known.

2. Rarity and Search Limits: Computer searches have tested all primes up to $> 1.45 \times 10^{17}$ as of 2023 — that is 145 quadrillion — without finding any additional base-2 Wieferich primes. PrimeGrid, Dorais-Klyve, and others use efficient sieving using $a^{p-1} \bmod p^2$ can be computed with $O(\log p)$ modular multiplications using fast exponentiation, but the sheer range makes it hard.

Heuristic: For random prime p , $a^{p-1} \bmod p^2$ is roughly uniformly distributed among the p multiples of p that are $1 \bmod p$. So probability it is $1 \bmod p^2$ is $\frac{1}{p}$. Expected number up to X is $\sum_{p \leq X} \frac{1}{p} \sim \log \log X$, which diverges but agonizingly slowly: $\log \log(10^{17}) \approx 3.66$, consistent with 2 found. This suggests infinitely many exist, but the next may be beyond 10^{18} .

It is conjectured there are infinitely many Wieferich primes base a for every a , but unproven for any a .

3. Connection to Fermat's Last Theorem: German mathematician Arthur Wieferich discovered in 1909 the result that made these primes famous:

If the first case of Fermat's Last Theorem fails for prime exponent p — i.e., there exist integers x, y, z not divisible by p with $x^p + y^p = z^p$ — then p must be a Wieferich prime base 2.

FLT's first case was eventually proved for all p up to huge bounds using this: if you could prove no Wieferich primes exist in a range, FLT first case holds there. This drove searches for Wieferich primes for 70 years, until Wiles proved FLT completely in 1995 by different methods. Mirimanoff in 1910 extended it: p must also be Wieferich base 3.

Even after FLT, the link remains: Wieferich primes are obstructions to the "first case" and to the abc conjecture — abc implies infinitely many *non*-Wieferich primes.

4. Other bases — a -Wieferich primes: Mathematicians study Wieferich primes to bases other than 2. Definition is same: $a^{p-1} \equiv 1 \bmod p^2$, $p \nmid a$.

- **Base 3:** 11, 1006003 — only two known below 10^{15} , plus huge ones like ...
- **Base 5:** 2, 20771, 40487, 53471161, ... — 2 is trivially Wieferich base 5 because $5^1 \equiv 1 \bmod 4$? Wait 2 base 5: $5^1 = 5 \equiv 1 \bmod 4$? $2^2 = 4$ divides $5^1 - 1 = 4$, yes.
- **Base 10:** 3, 487 — called repunit Wieferich because $10^{p-1} \equiv 1 \bmod p^2$ means p^2 divides repunit 999...9.
- Base a and prime $p = a$ is excluded because $p \mid a$.

For each base, the set is conjectured infinite but appears sparse. Base 2 is most studied because of FLT and Mersenne numbers.

5. The Math Behind It — Fermat Quotient: Define the Fermat quotient:

$$q_p(a) = \frac{a^{p-1} - 1}{p}$$

Fermat's theorem says $q_p(a)$ is integer. Wieferich condition is $q_p(a) \equiv 0 \bmod p$.

$q_p(a)$ behaves like a logarithm mod p — called the p -adic logarithm — satisfying $q_p(ab) \equiv q_p(a) + q_p(b) \pmod{p}$. Wieferich primes are where this logarithm vanishes.

Why They Matter Today

Fermat quotient and p -adic log:

$$q_p(a) = \frac{a^{p-1} - 1}{p} \in \mathbb{Z}$$

Properties:

$$\begin{aligned} q_p(ab) &\equiv q_p(a) + q_p(b) \pmod{p} \\ q_p(a^k) &\equiv kq_p(a) \pmod{p} \end{aligned}$$

So q_p mimics log. Wieferich base a means $q_p(a) \equiv 0 \pmod{p}$ — logarithm zero.

Heuristic rarity:

Write $a^{p-1} = 1 + kp$, $0 \leq k < p$ determines $k \pmod{p^2}$ via $k \pmod{p}$. $k \equiv 0 \pmod{p}$ is Wieferich. If k random uniform among p values, probability $\frac{1}{p}$.

Expected count up to X :

$$\begin{aligned} E(X) &= \sum_{p \leq X} \frac{1}{p} \sim \log \log X + \text{Mertens for primes} \\ X = 10^3 : \log \log X &\approx 1.93 \\ X = 10^{17} : \log \log X &\approx 3.66 \\ X = 10^{100} : \log \log X &\approx 5.43 \end{aligned}$$

So even up to googol, expect ~5 Wieferich. Explains 2 found and why third may be far.

Modern Developments and Search Technology:

- **Crandall, Dilcher, Pomerance 1997:** Efficient algorithm to compute $q_p(2) \pmod{p}$ using $O(p)$? Actually using FFT and splitting, enabling search to 10^{12} .
- **Dorais-Klyve 2011, PrimeGrid 2020-2023:** Distributed search to 6.7×10^{15} , then 1.45×10^{17} using GPUs. Technique: compute $2^{p-1} \pmod{p^2}$ with binary exponentiation, each multiplication $\pmod{p^2}$ fits 128-bit for $p < 10^{17}$ ($p^2 < 10^{34} < 2^{113}$), so need 128-bit modular multiply, implemented in hardware.
- **No third found:** Lower bound $> 1.45 \times 10^{17}$ means if p Wieferich base 2 exists beyond known two, $p \geq 1.45 \times 10^{17}$.
- **Near-Wieferich:** Primes where $q_p(2)$ small but non-zero, e.g., $|q_p(2)| < p^\epsilon$. These are also studied, many known.

Other Bases:

$$\begin{aligned} \text{Base 2 : } &1093, 3511 \\ \text{Base 3 : } &11, 1006003 \\ \text{Base 5 : } &2, 20771, 40487, 53471161, 1645333507, \dots \\ \text{Base 10 : } &3, 487, 56598313, \dots \text{ (repunit)} \\ \text{Base } a = p : &\text{excluded} \end{aligned}$$

2 is Wieferich base 5 because $5^1 = 5$, $5 - 1 = 4$ divisible by $2^2 = 4$.

Modern Day Uses

- **Cyclotomic fields and Mersenne:** p Wieferich base 2 iff $\mathbb{Q}(\zeta_p)$ has 2 with extra ramification: order of 2 $\pmod{p^2}$ equals order $\pmod{p}$. This affects class numbers of cyclotomic fields, Iwasawa invariants.
- **Primality testing / pseudoprimes:** Base- a Fermat test checks $a^{p-1} \equiv 1 \pmod{p}$. Stronger test $\pmod{p^2}$ would catch pseudoprimes. Wieferich primes are where stronger test also passes, making them "exceptional" pseudoprimes. Understanding them helps analyze failure probability of Fermat and Miller-Rabin variants that use p^2 .
- **abc conjecture and non-Wieferich infinitude:** J. H. Silverman 1988 proved abc implies for any a , infinitely many primes p with $a^{p-1} \not\equiv 1 \pmod{p^2}$ — i.e., infinitely many non-Wieferich. So proof of many non-Wieferich would support abc . Converse open.
- **Wieferich pairs and Wieferich cubes:** Pair (p, q) where $p^{q-1} \equiv 1 \pmod{q^2}$ and $q^{p-1} \equiv 1 \pmod{p^2}$. Only known small examples like $(2, 1093)$ not symmetric pair, but $(83, 4871)$ is double Wieferich? Research on mutual Wieferich.

- **Cryptography — not direct key, but testbed:** Fast exponentiation mod p^2 used in searches is same primitive as RSA exponentiation. Benchmarking $a^{p-1} \bmod p^2$ for p up to 10^{17} stress-tests big-integer libraries, GPU modular arithmetic.
- **Waring / Wolstenholme analogy:** Similar strengthening from $\bmod p$ to $\bmod p^2$ appears in Wolstenholme primes $\binom{2p-1}{p-1} \equiv 1 \bmod p^3$, related to Bernoulli numbers. Wieferich and Wolstenholme both measure p -adic zeta values vanishing.

Summary

$$\begin{aligned} \text{Fermat : } a^{p-1} &= 1 + p q_p(a) \\ \text{Wieferich : } q_p(a) \equiv 0 \pmod{p} &\iff a^{p-1} = 1 + p^2 k \\ \text{Base 2 known : } p &= 1093, 3511, \quad p^2 = 1194649, 12326721 \\ \text{Example non : } p = 5 : \quad 2^4 = 16 &= 1 + 3 \cdot 5, \quad q_5(2) = 3 \not\equiv 0 \pmod{5} \\ \text{Heuristic count : } \sum_{p \leq X} \frac{1}{p} &\sim \log \log X \rightarrow \infty \text{ slowly} \\ \text{FLT link : } x^p + y^p = z^p, \quad p \nmid xyz &\implies q_p(2) \equiv 0 \text{ and } q_p(3) \equiv 0 \pmod{p} \\ \text{Open : } \exists \text{ infinitely many } p : q_p(2) &\equiv 0 \pmod{p} ? \\ \text{Known lower bound : } p &> 1.45 \times 10^{17} \text{ for next base-2} \end{aligned}$$

Thus Wieferich primes began as technical lemma for FLT in 1909, became computational milestone with Meissner and Beeger finding 1093 and 3511, survived FLT's 1995 proof to become central example of $\bmod p^2$ phenomenon, p -adic logarithm zero, and rarity $\sim \frac{1}{p}$ — only two known, next beyond 10^{17} , conjecturally infinite but agonizingly sparse.

Wilson Primes

They come from one of the most elegant theorems in elementary number theory.

Wilson's Theorem (1770): For an integer $p > 1$,

$$p \text{ is prime} \iff (p-1)! \equiv -1 \pmod{p}$$

That is, p divides $(p-1)! + 1$ exactly when p is prime. For example, $4! + 1 = 25$, divisible by 5; $5! + 1 = 121$, not divisible by 6.

Wilson primes ask: when does this congruence hold one power higher?

Definition: A prime p is a Wilson prime if

$$(p-1)! \equiv -1 \pmod{p^2}$$

i.e., $p^2 \mid (p-1)! + 1$.

If Wilson's Theorem says p divides $(p-1)! + 1$ once, a Wilson prime is a prime where p divides it twice.

History and Discovery

Ibn al-Haytham ~ 1000 — Wilson 1770 — Lagrange 1771 — Beeger 1913-1950s — Modern search to 2×10^{13}

- **Pre-history:** Result $(p-1)! \equiv -1 \pmod{p}$ for prime p known to Arab mathematician Ibn al-Haytham (Alhazen, ~ 1000 AD) and to Leibniz 1683, but not proved in surviving manuscripts.
- **John Wilson (1741–1793):** English mathematician, student of Edward Waring. Waring announced in *Meditationes Algebraicae* (1770) theorem without proof: "If p prime, then $(p-1)! + 1$ divisible by p ", crediting student Wilson. Waring added "the theorem is hard because no notation exists for factorial" — factorial notation $!$ came later.
- **Lagrange 1771:** First published proof by Joseph-Louis Lagrange, using that polynomial $x^{p-1} - 1 - (x-1)(x-2)\cdots(x-p+1)$ has $p-1$ roots mod p and coefficient comparison yields $(p-1)! \equiv -1 \pmod{p}$. Converse: if n composite $\neq 4$, $(n-1)! \equiv 0 \pmod{n}$, so Wilson distinguishes primes.
- **Wilson primes defined:** Natural strengthening: when does $p^2 \mid (p-1)! + 1$? First computed by hand.
- $p = 5$: $4! = 24$, $4! + 1 = 25 = 5^2$ — trivially Wilson, known to 19th century.
- $p = 13$: $12! = 479001600$, $12! + 1 = 479001601 = 13^2 \cdot 2834329$. Found by 19th-century factorial tables, often credited to 1900s.
- $p = 563$: Discovered by Karl Goldberg 1952 using early computer? Actually Leo Moser? Goldberg announced 563 as Wilson prime in 1953 after computation on early SWAC computer. Verification needs $562! \bmod 563^2$, $562! \approx 10^{1301}$ huge, but product modulo 316969 feasible by iterated multiplication.

- **Search era:** After 1950s, no new Wilson primes found despite search to 500, 10^4 , 10^5 , 5×10^8 (Crandall et al. 1997), 2×10^{13} (Costa et al. 2024) using algorithms $O(p \log p)$ for factorial mod p^2 , p -adic gamma function, and Wilson quotient.

The Math Behind Wilson Primes

Normal Wilson:

For prime p , define Wilson quotient:

$$W(p) = \frac{(p-1)! + 1}{p} \in \mathbb{Z}$$

By Wilson's theorem integer.

Example $p = 5$:

$$\begin{aligned} 4! &= 24 \\ W(5) &= \frac{24 + 1}{5} = 5 \end{aligned}$$

Example $p = 7$:

$$\begin{aligned} 6! &= 720 \\ W(7) &= \frac{720 + 1}{7} = 103 \end{aligned}$$

$103 \mod 7 = 5 \neq 0$, so 7 not Wilson.

Squared rule:

$$p \text{ Wilson} \iff W(p) \equiv 0 \mod p \iff p^2 \mid (p-1)! + 1$$

Because $(p-1)! + 1 = pW(p)$, need $p \mid W(p)$.

$$(p-1)! + 1 = pW(p) = p \cdot (p \cdot m) = mp^2$$

This parallels Wieferich: Fermat quotient $q_p(a) = \frac{a^{p-1} - 1}{p}$, Wieferich condition $q_p(a) \equiv 0 \mod p$. Wilson quotient is factorial analogue.

Why Wilson's theorem true — quick pairing idea:

For prime p , each $a = 1, \dots, p-1$ has inverse $a^{-1} \mod p$ distinct unless $a \equiv \pm 1 \mod p$ (since $a^2 \equiv 1 \mod p \implies p \mid a^2 - 1$). Pair up 2 with inverse 3, 4 with 5, etc., product $\equiv 1$. Leftover 1 and $p-1 \equiv -1$, so total product $\equiv -1$.

$$(p-1)! \equiv 1 \cdot (p-1) \cdot \prod_{\text{pairs}} (a \cdot a^{-1}) \equiv -1 \mod p$$

Known Values and Verification

Only three known:

5, 13, 563

1. $p = 5$

$$\begin{aligned} 4! &= 24 \\ 4! + 1 &= 25 \\ \frac{25}{5^2} &= 1 \\ W(5) &= 5 \equiv 0 \mod 5 \end{aligned}$$

2. $p = 13$

$$\begin{aligned} 12! &= 479001600 \\ 12! + 1 &= 479001601 \\ \frac{12! + 1}{13} &= 36846277 = W(13) \\ \frac{W(13)}{13} &= \frac{36846277}{13} = 2834329 \\ 13^2 &= 169, 479001601/169 = 2834329 \\ \text{Thus } 12! &\equiv -1 \mod 169 \end{aligned}$$

Check $W(13) \mod 13 = 0$.

3. $p = 563$

$562!$ has about $\log_{10} 562! \approx 1306$ digits — impossible to write. Compute iteratively modulo $p^2 = 316969$:

$$\begin{aligned} r_1 &= 1 \\ r_{k+1} &= r_k \cdot (k+1) \pmod{316969}, \quad k = 1 \dots 562 \\ \text{Result } r_{562} &= 316968 \equiv -1 \pmod{316969} \end{aligned}$$

Each step numbers $< p^2$, feasible $O(p)$ multiplies. Modern search uses faster $O(p \log p)$ grouping via polynomial and p -adic gamma Γ_p to avoid p steps per prime.

Facts and Mysteries

Extreme rarity and search limits:

Search complexity per prime p is $O(p)$ multiplies mod p^2 . Sum over $p \leq X$ about $\frac{X^2}{\log X}$ operations naive, too large. Improved algorithms using Wilson primes via $(p-1)! \equiv \Gamma_p(p) \pmod{p^2}$ and splitting product into blocks using FFT reduce to near $O(X)$.

Current verified bound:

$$\text{No Wilson prime } p \text{ with } 5 < p < 2 \times 10^{13}, \quad p \neq 13, 563$$

That is 20 trillion.

Heuristic $\frac{1}{p}$:

For prime p , Wilson theorem says $(p-1)! \equiv -1 \pmod{p}$, so $(p-1)! \pmod{p^2}$ lies among p residues congruent $-1 \pmod{p}$: $-1, -1+p, -1+2p, \dots, -1+(p-1)p$.

If random uniform among those p possibilities, probability exactly $-1 \pmod{p^2}$ is $\frac{1}{p}$.

Expected number up to X :

$$\begin{aligned} E(X) &= \sum_{p \leq X} \frac{1}{p} \sim \log \log X \\ X = 13 : \sum_{p \leq 13} \frac{1}{p} &= \frac{1}{2} + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{11} + \frac{1}{13} \approx 1.17 \\ X = 563 : \log \log 563 &\approx 1.90 \text{ expects } 2 \text{ Wilson} \\ X = 2 \times 10^{13} : \log \log X &\approx 3.4 \text{ expects } 3\text{-}4 \\ X = 10^{100} : \log \log X &\approx 5.43 \text{ expects } 5 \end{aligned}$$

Matches observation 3 found, predicts next may be $\gg 10^{13}$, perhaps $\sim 10^{30}$.

Conjectured infinitely many, but no proof for any type of supercongruence prime (Wieferich, Wilson, Wolstenholme).

Connections:

- Wolstenholme's theorem:** For $p > 3$, $\binom{2p-1}{p-1} \equiv 1 \pmod{p^3}$. Wolstenholme primes where $\equiv 1 \pmod{p^4}$ are 16843, 2124679 — similarly rare.
- Bernoulli numbers:** Wilson quotient formula Lerch:

$$W(p) \equiv B_{p-1} - B_{2p-2} \pmod{p}$$

Actually $W(p) \equiv \frac{1}{p}((p-1)! + 1)$ linked to B_{p-1} numerator. Irregular primes ($p \mid$ numerator B_{2k}) include 37, 59, 67, $\dots$; 563 is irregular — $563 \mid B_{\text{something}}$, linking Wilson to class number of cyclotomic field $\mathbb{Q}(\zeta_p)$.

- Wilson primes are Lerch primes?** Lerch formula relates Wilson quotient to sum $\sum_{a=1}^{p-1} q_p(a)$.

Modern day uses and why they matter:

- Supercongruence prototype:** Wilson primes are canonical example where congruence $\pmod{p}$ lifts to $\pmod{p^2}$ unexpectedly. Study of supercongruences mod p^2, p^3 drives p -adic analysis, p -adic gamma function Γ_p , and p -adic L-functions. Wilson p -adic gamma: $\Gamma_p(p) = (p-1)!$ up to sign, so Wilson condition $\Gamma_p(p) \equiv -1 \pmod{p^2}$.

- **Algorithm development:** Fast factorial mod p^2 algorithms developed for Wilson search — splitting product $1 \cdot 2 \cdot \dots \cdot p - 1$ into blocks, using polynomial evaluation and Wilson's theorem for intervals — now used in primality testing, factorial prime searches ($n! \pm 1$), and computation of p -adic invariants.
- **Cryptography — not direct:** Not used as key, but failure mode for Wilson-based primality test: test n prime iff $(n - 1)! \equiv -1 \pmod n$ is true but factorial huge, not practical. Understanding Wilson primes shows where even stronger test $(n - 1)! \equiv -1 \pmod{n^2}$ would falsely suggest composite? Actually composite never satisfies Wilson mod n , but Wilson primes satisfy stronger condition.
- **p -adic and abc :** Like Wieferich, abc conjecture implies infinitely many non-Wilson primes. Existence of infinitely many Wilson primes also expected from $\log \log$ heuristic.

In short: Every prime p satisfies $(p - 1)! \equiv -1 \pmod p$. Wilson primes are the extremely rare primes where divisibility happens twice: $p^2 \mid (p - 1)! + 1$. Only 5, 13, 563 known, with 5 giving 25, 13 giving $169 \mid 12! + 1$, 563 verified via modular product, no fourth below 2×10^{13} , expected count $\sum \frac{1}{p} \sim \log \log X$ suggests infinitely many but next may be astronomically large, and their study drives p -adic gamma and supercongruence theory.

Carmichael Numbers

Carmichael numbers are composite impostors — they pretend to be prime when tested with Fermat's Little Theorem. They are the worst possible case for naive primality testing, and the reason we cannot use the converse of Fermat's Little Theorem as a primality test.

History and Discovery

Fermat 1640 — Korselt 1899 — Carmichael 1910 — Chernick 1939 — Alford-Granville-Pomerance 1994

- **Pierre de Fermat (1640):** Fermat's Little Theorem: if p prime, $\gcd(a, p) = 1$, then $a^{p-1} \equiv 1 \pmod p$. Stated without proof in letter to Frénicle.
- **Sarrus (1820):** First noted converse false: $341 = 11 \cdot 31$ composite but $2^{340} \equiv 1 \pmod{341}$. So some composites pass base-2 test — first Fermat pseudoprime.
- **Arthur Korselt (1899):** German mathematician in *Archiv der Mathematik und Physik* gave complete characterization of absolute pseudoprimes: n composite square-free, $p - 1 \mid n - 1$ for all $p \mid n$. He proved theorem but had no example, did not know if any n satisfying it existed — pure criterion.
- **Robert Daniel Carmichael (1879–1967):** American mathematician at University of Illinois. In 1910 paper, he found first example $561 = 3 \cdot 11 \cdot 17$ satisfying Korselt. In 1912 he found 1105, 1729, 2465, 2821, 6601, 8911, 10585, 15841, ... Hence name Carmichael numbers, though Korselt had priority for criterion. Carmichael also defined Carmichael function $\lambda(n)$.
- **Jack Chernick (1939):** Observed parametric form: if $6k + 1, 12k + 1, 18k + 1$ prime, then $n = (6k + 1)(12k + 1)(18k + 1)$ Carmichael. For $k = 1$: $7 \cdot 13 \cdot 19 = 1729$ — Hardy-Ramanujan taxicab $1729 = 1^3 + 12^3 = 9^3 + 10^3$ is also Carmichael. This gave infinite candidate family conditional on prime triplets.
- **Paul Erdős heuristic (1956):** Conjectured infinitely many Carmichael numbers, outlined strategy: choose many primes p where $p - 1$ smooth (all prime factors small), product of subset will satisfy $p - 1 \mid n - 1$ via combinatorial argument.
- **Alford, Granville, Pomerance (1994):** Landmark Annals of Mathematics paper "There are infinitely many Carmichael numbers". Proved $C(X) \geq X^{2/7}$ for large X , where $C(X)$ = count of Carmichael $\leq X$. Made Erdős heuristic rigorous using zero-free regions of L-functions and combinatorial number theory. Later improved by Harman to $X^{0.33}$, and best heuristic $X^{1-o(1)}$ expected? Actually Pomerance conjectured $C(X) = X^{1-o(1)}$? Current conjecture $C(X) = X^{1/2+o(1)}$? Data suggests $C(X)$ roughly exp.
- **Modern computations:** Pinch (2007) enumerated up to 10^{18} : 1, 401, 644 Carmichael. To 10^{21} , about 20 million. Used to test primality libraries.

Fermat's Little Theorem and Its Converse

Fermat's Little Theorem:

$$\text{If } p \text{ prime, } \gcd(a, p) = 1 \implies a^{p-1} \equiv 1 \pmod p$$

Many hope converse true: If $a^{n-1} \equiv 1 \pmod n$, then n prime. False.

Composite n with $a^{n-1} \equiv 1 \pmod n$ for given a called **Fermat pseudoprime base a** .

Classic $341 = 11 \times 31$:

$$2^{340} \equiv 1 \pmod{341}$$

Yet 341 composite. So 341 looks prime testing base 2. Known to Sarrus 1820, smallest base-2 pseudoprime.

Carmichael numbers are ultimate pseudoprimes — pseudoprime to *every* base coprime.

Definition: Composite n is **Carmichael** (absolute pseudoprime) if

$$a^{n-1} \equiv 1 \pmod n$$

for every a with $\gcd(a, n) = 1$. Equivalently $a^n \equiv a \pmod n$ for all integers a .

So for true prime p , $p - 1$ bases pass. For Carmichael n , $\varphi(n)$ bases pass — all.

First examples:

$$\begin{aligned} 561 &= 3 \cdot 11 \cdot 17 \\ 1105 &= 5 \cdot 13 \cdot 17 \\ 1729 &= 7 \cdot 13 \cdot 19 \\ 2465 &= 5 \cdot 17 \cdot 29 \\ 2821 &= 7 \cdot 13 \cdot 31 \\ 6601 &= 7 \cdot 23 \cdot 41 \\ 8911 &= 7 \cdot 19 \cdot 67 \end{aligned}$$

Check 561:

$$\begin{aligned} 2^{560} &\equiv 1 \pmod{561} \\ 5^{560} &\equiv 1 \pmod{561} \\ 7^{560} &\equiv 1 \pmod{561} \end{aligned}$$

for any of $\varphi(561) = 320$ numbers a with $\gcd(a, 561) = 1$.

Korselt's Criterion (1899)

Theorem: n Carmichael iff:

1. **Composite and square-free:** $n = p_1 p_2 \cdots p_k$ distinct primes. If $p^2 \mid n$, choose $a = 1 + p$, then $(1 + p)^p \equiv 1 + p^2 \not\equiv 1 \pmod{p^2}$, fails.
2. **At least three primes:** $k \geq 3$. Proof sketch: if $n = pq$, need $p - 1 \mid pq - 1$ and $q - 1 \mid pq - 1$. Write $pq - 1 = q(p - 1) + (q - 1)$, so $p - 1 \mid q - 1$ and $q - 1 \mid p - 1$ impossible unless $p = q$.
3. **Divisibility:** $p - 1 \mid n - 1$ for every $p \mid n$.

Why (3) key: If $a^{n-1} \equiv 1 \pmod n$, then $\pmod p$ also 1. Take a primitive root mod p — order $p - 1$. Then $p - 1 \mid n - 1$.

Examples with:

$$\begin{aligned} 561 &= 3 \cdot 11 \cdot 17 \\ 3 - 1 &= 2 \mid 560 \\ 11 - 1 &= 10 \mid 560 \\ 17 - 1 &= 16 \mid 560 = 35 \cdot 16 \end{aligned}$$

$$341 = 11 \cdot 31 \text{ not Carmichael: } 31 - 1 = 30 \nmid 340$$

Indeed $3^{340} \equiv 56 \not\equiv 1 \pmod{341}$, so base 3 catches 341.

$$\begin{aligned} 1105 &= 5 \cdot 13 \cdot 17 \\ 4 \mid 1104, 12 \mid 1104 &= 92 \cdot 12, 16 \mid 1104 = 69 \cdot 16 \\ 1729 &= 7 \cdot 13 \cdot 19 \\ 6 \mid 1728, 12 \mid 1728 &= 144 \cdot 12, 18 \mid 1728 = 96 \cdot 18 \end{aligned}$$

Constructing via Chernick:

If $6k + 1, 12k + 1, 18k + 1$ prime, then product Carmichael because:

$$\begin{aligned} n &= (6k + 1)(12k + 1)(18k + 1) \\ n - 1 &= (6k + 1)(12k + 1)(18k + 1) - 1 \\ &\equiv 0 \pmod{6k}, 0 \pmod{12k}, 0 \pmod{18k} \end{aligned}$$

Since $L = \text{lcm}(6k, 12k, 18k) = 36k$, and one checks divisibility holds. For $k = 1$: 7, 13, 19 prime $\rightarrow 1729$.

Key Facts

- **Carmichael function** $\lambda(n)$: Smallest m with $a^m \equiv 1 \pmod n$ for all $\gcd(a, n) = 1$. For $n = \prod p_i$ odd square-free,

$$\lambda(n) = \text{lcm}(p_1 - 1, \dots, p_k - 1)$$

Then Korselt says

$$n \text{ Carmichael} \iff \lambda(n) \mid n - 1$$

Example $n = 561$:

$$\begin{aligned}\lambda(561) &= \text{lcm}(2, 10, 16) = 80 \\ 80 &\mid 560\end{aligned}$$

So exponent of group $(\mathbb{Z}/n\mathbb{Z})^\times$ divides $n - 1$.

- **Infinitude:** Alford-Granville-Pomerance 1994 proved infinitely many, at least $X^{2/7}$ up to X . Current best $X^{0.333\dots}$ by Harman. Conjectured $C(X) = X^{1-o(1)}$? Data:

$$\begin{aligned}C(10^3) &= 1 \\ C(10^6) &= 43 \\ C(10^9) &= 646 \\ C(10^{12}) &= 8241 \\ C(10^{15}) &= 105212 \\ C(10^{18}) &= 1401644 \\ C(10^{21}) &\approx 20138200\end{aligned}$$

Compare primes $\pi(10^{21}) \approx 2 \times 10^{19}$ — ratio $\sim 10^{-12}$.

- **Cryptographic importance:**

Fermat test picks random a , checks $a^{n-1} \equiv 1 \pmod n$. For prime, always passes. For typical composite, fails with probability $\geq 1/2$. For Carmichael, fails with probability 0 — fools every a coprime.

$$\Pr_a[\text{Fermat passes} \mid n \text{ Carmichael}] = 1$$

Hence Fermat test not suitable for crypto.

Miller-Rabin fixes: write $n - 1 = d \cdot 2^s$, d odd, check $a^d \equiv 1$ or $a^{d2^r} \equiv -1$ for some $0 \leq r < s$. Carmichael numbers fail this stronger condition.

Example $561 = 2^4 \cdot 35 + 1$, base 2:

$$\begin{aligned}2^{35} &\equiv 263 \pmod{561} \\ 2^{70} &\equiv 166 \pmod{561} \\ 2^{140} &\equiv 67 \pmod{561} \\ 2^{280} &\equiv 1 \pmod{561}\end{aligned}$$

Never ± 1 before 1, so Miller-Rabin declares composite — catches 561.

- **Distinction:**

$$\begin{aligned}\text{Fermat pseudoprime base } a : n \text{ composite, } a^{n-1} &\equiv 1 \pmod n \text{ for one } a \\ \text{Carmichael} : n \text{ composite, } a^{n-1} &\equiv 1 \pmod n \text{ for all } a, \gcd(a, n) = 1\end{aligned}$$

341 pseudoprime base 2 only, not Carmichael. 561 pseudoprime to all 320 bases.

Modern developments:

- **Distribution:** Granville-Pomerance conjecture $C(X) \sim X^{1 - \frac{(1+o(1)) \log \log \log X}{\log \log X}}$ — grows faster than any power X^ϵ ? Still open.
- **Erdős construction:** Modern proofs use primes p where $p - 1$ is y -smooth (all prime factors $\leq y$) and large $L = \text{lcm}$ of many $p - 1$, then product of subset $\equiv 1 \pmod L$ via pigeonhole.
- **Use in primality libraries:** Lists of Carmichael up to 10^{18} used as regression tests for Miller-Rabin, Baillie-PSW. OpenSSL, GMP test against them.
- **Absolute Euler pseudoprimes:** Subclass where $a^{(n-1)/2} \equiv \pm 1 \pmod n$ for all a with $\gcd(a, n) = 1$ and Legendre symbol condition — even rarer, e.g. 1729, 2465.

In short: Korselt 1899 characterized them before any example known, Carmichael 1910 found $561 = 3 \cdot 11 \cdot 17$ first, Chernick gave prime-triple construction 1729, Alford-Granville-Pomerance 1994 proved infinite, modern search enumerates to 10^{21} , and their existence forces cryptography to use Miller-Rabin, not naive Fermat — they are composites where $\lambda(n) = \text{lcm}(p_i - 1) \mid n - 1$, fooling $a^{n-1} \equiv 1 \pmod n$ for every a coprime.