

EECE 5610 Digital Control Systems

Lecture 7

Milad Siami

Assistant Professor of ECE

Email: m.siami@northeastern.edu



Northeastern University
College of Engineering

- Ideal sampler :

- Ideal sampler:

Consider $E^*(s) = \sum_{k=0}^{\infty} e(kT) e^{-kTs}$

$\Downarrow \mathcal{L}^{-1}$

$$\begin{aligned} e^*(t) &= \sum_{k=0}^{\infty} e(kT) \mathcal{L}^{-1} [e^{-kTs}] = \sum_{k=0}^{\infty} e(kT) \delta(t - kT) = \sum_{k=0}^{\infty} e(t) \delta(t - kT) \\ &= e(t) \sum_{k=0}^{\infty} \delta(t - kT) \end{aligned}$$

- Ideal sampler:

Consider $E^*(s) = \sum_{k=0}^{\infty} e(kT) e^{-kTs}$

$\Downarrow \mathcal{L}^{-1}$



$$\begin{aligned} e^*(t) &= \sum_{k=0}^{\infty} e(kT) \mathcal{L}^{-1} [e^{-kTs}] = \sum_{k=0}^{\infty} e(kT) \delta(t - kT) = \sum_{k=0}^{\infty} e(t) \delta(t - kT) \\ &= e(t) \sum_{k=0}^{\infty} \delta(t - kT) \end{aligned}$$

Let: $\delta_T = \sum_{k=0}^{\infty} \delta(t - kT)$ (a train of impulses)

Let: $\delta_T = \sum_0^{\infty} \delta(t - kT)$ (a train of impulses)



Let: $\delta_T = \sum_{k=-\infty}^{\infty} \delta(t - kT)$ (a train of impulses)

\Rightarrow The action of the ideal sampler can be thought of as modulating a train of impulses with the input signal $e(t)$



Let: $\delta_T = \sum_{k=0}^{\infty} \delta(t - kT)$ (a train of impulses)

\Rightarrow The action of the ideal sampler can be thought of as modulating a train of impulses with the input signal $e(t)$



\Rightarrow In the future we will think of $E^*(s)$ as the output of an ideal sampler.

- As mentioned before neither the ideal sampler nor the $\left[\frac{1-e^{-Ts}}{s} \right]$ block by themselves model the operation of a physical device. However, their combination does give the correct mathematical description of the sample & hold operation.


- As mentioned before neither the ideal sampler nor the $\left[\frac{1-e^{-Ts}}{s} \right]$ block by themselves model the operation of a physical device. However, their combination does give the correct mathematical description of the sample & hold operation.

- Q: Why do we go through all this trouble?

- As mentioned before neither the ideal sampler nor the $\left[\frac{1-e^{-Ts}}{s}\right]$ block by themselves model the operation of a physical device. However, their combination does give the correct mathematical description of the sample & hold operation.

• Q: Why do we go through all this trouble?

A: We want to get a T.F model of the hold operation and an input-output model of the sample and hold suitable for using in combination with linear systems analysis tools, such as the Z-transform we'll see later



$$\text{where } G_3(z) = Z \left[\left(\frac{1-e^{-Ts}}{s} \right) \cdot \text{Plant} \right]$$

Example: $e(t) = v(t)$

e^* ?

Example: $e(t) = u(t)$

e^* ?

$$e(nT) = u(nT) = 1 \quad \Rightarrow \quad e^*(t) = \sum_{n=0}^{\infty} \overset{1}{e(nT)} \delta(t - nT) = \sum_{n=0}^{\infty} \delta(t - nT)$$

Example: $e(t) = u(t)$

e^* ?

$$e(nT) = u(nT) = 1 \quad \Rightarrow \quad e^*(t) = \sum_{n=0}^{\infty} \overset{1}{e(nT)} \delta(t - nT) = \sum_{n=0}^{\infty} \delta(t - nT)$$

$$E^*(s) = \sum_{n=0}^{\infty} e^{-nTs} \overset{\text{LT}}{=} \frac{1}{1 - e^{-Ts}} \quad \left(\text{ROC: } |e^{-Ts}| < 1 \right)$$

Example: $e(t) = u(t)$

e^* ?

$$e(nT) = u(nT) = 1 \quad \Rightarrow \quad e^*(t) = \sum_{n=0}^{\infty} \overset{1}{e(nT)} \delta(t - nT) = \sum_{n=0}^{\infty} \delta(t - nT)$$

$$E^*(s) = \sum_{n=0}^{\infty} e^{-nTs} \xleftrightarrow{LT} = \frac{1}{1 - e^{-Ts}} \quad \left(\text{ROC: } |e^{-Ts}| < 1 \right)$$

(Note that the expression for $E^*(s)$ resembles that of $E(z)$, in fact, they are identical if we define $z = e^{sT}$. This is no accident, more on this later)

- How do we compute $E^*(s)$? (We'd like to have a closed form like in the example, rather than the infinite series)

- How do we compute $E^*(s)$? (We'd like to have a closed form like in the example, rather than the infinite series)

Facts:

- How do we compute $E^*(s)$? (We'd like to have a closed form like in the example, rather than the infinite series)

Facts:

$$(a) E^*(s) = \sum \left[\text{residues of } \left\{ E(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\} \text{ at poles of } E(\lambda) \right]$$

$$(b) E^*(s) = \frac{1}{T} \left[\sum_{-\infty}^{+\infty} E(s + jn\omega_s) + \frac{e(0)^+}{2} \right] \quad \text{where } \omega_s = 2\frac{\pi}{T}$$

= sampling frequency

- How do we compute $E^*(s)$? (We'd like to have a closed form like in the example, rather than the infinite series)

Facts:

$$(a) E^*(s) = \sum \left[\text{residues of } \left\{ E(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\} \text{ at poles of } E(\lambda) \right]$$

$$(b) E^*(s) = \frac{1}{T} \left[\sum_{-\infty}^{+\infty} E(s + jn\omega_s) + \frac{e(0)^+}{2} \right] \quad \text{where } \omega_s = 2\frac{\pi}{T}$$

= sampling frequency

Which one is useful to find $E^*(s)$?

- How do we compute $E^*(s)$? (We'd like to have a closed form like in the example, rather than the infinite series)

Facts:

$$(a) E^*(s) = \sum \left[\text{residues of } \left\{ E(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\} \text{ at poles of } E(\lambda) \right]$$

$$(b) E^*(s) = \frac{1}{T} \left[\sum_{-\infty}^{+\infty} E(s + jn\omega_s) + e^{\frac{(0)^+}{2}} \right] \quad \text{where } \omega_s = 2\frac{\pi}{T}$$

= sampling frequency

Which one is useful to find $E^*(s)$?

Which one is useful to determine properties of $E^*(s)$?

The proof uses again Cauchy's theorem. Sketch of the proof:

$$e^*(t) = e(t) \delta_T(t) \quad \text{where} \quad \delta_T(t) = \sum_{n=0}^{\infty} \delta(t - nT)$$

$$E^*(s) = \frac{1}{2\pi j} \int_{C-j\infty}^{C+j\infty} E(\lambda) \Delta_T(s-\lambda) d\lambda$$

$$\Delta_T(s) = \mathcal{L}[\delta_T(t)]$$

The proof uses again Cauchy's theorem. Sketch of the proof:

$$e^*(t) = e(t) \delta_T(t) \quad \text{where} \quad \delta_T(t) = \sum_{n=0}^{\infty} \delta(t-nT)$$

$$E^*(s) = \frac{1}{2\pi j} \int_{c-j\infty}^{c+j\infty} E(\lambda) \Delta_T(s-\lambda) d\lambda$$

$$\Delta_T(s) = \mathcal{L}[\delta_T(t)]$$

(Here we used the fact that product in time domain $\xleftrightarrow{\mathcal{L}}$ convolution in s ("frequency") domain)
where c must be chosen such that all poles of $E(\lambda)$ are to the left and all poles of Δ_T are to the right

The proof uses again Cauchy's theorem. Sketch of the proof:

$$e^*(t) = e(t) \delta_T(t) \quad \text{where} \quad \delta_T(t) = \sum_{n=0}^{\infty} \delta(t - nT)$$


$$E^*(s) = \frac{1}{2\pi j} \int_{c-j\infty}^{c+j\infty} E(\lambda) \Delta_T(s-\lambda) d\lambda$$

$$\Delta_T(s) = \mathcal{L}[\delta_T(t)]$$

(Here we used the fact that product in time domain $\xleftrightarrow{\mathcal{L}}$ convolution in s ("frequency") domain)
where c must be chosen such that all poles of $E(\lambda)$ are to the left and all poles of Δ_T are to the right

$$\text{Note that} \quad \Delta_T(s) = \mathcal{L}[\delta_T(t)] = \sum_{n=0}^{\infty} e^{-nTs} = \frac{1}{1 - e^{-Ts}} \Rightarrow \text{poles at } e^{-Ts} = 1 \\ \Rightarrow s = \pm j \frac{2\pi n}{Ts} = \pm j \omega_s n$$

so that $\Delta_T(s)$ has an infinite number of poles, all on the $j\omega$ axis, spaced ω_s



$\Delta_T(s)$ poles at $s = \pm j\omega_s$

$\Delta_T(s-\lambda)$ poles at $\lambda = s \pm j\omega_s$

Closing the contour with C_1 (a semicircle in the LHP with radius $r \rightarrow \infty$) yields the first equality. If, on the other hand, we close the contour with C_2 (semicircle in RHP, $r \rightarrow \infty$) we obtain the second formula

Example 1:

$$e(t) = v(t)$$

Example 1: $e(t) = u(t)$

$$\rightarrow E(\lambda) = \frac{1}{\lambda}; \quad E(\lambda) \frac{1}{1 - e^{-T(\lambda - \lambda)}} = \frac{1}{\lambda (1 - e^{-T(\lambda - \lambda)})}$$

Example 1: $e(t) = u(t)$

$$\rightarrow E(\lambda) = \frac{1}{\lambda}; \quad E(\lambda) \frac{1}{1 - e^{-T(\lambda)}} = \frac{1}{\lambda(1 - e^{-T(\lambda)})}$$

$$\sum_{\substack{\text{poles} \\ \text{of } E(\lambda)}} \text{Res} =$$

$$\sum_{\lambda=0} \text{Res} =$$

$$\frac{1}{1 - e^{-Ts}} \neq$$

Example 1: $e(t) = u(t)$

$$\rightarrow E(\lambda) = \frac{1}{\lambda}; \quad E(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} = \frac{1}{\lambda(1 - e^{-T(s-\lambda)})}$$

$$\sum_{\text{poles of } E(\lambda)} \text{Res} = \sum_{\lambda=0} \text{Res} = \frac{1}{1 - e^{-Ts}} \quad \#$$

Hint:

$$(a) E^*(s) = \sum \left[\text{residues of } \left\{ E(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\} \text{ at poles of } E(\lambda) \right]$$

$$(b) E^*(s) = \frac{1}{T} \left[\sum_{-\infty}^{+\infty} E(s + jn\omega_s) + \frac{e(0)^+}{2} \right] \quad \text{where } \omega_s = \frac{2\pi}{T}$$

= sampling frequency

$$\text{Res}(z_i) = \frac{1}{(n-1)!} \frac{d^{n-1}}{dz^{n-1}} \left[(z - z_i)^n F(z) \right] \Big|_{z=z_i} \quad \text{if } f(z) \text{ has a singularity of order } n \text{ at } z_i$$

$$\text{Res}(z_i) = (z - z_i) F(z) \Big|_{z=z_i} \quad \text{for singularities of order 1}$$

Example 2 :

Suppose that $f(t) = 1 - e^{-t}$

Example 2 : Suppose that $f(t) = 1 - e^{-t}$

$$\rightarrow F^*(s) = \sum_{k=0}^{\infty} (1 - e^{-kT}) e^{-kTs}$$

Example 2: Suppose that $f(t) = 1 - e^{-t}$

$$\rightarrow F^*(s) = \sum_{k=0}^{\infty} (1 - e^{-kT}) e^{-kTs}$$

$$= \sum_{k=0}^{\infty} e^{-kTs} - \sum_{k=0}^{\infty} e^{-k(s+1)T}$$

$$\boxed{\frac{1}{1 - e^{-Ts}} - \frac{1}{1 - e^{-(s+1)T}}}$$

Example 2: Suppose that $f(t) = 1 - e^{-t}$

$$\rightarrow F^*(s) = \sum_{k=0}^{\infty} (1 - e^{-kT}) e^{-kTs}$$

$$= \sum_{k=0}^{\infty} e^{-kTs} - \sum_{k=0}^{\infty} e^{-k(s+1)T} = \boxed{\frac{1}{1 - e^{-Ts}} - \frac{1}{1 - e^{-(s+1)T}}}$$

Alternatively: $F(s) = \frac{1}{s} - \frac{1}{(s+1)} = \frac{1}{s(s+1)}$, poles at $s=0, s=-1$

$$\sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \left\{ F(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\} = \sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \frac{1}{\lambda(\lambda+1)} \cdot \frac{1}{(1 - e^{-T(s-\lambda)})} = \frac{1}{1 - e^{-Ts}} - \frac{1}{1 - e^{-T(s+1)}} \neq$$

(same as before)

Example 2: Suppose that $f(t) = 1 - e^{-t}$

$$\rightarrow F^*(s) = \sum_{k=0}^{\infty} (1 - e^{-kT}) e^{-kTs}$$

$$= \sum_{k=0}^{\infty} e^{-kTs} - \sum_{k=0}^{\infty} e^{-k(s+1)T} = \boxed{\frac{1}{1 - e^{-Ts}} - \frac{1}{1 - e^{-(s+1)T}}}$$

Alternatively: $F(s) = \frac{1}{s} - \frac{1}{(s+1)} = \frac{1}{s(s+1)}$, poles at $s=0, s=-1$

$$\sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \left\{ F(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\} = \sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \frac{1}{\lambda(\lambda+1)} \cdot \frac{1}{(1 - e^{-T(s-\lambda)})} = \frac{1}{1 - e^{-Ts}} - \frac{1}{1 - e^{-T(s+1)}} \neq$$

(same as before)?

$T=1/2$

Example 3:

A function with a time-delay. This example will become relevant later on when we will look into the effects of sampling:


$$T=1/2$$

Example 3:

A function with a time-delay. This example will become relevant later on when we will look into the effects of sampling:

$$\text{Let } e(t) = [1 - e^{-(t-1)}] u(t-1)$$

(i.e: $e(t) = (1 - e^{-t})$ delayed by 1 second)



$$e(k) = [1 - e^{-(0.5k-1)}] \quad k \geq 2; \quad e(k) = 0 \quad k = 0, 1$$


$$T = 1/2$$

Example 3:

A function with a time-delay. This example will become relevant later on when we will look into the effects of sampling:

$$\text{Let } e(t) = [1 - e^{-(t-1)}] u(t-1)$$

(i.e: $e(t) = (1 - e^{-t})$ delayed by 1 second)



Starting Summation from $k = 2$ because first 2 terms are zero due to step delay

$$e(k) = [1 - e^{-(0.5k-1)}] \quad k \geq 2;$$

$$e(k) = 0 \quad k = 0, 1$$

$$T = 1/2$$

From the definition we have:

$$E^*(s) = \sum e(k) e^{-kTs} = \sum_{k=2}^{\infty} (1 - e^{-(0.5k-1)}) e^{-kTs} = \frac{e^{-2Ts}}{1 - e^{-Ts}} - \frac{e^{-2(0.5+Ts)}}{1 - e^{-(0.5+Ts)}}$$


$$= \frac{e^{-s}}{1 - e^{-0.5s}} - \cancel{\frac{e^{-1}}{1 - e^{-0.5s}}} \cdot \frac{e^{-s}}{1 - e^{-0.5(s+1)}} = \boxed{\frac{(1 - e^{-0.5}) e^{-1.5s}}{(1 - e^{-0.5s})(1 - e^{-0.5(s+1)})}} \quad \#$$

Example 3:

A function with a time-delay. This example will become relevant later on when we will look into the effects of sampling:

$$\text{Let } e(t) = [1 - e^{-(t-1)}] u(t-1)$$

(i.e: $e(t) = (1 - e^{-t})$ delayed by 1 second)




$$e(k) = [1 - e^{-(0.5k-1)}] \quad k \geq 2; \quad e(k) = 0 \quad k = 0, 1$$

Example 3:

A function with a time-delay. This example will become relevant later on when we will look into the effects of sampling:

$$\text{Let } e(t) = [1 - e^{-(t-1)}] u(t-1) \quad (\text{i.e.: } e(t) = (1 - e^{-t}) \text{ delayed by 1 second})$$



$$e(k) = [1 - e^{-(0.5k-1)}] \quad k \geq 2; \quad e(k) = 0 \quad k = 0, 1$$

Now let's try our residues formula:

$$E(s) = \frac{e^{-s}}{s(s+1)} \Rightarrow E^*(s) = \sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \left\{ \frac{e^{-\lambda}}{\lambda(\lambda+1)} \frac{1}{1 - e^{-T(s-\lambda)}} \right\} =$$

$$= \frac{1}{1 - e^{-Ts}} + \frac{e^1}{(-1)} \frac{1}{1 - e^{-T(s+1)}} = \boxed{\frac{1}{1 - e^{-0.5s}} - \frac{e^1}{1 - e^{-0.5(s+1)}}}$$

Surprise! we got different answers

Q: What went wrong here?

Surprise! we got different answers

Q: What went wrong here?  Cauchy's Integral does not work well with time-delays!!

A: A technical point: the "proof" of the residues formula is not valid for systems having time delays

The reason is that $e^{-sT} \not\rightarrow 0$ on the infinite portion of the contour \mathcal{B}_1 and thus we can't close the contour and compute the \int using residues

Solution

- (a) don't use the residues formula for systems with delays
Not too convenient. It defeats the whole purpose
of introducing the $*$ transform!

Solution

(a) don't use the residues formula for systems with delays
Not too convenient. It defeats the whole purpose
of introducing the * transform!

(b) Modify the formula:

It can be shown that if the delay is an integer number of periods
then:

$$E^*(s) = \left[e^{-kTs} E_1(s) \right]^* = e^{-kTs} \sum_{\substack{\text{at poles} \\ \text{of } E_1}} \left\{ \text{Res } E_1(\lambda) \frac{1}{1 - e^{-T(s-\lambda)}} \right\}$$

↑
non delayed
signal

Applying this modified formula to our earlier example we get

$$E_1(s) = \frac{1}{s(s+1)}$$

$$\sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \frac{1}{\lambda(\lambda+1)} \cdot \frac{1}{(1 - e^{-\tau(s-\lambda)})}$$

$$= \frac{1}{(1 - e^{-\tau s})} - \frac{1}{(1 - e^{-\tau(s+1)})}$$

Applying this modified formula to our earlier example we get

$$E_1(s) = \frac{1}{s(s+1)}$$

$$\sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \frac{1}{\lambda(\lambda+1)} \left(\frac{1}{1 - e^{-T(s-\lambda)}} \right)$$

$$= \frac{1}{(1 - e^{-Ts})} - \frac{1}{(1 - e^{-T(s+1)})}$$

$$\Rightarrow E^*(s) = e^{-2Ts} \left[\frac{1}{1 - e^{-0.5s}} - \frac{1}{1 - e^{-0.5(s+1)}} \right] = e^{-s} \left[\frac{1}{1 - e^{-0.5s}} - \frac{1}{1 - e^{-0.5(s+1)}} \right]$$

Applying this modified formula to our earlier example we get

$$E_1(s) = \frac{1}{s(s+1)}$$

$$\begin{aligned} & \sum_{\substack{\lambda=0 \\ \lambda=-1}} \text{Res} \frac{1}{\lambda(\lambda+1)} \left(\frac{1}{1 - e^{-T(s-\lambda)}} \right) \\ &= \frac{1}{(1 - e^{-Ts})} - \frac{1}{(1 - e^{-T(s+1)})} \end{aligned}$$

$$\Rightarrow E^*(s) = e^{-2Ts} \left[\frac{1}{1 - e^{-0.5s}} - \frac{1}{1 - e^{-0.5(s+1)}} \right] = e^{-s} \left[\frac{1}{1 - e^{-0.5s}} - \frac{1}{1 - e^{-0.5(s+1)}} \right]$$

which coincides with our earlier result

Properties of $E^*(s)$

Properties of $E^*(s)$

1) $E^*(s)$ is periodic, with period $T_s = j\omega_s = j\frac{2\pi}{T}$

Properties of $E^*(s)$

1) $E^*(s)$ is periodic, with period $T_s = j\omega_s = j\frac{2\pi}{T}$

proof:

$$E^*(s) = \sum_{k=0}^{\infty} e(kT) e^{-skT}$$

$$E^*(s + j\omega_s) = \sum_{k=0}^{\infty} e(kT) e^{-k[s + j\frac{2\pi}{T}]T} = \sum_{k=0}^{\infty} e(kT) e^{-skT} \underbrace{e^{-jk2\pi}}_1 = E^*(s) \quad \#$$

Properties of $E^*(s)$

1) $E^*(s)$ is periodic, with period $T_s = j\omega_s = j\frac{2\pi}{T}$

proof:


$$E^*(s) = \sum_0^{\infty} e(kT) e^{-skT}$$

$$E^*(s + j\omega_s) = \sum_0^{\infty} e(kT) e^{-k[s + j\frac{2\pi}{T}]T} = \sum_0^{\infty} e(kT) e^{-skT} \underbrace{e^{-jk2\pi}}_1 = E^*(s) \quad \#$$


2) If $E(s)$ has a pole at $s = s_1 \Rightarrow E^*(s)$ has poles at $s = s_1 + jm\omega_s$ $m = 0, \pm 1, \dots$

Note: some property does not apply to zeros of $E^*(s)$

Example: assume that $E(s)$ has a pole at $s=0 \Rightarrow E^*(s)$ poles at $s_m = \pm j n \omega_s$




Example: assume that $E(s)$ has a pole at $s=0 \Rightarrow E^*(s)$ poles at $s_m = \pm j n \omega_s$



proof:
$$E^*(s) = \frac{1}{T} \sum_{n=-\infty}^{n=+\infty} E(s + j n \omega_s) = \frac{1}{T} \left[E(s) + E(s + j \omega_s) + \dots \right]$$

Example: assume that $E(s)$ has a pole at $s=0 \Rightarrow E^*(s)$ poles at $s_m = \pm j n \omega_s$



proof:
$$E^*(s) = \frac{1}{T} \sum_{n=-\infty}^{n=+\infty} E(s + j n \omega_s) = \frac{1}{T} \left[E(s) + E(s + j \omega_s) + \dots \right]$$

if $E(s)$ has a pole at $s=s_1$, the first term contributes a pole at $s=s_1$
 second $s=s_1 - j\omega_s$
 with n $s=s_1 - j n \omega_s$

Example 2: Recall that we have shown that:

$$F(s) = \frac{1}{s} \cdot \frac{1}{(s+1)} \iff F^*(s) = \frac{1}{1-e^{-sT}} - \frac{1}{1-e^{-T(1+1)}}$$

\Downarrow
poles at $s=0$

$$s=-1$$

\Downarrow
poles at $s = \pm jn \frac{2\pi}{T}$ $(e^{jTs}=1)$
 $s = -1 \pm jn \frac{2\pi}{T}$

- Spectrum of a Sampled Signal

- Spectrum of a Sampled Signal

$$\underline{e(t)} \quad / \quad \underline{e^*(t)}$$

• Spectrum of a Sampled Signal

$$\underline{e(t)} \quad \underline{e^*(t)}$$


We'd like to relate the spectrum (i.e. Fourier transform) of $e(t)$ and $e^*(t)$. This will become relevant when we discuss how to reconstruct (if possible) $e(t)$ from $e^*(t)$.


$$\text{Let } e^*(t) = \sum_{k=-\infty}^{k=+\infty} e(t) \delta(t - kT)$$

$$E^*(s) = \frac{1}{T} \sum_{n=-\infty}^{+\infty} E(s + jn\omega_s) = \frac{1}{T} [E(s) + E(s + j\omega_s) + \dots + E(s + jn\omega_s) + \dots]$$


gives the spectrum
of E , shifted by
 $n\omega_s$

gives the spectrum
of $E(s)$








← original spectrum
replicated at
 $\omega_s, 2\omega_s, \dots$



From these plots it follows that we can recover $E(s)$ from $E^*(s)$ only if the highest frequency present in $e(t)$ is smaller than $\frac{\omega_s}{2}$ (the "Nyquist frequency")



We can't have any overlapping:





From these plots it follows that we can recover $E(s)$ from $E^*(s)$ only if the highest frequency present in $e(t)$ is smaller than $\frac{w_s}{2}$ (the "Nyquist frequency")

We can't have any overlapping:



This is the celebrated

Shannon's Sampling Theorem:

This is the celebrated Shannon's Sampling Theorem:


This is the celebrated Shannon's Sampling Theorem:

- A function $e(t)$ which contains no frequency component higher than f_0 is uniquely determined by the values of $e(t)$ at any set of sampling points spaced $T = \frac{1}{2f_0}$

This is the celebrated Shannon's Sampling Theorem:

- A function $e(t)$ which contains no frequency component higher than f_0 is uniquely determined by the values of $e(t)$ at any set of sampling points spaced $T = \frac{1}{2f_0}$

If $e(t)$ has components above the Nyquist frequency we have the following situation:




In the sampled signal the contributions from the frequencies ω_1 and $\omega_2 = \omega_1 - \omega_s$ both show up at ω_1 . This phenomenon is called aliasing.

Implications: 2 sinusoids of different frequencies appear at the same place when sampled \Rightarrow can't tell them apart


Example:

$$f_1(t) = 1$$

$$f_2(t) = \cos 2\pi t$$




If sampled at $f_s = 1$ Hz both yield:



(the same spectrum!)

With 20/20 hindsight this is not surprising:



Related phenomenon: Hidden oscillations:

We can have a signal that does not show up at all when sampled

