

Aquifer depletion exacerbates agricultural drought losses in the US High Plains

Taro Mieno* Timothy Foster† Shunkei Kakimoto‡ Nicholas Brozovic§

Abstract

Aquifer depletion poses a major threat to the ability of farmers, food supply chains, and rural economies globally to use groundwater as a means of adapting to climate variability and change. Empirical research has demonstrated the large differences in drought risk exposure that exist between rainfed and irrigated croplands, but previous work commonly assumes water supply for the latter is unconstrained. In this paper, we evaluate how aquifer depletion affects the resilience of irrigated crop production to drought risk using over 30 years of data on historical corn and soybean yields, production areas, and aquifer conditions for the High Plains region in the United States. We show that aquifer depletion reduces the ability of farmers to sustain irrigated crop yields and production areas in years and locations with large growing season water deficits. Our findings demonstrate that drought-related production losses on irrigated croplands increase non-linearly with aquifer depletion, highlighting the need for proactive aquifer conservation interventions to support adaptation and resilience to future increases in rainfall variability under climate change.

1 Main

Groundwater is an essential input for agricultural production, providing a critical buffer against limited or variable surface water supplies for farmers in many parts of the world [Scanlon et al., 2023]. Globally, agriculture’s dependence on groundwater for irrigation is expected to increase in the future because of higher crop water requirements, more erratic rainfall, and more frequent and extreme drought events caused by climate change [Zhou et al., 2010, Wada et al., 2013, Wada and Bierkens, 2014, Kreins et al., 2015, Flörke et al., 2018]. However, many of the world’s most important aquifer systems have experienced large reductions in storage over recent decades [Wada et al., 2010, Famiglietti et al., 2011, Scanlon et al., 2012, Konikow, 2015, Bierkens and Wada, 2019]. If not addressed, will negatively affect the ability of farmers, food supply chains, and rural economies to use groundwater as a means of adapting to climate change.

Despite widespread alarm about the risks that aquifer depletion poses to crop productivity and food security, empirical evidence is limited about how reductions in groundwater availability will alter farmers capacity to adapt successfully to drought and rainfall variability. Most empirical studies that have assessed impacts of drought on agricultural productivity have focused primarily on rainfed production areas due to the greater exposure of rainfed agriculture to drought-related shocks [Schlenker and Roberts, 2009, Lobell et al., 2014, Schlenker and Lobell, 2010, Zhou et al., 2020, Borgomeo et al., 2020]. Existing studies also demonstrate the importance of existing irrigated areas or future irrigation intensification and expansion to mitigate negative impacts of drought [Kuwayama et al., 2019, Zipper et al., 2016, Zhu and Burney, 2022, Zhu et al., 2022, Lu et al., 2020, Davis et al., 2019, Li and Troy, 2018]. In contrast, most research assessing impacts of aquifer depletion on resilience of irrigated farmland to drought has relied on simulation modeling [Foster et al., 2015b, Cotterman et al., 2018, Kahil et al., 2015, Yoon et al., 2021, Rouhi-Rad et al., 2020c], with more limited work that attempts to empirically evaluate change in drought-related production losses and risks on irrigated lands as a function of changing aquifer storage [Jain et al., 2021, Suter et al., 2021].

*Department of Agricultural Economics, University of Nebraska Lincoln, tmieno2@unl.edu

†Department of WWW, University of Manchester, timothy.foster@manchester.ac.uk

‡Department of Applied Economics, University of Minnesota, kakim002@umn.edu

§Department of Agricultural Economics, University of Nebraska Lincoln, nbrozovic@nebraska.edu

Two pathways exist by which aquifer depletion will impact farmers’ ability to access and use groundwater as a buffer against drought and rainfall variability [Foster et al., 2015a]. First, depletion increases the energy requirements and costs of pumping water, potentially making it less economically viable for farmers to satisfy crop water needs [Mieno and Brozović, 2017, Bhattarai et al., 2021]. Second, depletion also reduces the aquifer thickness of an aquifer, lowering the aquifer’s transmissivity (i.e., the potential rate of water supply to wells). This latter effect reduces the rate at which water can be pumped and applied to a field as irrigation [Konikow and Kendy, 2005, Foster et al., 2014, Hrozencik et al., 2017]. If the reduction in well yields is substantial - for example as would occur when a large proportion of aquifer thickness is depleted [Hecox et al., 2002, Korus and Hensen, 2020] - then a farmer may be unable to extract sufficient water to meet crop water demands, in particular during periods of limited rainfall and in the peak of the growing season when crops are most sensitive to water deficits [Foster et al., 2015b, Rouhi-Rad et al., 2020a]. The combined effect of declining well yields and increasing pumping costs could force farmers to reduce irrigation depths, resulting in greater risk of drought-related crop stress. Alternatively, or in addition, a farmer may be forced to limit irrigated area to ensure crop water demands can be met, leading to reductions in total crop production [Foster et al., 2014, Rouhi-Rad et al., 2020b]. Understanding how these two responses vary as a function of aquifer conditions is crucial to guide groundwater management planning decisions, in particular given projected increases in frequency and intensity of droughts under climate change [Ukkola et al., 2020, Chiang et al., 2021, Cook et al., 2020].

In this paper, we present empirical evidence about how aquifer depletion affects farmers ability to effectively and reliably buffer crops against drought using data on historical crop yields, production areas, and aquifer conditions for the High Plains region in the United States. Our analysis extends previous research on the impacts of drought on agricultural production by evaluating how drought-related production losses are influenced by severity of groundwater scarcity, moving beyond prior binary comparisons of rainfed and irrigated production [Schlenker and Roberts, 2009, Lobell et al., 2014, Lu et al., 2018]. We also provide new empirical insights about the relative roles of extensive (i.e., irrigated area) and intensive (i.e., per-area irrigation rates) margin water use adjustments in determining drought-related production losses, which previously had only been demonstrated through theoretical modeling [Foster et al., 2014, 2017, Rouhi-Rad et al., 2020b]. Our findings provide important evidence of the value of preserving aquifer storage as a buffer against drought events. In doing so, our analysis adds weight to the need to improve groundwater resource conservation and sustainability to enable adaptation to climate change in groundwater-dependent agricultural systems globally [Jain et al., 2021, Scanlon et al., 2023].

2 Results

2.1 Aquifer depletion increases sensitivity of crop yields to water deficits

Figure 1 shows the estimated crop yield response to water deficit for rainfed production and groundwater-irrigated production at different levels of aquifer thickness categories for corn (panel A) and soybean (panel B)¹. The histogram of annual water deficit is presented at the top for each crop. In wetter years with low crop irrigation requirements (i.e., negative values of water deficit), rainfed and irrigated yields are very similar for both corn and soybean. As drought severity increases (i.e., as water deficit increases), rainfed yields of both crops decline significantly whereas irrigated yields are increasing or stable as deficit increases. This result reflects the positive benefits irrigation provides both as a buffer against precipitation variability, and as a means of raising overall yields through increased evapotranspiration and associated biomass accumulation, reductions of heat-related stressors, and other factors [Zhu and Burney, 2022, Li et al., 2020].

We now focus on comparing yield response to water deficit among the three aquifer thickness categories (all irrigated). First, for water deficits above around 400mm, yield changes become negative for counties with the lowest levels of aquifer thickness (i.e., 1st quantile in Figure 1). This likely reflects the restrictions that lower aquifer thickness and associated reductions in well yields place on farmers ability to fully meet crop water requirements during periods of more severe or extended water deficits [Rouhi-Rad et al., 2020b, Foster et al.

¹Confidence intervals were not presented to avoid over-crowding the figures. Please see Appendix A for estimated yield response curves with 95 percent confidence interval.

[2014]; Hrozencik et al., 2017]. In contrast, yield continues to increase gradually as water deficit increases beyond 400 mm for counties with the highest level of aquifer thickness.

Figure 2 shows the difference in yield of the second and third (highest) aquifer quantiles relative to the first (lowest) quantile with the 95% confidence interval for corn and soybean. For corn, in years with water deficit greater than about 700 mm, the highest aquifer thickness group has statistically significantly higher yields than the lowest aquifer thickness group. At an extreme water deficit level of 950 mm (average water deficit in 2012, the most extreme drought in our historical record), the difference in yields between the smallest and largest aquifer thickness groups is 1.15 tonne/ha. Unlike corn, irrigated soybean yields are not statistically distinguishable among the three aquifer thickness groups. This is likely because soybean is not as water demanding a crop as corn, meaning that even counties with lower saturated thickness can meet the majority of crop water demands as well yield is not a significantly binding constraint on irrigation management decisions.

At low water deficits (below 200mm), Figure 2 illustrates that counties with the highest levels of aquifer thickness (i.e., 3rd quantile in Figure 1 experience statistically lower yields for corn and soybean than counties with the lowest level of aquifer thickness (1st quantile). In case of soybean, irrigated yields for the lowest aquifer thickness category converge to levels comparable to those obtained from rainfed production. These counter-intuitive results reflect a combination of two factors. First, in years with small water deficits there is less need for irrigation and hence irrigated and rainfed yield differences are reduced. Second, years with small or even negative water deficits are associated with wetter conditions during the growing season, which can negatively impact yield if waterlogging occurs leading to greater risks of physical crop damage (e.g., lodging of grains), delays to planting/harvesting, nutrient leaching, and restricted root growth in waterlogged soils [Li et al., 2019]. Declines in irrigated yields could be indicative of over-irrigation by farmers, which has been shown to occur in the HPA region in wet years [Foster et al., 2019, Gibson et al., 2017, 2019], exacerbating these risks. However, the small declines in irrigated yields at low or negative levels of water deficit may also reflect the fact that wetter years are typically cooler, reducing positive heat effects on crop growth and yield formation.




Figure 1: The impact of water deficit and aquifer thickness on rainfed and irrigated per-area yields of corn and soybean in US High Plains. Irrigated yield responses to water deficits are disaggregated by aquifer thickness level, considering three quantile groups that capture the range of aquifer thickness values in our sample: (i) 9 m to 30.1 m; (ii) 30.1 m to 67.2 m; and (iii) 67.2 m to 215.8 m. The histograms on the top of each panel show the distribution of county-year water deficit values included in regressions for each crop.




Figure 2: The difference in estimated county-level irrigated corn (top panels) and soybean (bottom panels) yields for counties in the second and third (largest) aquifer thickness quantiles relative to the first (smallest) aquifer thickness quantile.

2.2 Aquifer depletion substantially reduces the share of agricultural land under irrigation

While aquifer thickness has a statistically significant impact on per-area irrigated crop yields in drought conditions (water deficit greater than 700mm), the magnitude of the yield impact is small when compared to differences between irrigated and rainfed production for the same weather conditions. One explanation for this result is that farmers may choose to retain a smaller proportion of land in irrigated production, so that limited groundwater pumping capacity can be used to adequately buffer crops against drought on land that remains under irrigation. Reductions in irrigated areas as an aquifer is depleted have been demonstrated theoretically [Rouhi-Rad et al., 2020b, Foster et al., 2014, Hrozencik et al., 2017, Deines et al., 2020], but there has been little empirical research on farmers' irrigated area choices or how these are affected by combinations and magnitudes of weather risks and groundwater scarcity.

Figure 3 shows the estimated relationship between aquifer thickness and the share of production area that is irrigated for corn and soybean after accounting for spatio-temporal variations in weather conditions and drought risk exposure across the aquifer. Figure 3 demonstrates that aquifer thickness has meaningfully large impacts on the share of irrigated production for both corn and soybean. As aquifer thickness declines, the share of irrigated acres for corn decreases from around 0.86 (86%) for an average county-level aquifer thickness of 130m to 0.61 (61%) for an average county-level aquifer thickness of 10m (See Appendix B for formal

statistical testings). Reductions in the share of production area under irrigation is also evident for soybean, although the effect is less pronounced than for corn, potentially due to soybean's lower water requirements and susceptibility to drought relative to corn [Zipper et al., 2016, Ruess et al., 2022]




Figure 3: The estimated impact of aquifer thickness on the share of agricultural production under irrigation for corn and soybean producing counties in the US High Plains (Note: shaded area represents 95 percent confidence interval)

2.3 Aquifer depletion increases overall drought-related production losses by up to 25%

The overall influence of aquifer depletion on drought-related production losses is a function of both per-area irrigated yield and irrigated area changes highlighted in Sections 2.1 and 2.2. Combining both responses, Figure 4 shows the relationship between water deficit and average per-area production (weighted-average of irrigated and rainfed yields, where the weights are production area shares) for corn and soybean at four values of aquifer thickness: 10, 50, 90, and 130 meters. While similar trends are observed as in Figure 1, the difference in production loss caused by high water deficits among the aquifer thickness levels are much more pronounced due to lower estimated irrigated area share for smaller aquifer thickness values. For example, for an aquifer thickness of 10 m, reductions in average yields occur for smaller water deficits than are required to trigger reductions in per-area irrigated yields alone. Similarly, the overall magnitude of yield reduction is greater for average yields than for per-area irrigated yields. This is because low aquifer thickness values have a greater estimated share of crop area under rainfed production, which magnifies average yield declines with increasing water deficits.

Figure 5 shows the difference in average per-area productivity for 50 m, 90 m, and 130 m aquifer thickness levels relative to an aquifer thickness of 10 m for corn and soybean. For corn, aquifer thickness does not have a significant impact on productivity when water deficits are below about 350mm. However, beyond this threshold, higher levels of water deficit result in noticeable and statistically significant reductions to average corn production. This reflects the combination of lower irrigated yields under drought conditions, and, more

importantly, reduced overall productivity due to smaller share of land under irrigation in areas and years where aquifer thickness is lower. Similar effects are also observed for soybean, although reductions in average production only occur at a higher deficit threshold (greater than around 600mm) and are smaller in overall magnitude. For example, for a seasonal water deficit of 950mm (average water deficit in 2012, the most extreme drought in our historical record), average per-area corn and soybean productivity are about 25% and 15% lower, respectively, when differentiating between areas with the 130m and 10m aquifer thicknesses.

Figure 4 and Figure 5 also demonstrate that impact of aquifer thickness on average per-area production is highly non-linear at high seasonal water deficits. At a water deficit of 950mm, for example, reducing from 130m aquifer thickness to 90m (decline of 40m) would result in only 0.11 tonne/ha of production loss for corn. This negligible difference in average yield from the decline is due to the insensitivity of the share of irrigated acres to a decline in aquifer thickness when the initial aquifer thickness level is high as can be seen in Figure 3. Reducing from 90m aquifer thickness to 50m (decline of 40m) would result in 0.77 tonne/ha of production loss for corn, with a further drop from 50m aquifer thickness to 10m (decline of 40m) leading in 2.09 tonne/ha of production loss for corn. Similar but less pronounced effects are also observed for soybean. This non-linear impact of aquifer thickness is due to the influence of aquifer thickness on the share of irrigated corn acres (Figure 3), which implicitly reflects how aquifer thickness alters well yields and, in turn, farmers ability to effectively and reliably buffer crops against water deficits during the growing season.




Figure 4: Average productivity of corn and soybean in the US High Plains for different levels of water deficit and aquifer thickness, considering both per-area yield responses and farmers choices about irrigated area shares in response to different water deficit and aquifer conditions. See Appendix A for estimated average yield response curves with 95 percent confidence intervals displayed, which are omitted here for visual clarity.




Figure 5: Difference in average yield of corn and soybean for aquifer thickness levels of 50 m, 90 m, and 130 m, relative to a baseline aquifer thickness of 10 m. Shaded areas denote 95 percent confidence interval, which when greater than zero indicate average yields for the given aquifer thickness level are statistically significantly greater than for an aquifer thickness of 10 m at the 5 percent level of significance.

3 Discussion

We demonstrate the important role of aquifer thickness in determining farmers ability to successfully buffer crops against drought risks in groundwater irrigated agricultural systems such as the High Plains of the United States. Previous research focused on impacts of drought on agricultural systems has largely focused either on risks for rainfed production [Schlenker and Roberts, 2009, Lobell et al., 2014, Schlenker and Lobell, 2010, Zhou et al., 2020, Borgomeo et al., 2020] or for irrigated lands where water supply constraints are assumed to be minimal [Kuwayama et al., 2019, Zipper et al., 2016, Zhu and Burney, 2022, Zhu et al., 2022, Lu et al., 2020, Davis et al., 2019, Li and Troy, 2018, Luan et al., 2021]. Our analysis adds new empirical insights to this literature by quantifying how reductions in aquifer thickness increase the vulnerability of irrigated agricultural production areas to drought events. Critically, our findings suggest that commonly adopted binary definitions of agricultural land areas into rainfed and irrigated categories are likely to underestimate exposure of crop production to drought risks, in particular in areas where water supplies for irrigation are insufficient to fully meet crop water needs in all years due to constraints imposed by hydrology, policies, and/or economics.

An important implication of our findings is that depletion of groundwater resources is likely to increase significantly the vulnerability of agricultural production to current and future drought risks. Climate change is projected to increase the frequency and severity of extreme drought events in many parts of the world [Ukkola et al., 2020, Chiang et al., 2021, Cook et al., 2020], including in parts of Kansas and Texas [Bradford et al., 2020, Cook et al., 2022, Mullens and McPherson, 2019] where the High Plains Aquifer continues to experience prolonged and significant declines in aquifer thickness [Scanlon et al., 2012, Haacker et al., 2016, Cotterman et al., 2018]. Similar trends of increasing drought risks coupled with declining aquifer storage have been reported in other major groundwater irrigated farming systems worldwide [Wada et al., 2010, Döll et al., 2014, Famiglietti, 2014, Feng et al., 2018, Bierkens and Wada, 2019]. Our findings suggest that these simultaneous changes in aquifer conditions will amplify impacts of future climate change on crop production, with associated ramifications for resilience of supply chains, commodity markets, rural economies, and food security.

Our findings also provide useful guidance for water managers and policymakers seeking to address unsustainable groundwater use in the US High Plains and globally. Consistent with prior theoretical modeling, we demonstrate that the relationship between aquifer thickness and climatic water deficits during the growing season is highly non-linear when accounting for farmers decisions about irrigated production areas. This indicates that sustainable groundwater conservation efforts [MacEwan et al., 2017, Butler et al., 2020, Elshall et al., 2020] should be targeted in space and time to avoid depletion exceeding critical thresholds which impair farmers’ ability to effectively buffer production against drought. Analysis such as ours can provide valuable evidence about these critical thresholds and tipping points to guide proactive management of groundwater stocks, and, in doing so, limit economically damaging transitions from irrigated to rainfed production [Foster et al., 2017, Deines et al., 2020]. For example, several groundwater management agencies in the Texas portion of the HPA have in recent years set out targets to ensure 50% of current aquifer storage and thickness remains in 50 years time (so called 50/50 rule) [Closas and Molle, 2018]. The effectiveness of such targets could likely be improved substantially by setting conservation targets based on empirical evidence about the level of aquifer thickness that is required to ensure adequate levels of drought protection, both now and with future climate change.

Several opportunities exist for future research to build upon the methods and analyses presented here. First, due to the limited resolution of our observational data, we were only able to estimate the impacts of different aquifer thickness categories on drought risk. Availability of sub-county crop yield data, for example from satellite remote sensing and/or in-situ monitoring [Edreira et al., 2020, Deines et al., 2021], would allow models of the continuous effect of aquifer thickness on drought risk to be developed. Unobserved variations in aquifer properties will also introduce noise into the relationships between aquifer thickness, well yields, and drought risk. For example, a thinner aquifer interval composed of coarse sands and gravels may be able to supply comparable or higher well yields than a thicker aquifer dominated by fine sands and silt deposits, and hence offer greater resilience to water deficits despite similar or larger aquifer thickness [Butler et al., 2013, Korus and Hensen, 2020]. These measurement errors in the relationship between aquifer thickness and well yield will lead to attenuation bias [Bound and Krueger, 1991, Hyslop and Imbens, 2001], suggesting

that our analysis is likely to be an underestimate of the true impacts of groundwater depletion on drought risk. Nonetheless, our analysis shows that - even in the presence of these measurement uncertainties - we are still able to identify statistically significant increases in drought risk exposure for irrigated agriculture as a function of declining saturated thickness. Improved data on spatial variations in aquifer properties and greater monitoring of real-world changes in well capacities alongside traditional water level measurements would help to further refine understanding of patterns of drought vulnerability across the HPA. This would enable more precise estimates to be made of critical tipping points in the resilience of irrigated agriculture to drought, which are needed to effectively design and target groundwater conservation policies.

Second, our analysis considered the effects of aquifer thickness on farmers decisions to switch from irrigated to rainfed production, but did not assess other potential responses to aquifer depletion such as changing crop types or irrigation management practices. Switching to more drought resistant crops or varieties may enable farmers to maintain irrigated production at lower levels of aquifer thickness, and has been shown to be an important response to increasing agricultural water scarcity and abstraction policies in other studies [Bhattarai et al., 2021, Deines et al., 2019, Manning et al., 2017]. Improvements to irrigation scheduling practices and technologies, such as have been implemented as part of the SD6-LEMA (Local Enhanced Management Area) in Kansas [Deines et al., 2019, Glose et al., 2022] and in parts of Texas [Mrad et al., 2020] and which are proposed as part of California’s Sustainable Groundwater Management Act (SGMA) [Berbel and Esteban, 2019, Lubell et al., 2020], could also help to improve the efficiency of irrigation and, hence, the ability to more effectively meet crop water demands with lower well capacities. Such interventions may reduce - but not eliminate - increases in drought risk caused by lower or declining aquifer thickness, and could also help to mitigate against potential yield losses caused by over-use of water in wetter years via raising awareness of efficiency saving potentials [Foster et al., 2019] and the importance of conservation [Marston et al., 2022]. However, risks also exist that irrigation efficiency improvements will increase net extraction, ultimately causing a rebound effect that exacerbates future aquifer depletion and drought risks [Grafton et al., 2018, Pérez-Blanco et al., 2021]. Future research should explore potential strategies that farmers could adopt to lessen the combined impacts of drought and aquifer depletion, alongside efforts to curb unsustainable groundwater extraction and climate change.

Finally, we focus on the impacts of aquifer thickness on agricultural resilience to seasonal water deficits. However, where water deficits are distributed unevenly during the season, negative impacts of aquifer depletion may be even larger. This is because reductions in well yields caused by aquifer depletion will constrain farmers ability to meet peak crop water demands, increasing yield losses compared with if water deficits had been distributed evenly throughout the season [Ortiz-Bobea et al., 2019]. Furthermore, irrigation also has broader benefits for crops such as reducing heat-related stress through cooling of land surface temperatures [Adegoke et al., 2003, Bonfils and Lobell, 2007, Lobell and Bonfils, 2008, Zhu and Burney, 2022]. Reductions in aquifer thickness therefore may have other negative impacts on irrigated crop productivity, if constraints to water supply due to aquifer depletion are significant enough to limit cooling benefits of irrigation during the growing season.

4 Methods

4.1 Data-sets

Our study focuses on the High Plains Aquifer (HPA) in the Central United States. Irrigated agriculture is integral to the landscape and economy of the HPA, with the region accounting for the majority of grain production in the United States valued in the billions of dollars annually [Smidt et al., 2016, Fenichel et al., 2016]. To evaluate the impacts of aquifer depletion on agricultural production overlying the HPA, we rely on data from several sources, which are described below.

County-level annual irrigated and rainfed yield data for corn and soybean for 1985 through 2016 were obtained from the U.S. Department of Agriculture’s National Agricultural Statistics Service (USDA-NASS). USDA-NASS does not report irrigated yields disaggregated by the source of water used for irrigation. However, it is known that groundwater is the main source of water for irrigation for counties that overlie the HPA due to the limited availability of surface water resources in this area. Therefore, we retain both rainfed and

irrigated yield observations for counties where greater than 75% of the county area overlies the boundary of the HPA. For counties that do not meet this condition, only rainfed yield observations are obtained to avoid the possibility that irrigated production may be only partly dependent on use of groundwater from the HPA. We also include additional rainfed yield observations for counties that are outside the HPA, but within the main states covering the aquifer (i.e., South Dakota, Wyoming, Colorado, New Mexico, Nebraska, Kansas, and Texas), to enable more accurate estimation of the impact of drought for rainfed production as a point of comparison to our main irrigated crop yield analyses.

Meteorological data needed to estimate spatial and temporal variations in drought conditions across the HPA were obtained from the gridMET dataset [Abatzoglou, 2013]. gridMET is a gridded daily weather dataset with a spatial resolution of $\sim 4\text{km}$, and has been used in a range of climate impacts studies for agriculture and other sectors in the United States [Abatzoglou and Williams, 2016, Pereira et al., 2015, Crane-Droesch, 2018, Venkatappa et al., 2021, Zhu et al., 2019]. Daily precipitation values were extracted for each gridMET cell overlapping counties in our sample. Daily values of maximum and minimum temperature, surface radiation, wind speed, and humidity were also obtained from gridMET in order to calculate daily reference crop evapotranspiration for each grid cell. From these data, we computed the seasonal water deficit ($WB_{i,t}$) for each grid cell by subtracting total precipitation from total reference evapotranspiration for the period May through September, which represents the typical growing season for corn and soybean in the HPA region. Seasonal water balance estimates were subsequently aggregated to the county-level to match our agricultural production data using an area-weighted average based on the proportion of the county covered by each overlapping gridMET cell. Figure 6 presents annual water deficit by year for the corn and soybean data, illustrating major drought events in the region in years such as 2000, 2002, and 2012. The importance of irrigation is clearly evident in years with large water deficits, with rainfed yields dipping substantially whereas irrigated yields are, on average, largely unaffected (Figure 6).




Figure 6: Historical rainfed yield, irrigated yield, and water deficit

As a proxy for the extent of aquifer depletion we use annual estimates of aquifer thickness generated by existing research [Haacker et al., 2016, 2023]. These data provide annual estimates of the aquifer thickness of the aquifer at 250m resolution since 1935. Annual aquifer thickness maps were derived based on interpolation of 14,000 to 20,000 individual observations of water levels per year, along with detailed maps of both the land surface elevation and bedrock elevation of the HPA. Aquifer thickness estimates for each year were aggregated to the county level using an area-weighted average to match the resolution of agricultural production data. Aquifer thickness values were only estimated for counties with irrigated yield observations. Furthermore, during the aggregation process, any aquifer cells within a county that had a aquifer thickness value lower than 9 m were removed before aggregation to reflect the fact that aquifer thickness lower than this threshold are not considered to be viable for extraction and therefore it is unlikely these areas contribute groundwater for irrigation production [Fenichel et al., 2016, Haacker et al., 2016, Deines et al., 2020]. Figure 7 presents the distribution of aquifer thickness across our study area for counties included in corn and soybean analyses. aquifer thickness levels vary substantially over space and time, which we will exploit in our analysis to quantify how aquifer influences farmers' exposure and responses to drought.

306 These data were combined to form four sets of data for regression analysis: (1) per-acre corn yield (8773
 307 observations), (2) per-acre soybean yield (5977 observations), (3) share of irrigated corn (2686 observations),
 308 and (4) share of irrigated soybean (2014 observations).




Figure 7: Map showing county-average aquifer thickness in 2016 for each county included in our corn and soybean regression analyses. Counties highlighted in grey are rainfed only, and the red polygon denotes the boundary of the High Plains Aquifer system

4.2 Regression Models

Aquifer depletion may impact resilience of irrigated crop production to drought through two main channels: (1) a reduction in per-area irrigated yields due to the economic or physical inability to meet crop water requirements; and (2) a reduction in the area of land irrigated, resulting in reduced production due to lower yields obtained through rainfed agriculture.

To assess the first of these impacts, we quantify the impact of seasonal water deficits (WB) during the growing season (May through September) on county-level rainfed and irrigated yields (Y) using a hierarchical generalized additive model, where irrigated yield response to seasonal water deficits is conditioned on the level of aquifer thickness (AT) in the county in a given year. Our regression model is given in Equation (1), with the same model formulation used for corn and soybean:

$$y_{i,t} = \sum_{j=1}^4 \sigma_j C_j + \sum_{j=1}^4 \sum_{k=1}^K \beta_{j,k} \phi_{j,k}(WB_{i,t}) \cdot C_j + \alpha_i + \lambda_t + v_{i,t} \quad (1)$$

where county and year are indicated by subscripts i and t , respectively, y is crop yield, and WB is seasonal water deficit. C_j is a dummy variable that takes 1 if observation i is under category C_j , 0 otherwise. The observations are grouped into four categories: rainfed production, and county-years with aquifer thickness falling in the $[0 - 33.3]$, $[33.3 - 66.7]$, and $[66.7 - 100]$ quantile ranges of aquifer thickness. The three irrigated production groups have about the same number of observations, which ensures statistical stability in estimating yield response to drought by category. Spline functions are used to capture the potential non-linearity of the impact of water deficit on yield in a flexible manner, with σ_j represents the category specific intercept, $\phi_{j,k}$ k th spline basis function, and β_k is its coefficient (see Appendix D for further explanations on how spline-based regression works). To control for time-invariant heterogeneity across counties and yearly shocks, county fixed effects (α_i) and year fixed effects (λ_t) are also included and standard error estimation is clustered by county.

To estimate the additional impact of aquifer thickness on the share of total acres that are irrigated in a county, a fractional logit model under the generalized additive model framework is used in which the impact of aquifer thickness, average water deficit, and their interactions are estimated without the prior requirement to assume any functional forms. This allows for flexible representation of the non-linear relationships between drought risks, aquifer conditions, and agricultural land use, including potential thresholds in farmers irrigated area choices under water supply constraints that have been noted in prior work [Foster et al., 2015b,a, Hrozencik et al., 2017]. The model is given in Equation (2) below:

$$\log\left(\frac{s_{i,t}}{1 - s_{i,t}}\right) = \beta_0 + \sum_k^K \alpha_k \phi_k(WB_i) + \sum_l^L \beta_l \tau_l(AT_{i,t}) + X_i + \theta_{s,t} + \mu_t + v_{i,t} \quad (2)$$

where $s_{i,t}$ is the share of irrigated production. The impact of average water balance is captured by $\sum_k^K \alpha_k \phi_k(WB_i)$, where $\phi_k(\cdot)$ is the k th spline basis function and α_k is its coefficient. The impact of aquifer thickness is captured in a similar way with $\tau_l(\cdot)$ and β_l being the l th spline basis function and its coefficient, respectively. County-level soil characteristics including sand percentage, silt percentage, and water holding capacity, are represented by X_i . State-year fixed effects and error term are represented by $\theta_{s,t}$ and $v_{i,t}$, respectively. Unlike for our regression models of per-area yields, we do not include county fixed effects to avoid eliminating the majority of variation in the aquifer thickness variable. The vast majority of variation in aquifer thickness is cross-sectional (across counties), rather than over time. Including county fixed effects would effectively eliminate the cross-sectional variation in aquifer thickness. Instead, through robustness checks, we demonstrate that our results are consistent when using state fixed effects. A formal discussion of these results and trade-offs associated with the choice of fixed effects is provided in Appendix C.

In the final step of our analysis, we then estimate the average crop yield as a function of aquifer thickness and seasonal water deficit. The average crop yield is estimated as the average of rainfed and irrigated yields estimated on the basis of Equation (1), weighted by the estimated share of total acres that are irrigated in

the county estimated using Equation (2). Comparing the estimated average crop yield across different levels of aquifer thickness for a given seasonal water deficit provides an indication of the effect of aquifer conditions on drought impacts, considering impacts of drought through changes in both per-area irrigated yields and the area of land irrigated. Standard errors of this analysis are estimated using bootstrap methods clustered by county. Note that irrigated area share and average crop yield estimates use data only from counties that have both irrigated and rainfed production observations. Counties with solely rainfed production observation data (Section 4.1) are excluded from these analyses, and are used solely for estimating impacts of drought on per-area crop yields (Equation (1)).

4.3 Data and Code Availability

All the codes used for this study are available at the following Github repository (link). All analyses were performed in R [R Core Team, 2021] and all the raw data-sets used in this study are publicly accessible. The boundary shape file of the High Plains aquifer was obtained from the U.S. Geological Survey at <https://water.usgs.gov/GIS/metadata/usgswrd/XML/ds543.xml#stdorder>. This data is used for Figure 7. Corn and soybean yield data were obtained from the U.S. Department of Agriculture’s National Agricultural Statistics Service (USDA-NASS), available at https://www.nass.usda.gov/Quick_Stats/. The R tidyUSDA package [Lindblad, 2022] was used to automatically download the relevant data. This data is used for Figure 6 and the regression analysis. Weather data (precipitation, temperature, and evapotranspiration) was obtained from the gridMET dataset [Abatzoglou, 2013], which is available at <https://www.climatologylab.org/gridmet.html>. This data is used for Figure 6 and the regression analysis. The aquifer thickness data were obtained from <https://www.hydroshare.org/resource/7d925c7944244032af98c9ed20c22db6/> [Haacker et al., 2023]. This data is used for Figure 7 and the regression analysis. Soil characteristics data was obtained from the Soil Survey Geographic Dataset, which is used for regression analysis. The R soilDB package [Beaudette et al., 2023] was used to download the data. Regression analyses are implemented using the R fixest package [Bergé, 2018]. Other key R packages include ggplot2 [Wickham, 2016] for creating plots, data.table [Dowle and Srinivasan, 2022] for data wrangling, and sf [Edzer, 2018] for spatial data operations. The data used for the regression analysis can be generated using our codes by following the instructions available at the linked Github repository.

A Uncertainties in estimated impacts of water deficit on irrigated and average yield

In the main article, the confidence intervals of the estimated yield responses to water deficit were not presented to enable easier comparison between response curves. This section provides visualization of yield response with the 95% confidence interval to convey the degree of uncertainty in our estimation. Figures A.1 and A.2 present the impact of water deficit on crop yield by aquifer thickness category with the 95% confidence interval for irrigated corn and soybean, respectively. Figure A.3 presents the impact of water deficit on average crop yield (irrigated and rainfed combined) by aquifer thickness category for corn and soybean with the 95% confidence interval.




Figure A.1: The impact of water deficit and aquifer thickness on rainfed and irrigated per-area irrigated yields of corn in US High Plains (solid black line) with 95 percent confidence interval (grey shaded area)




Figure A.2: The impact of water deficit and aquifer thickness on rainfed and irrigated per-area irrigated yields of soybean in US High Plains (solid black line) with 95 percent confidence interval (grey shaded area)




Figure A.3: Average productivity of corn and soybean in the US High Plains for different levels of water deficit and aquifer thickness (solid black line) with 95 percent confidence interval (grey shaded area)

B Test of the Impact of Aquiefr Thickness on the Share of Irrigated Acres

Figure C.1 presents the difference in the share of irrigated acres at different levels of aquifer thickness relative to the aquifer thickness level of 130m.




Figure C.1: Decline in the share of irrigated acres as a function of aquifer thickness relative to aquifer thickness level of 130m (The shaded areas represent the 95 percent confidence interval)

C County Fixed Effects in Irrigation Share Regression

Figure B.1 presents the histogram of aquifer thickness after demeaning them by its overall average (A), state average (B), and county average (C) to gain insights into how much variation for identification is left under no county or state fixed effects (A), state fixed effects (B), and county fixed effects (C). As you can see in the figure, while including state fixed effects do not reduce the total variation in aquifer thickness, including county fixed effects dramatically reduces its total variation. Consequently, we found that including county fixed effects results in extremely inaccurate estimates (high standard error). For this reason, we just included state fixed effects.




Figure B.1: Comparison of variations in aquifer thickness after demeaning by overall average, county average, and state average.

Figure B.2 compares the impact of aquifer thickness on the share of irrigated acres for corn and soybean with different sets of fixed effects. For both corn and soybean, the negative impact of reduction in aquifer thickness on the share of irrigated acres are virtually the same across the sets of fixed effects. Importantly, adding state fixed effects, which control for unobservables at the state level, does not alter the estimated impacts of aquifer thickness on share of acres irrigated. This indicates that, even though there may be unobservables that are spatially correlated with aquifer thickness beyond soil characteristics, their correlation is unlikely to cause significant bias to our estimation and findings.




Figure B.2: Comparison of estimated impact of aquifer thickness on the share of irrigated acres with and without state fixed effects

D Spline basis functions and GAM estimation

We used generalized additive models (GAM) for our regression analysis. In GAM, the impact of a variable on the dependent variable can be expressed as a series of spline basis functions and their coefficients. Suppose you are interested in modeling the following model.

$$y = f(x) + \mu \quad (3)$$

Where $f(x)$ represents the impact of variable x on y . The shape of $f(x)$ is not known to the researcher. GAM attempts to represent $f(x)$ with a series of spline basis functions and their coefficients like below.

$$y = \sum_{k=1}^K \beta_k \phi_k(x) + \mu \quad (4)$$

Where $\phi_k(\cdot)$ is the k th spline basis function and β_k is its coefficient. Figure C.1 provides a visual illustration of a series of thin-plate spline basis functions.




Figure C.1: Example of a series of thin-plate spline basis functions

Different coefficients (or namely weights) given to the spline basis functions result in different shapes of the impact of x on y . Figure C.2 presents two examples of two different sets of coefficients and their

416 resulting representation of $f(x)$. In the first case, $\{\beta_1, \dots, \beta_5\} = \{60, 60, 60, 60, 60\}$. In the second case,
 417 $\{\beta_1, \dots, \beta_5\} = \{20, -20, -40, 40, 60\}$. Among all the possibilities of the combinations of the coefficients,
 418 GAM finds the set of coefficient values that fits the data best. Notice that unlike non-parametric regression,
 419 GAM is actually a linear in parameter model and estimate β_k to fit the data best.




Figure C.2: Example of a function represented by different coefficient values

References

- J.T. Abatzoglou. Development of gridded surface meteorological data for ecological applications and modelling. *International Journal of Climatology*, 33(1):121–131, 2013. doi: <https://doi.org/10.1002/joc.3413>. URL <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.3413>.
- J.T. Abatzoglou and A.P. Williams. Impact of anthropogenic climate change on wildfire across western US forests. *Proceedings of the National Academy of Sciences*, 113(42):11770–11775, 2016.
- J.O. Adegoke, R.A. Pielke Sr, J. Eastman, R. Mahmood, and K.G. Hubbard. Impact of irrigation on midsummer surface fluxes and temperature under dry synoptic conditions: A regional atmospheric model study of the us high plains. *Monthly Weather Review*, 131(3):556–564, 2003.
- D. Beaudette, J. Skovlin, S. Roecker, and A. Brown. *soilDB: Soil Database Interface*, 2023. URL <https://CRAN.R-project.org/package=soilDB>. R package version 2.7.7.
- J. Berbel and E. Esteban. Droughts as a catalyst for water policy change. analysis of Spain, Australia (MDB), and California. *Global Environmental Change*, 58:101969, 2019.
- R. Bergé. Efficient estimation of maximum likelihood models with multiple fixed-effects: the R package FENmlm. *CREA Discussion Papers*, (13), 2018.
- N. Bhattarai, A. Pollack, D.B. Lobell, R. Fishman, Balwinder-Singh, A. Dar, and M. Jain. The impact of groundwater depletion on agricultural production in India. *Environmental Research Letters*, 16(8):085003, 2021.
- M.F. Bierkens and Y. Wada. Non-renewable groundwater use and groundwater depletion: a review. *Environmental Research Letters*, 14(6):063002, 2019.
- Céline Bonfils and David Lobell. Empirical evidence for a recent slowdown in irrigation-induced cooling. *Proceedings of the National Academy of Sciences*, 104(34):13582–13587, 2007.
- E. Borgomeo, H.F. Khan, M. Heino, E. Zaveri, M. Kumm, C. Brown, and A. Jägerskog. Impact of green water anomalies on global rainfed crop yields. *Environmental Research Letters*, 15(12):124030, 2020.
- J. Bound and A.B. Krueger. The extent of measurement error in longitudinal earnings data: Do two wrongs make a right? *Journal of Labor Economics*, 9(1):1–24, 1991.
- J.B. Bradford, D.R. Schlaepfer, W.K. Lauenroth, and K.A. Palmquist. Robust ecological drought projections for drylands in the 21st century. *Global Change Biology*, 26(7):3906–3919, 2020. doi: <https://doi.org/10.1111/gcb.15075>.
- J.J. Butler, R.L. Stotler, D.O. Whittemore, and E.C. Reboulet. Interpretation of water level changes in the High Plains aquifer in western Kansas. *Groundwater*, 51(2):180–190, 2013.
- J.J. Butler, G.C. Bohling, D.O. Whittemore, and B.B. Wilson. Charting pathways toward sustainability for aquifers supporting irrigated agriculture. *Water Resources Research*, 56(10):e2020WR027961, 2020.
- F. Chiang, O. Mazdiasni, and A. AghaKouchak. Evidence of anthropogenic impacts on global drought frequency, duration, and intensity. *Nature communications*, 12(1):2754, 2021.
- A. Closas and F. Molle. Chronicle of a demise foretold: State vs. local groundwater management in Texas and the High Plains Aquifer system. *Water Alternatives*, 11(3), 2018.
- B.I. Cook, J.S. Mankin, K. Marvel, A.P. Williams, J.E. Smerdon, and K.J. Anchukaitis. Twenty-first century drought projections in the CMIP6 forcing scenarios. *Earth’s Future*, 8(6):e2019EF001461, 2020.
- B.I. Cook, A.P. Williams, and K. Marvel. Projected changes in early summer ridging and drought over the Central Plains. *Environmental Research Letters*, 17(10):104020, 2022.
- K.A. Cotterman, A.D. Kendall, B. Basso, and D.W. Hyndman. Groundwater depletion and climate change: future prospects of crop production in the central high plains aquifer. *Climatic change*, 146(1):187–200, 2018.

- A. Crane-Droesch. Machine learning methods for crop yield prediction and climate change impact assessment in agriculture. *Environmental Research Letters*, 13(11):114003, 2018.
- K.F Davis, A. Chhatre, N.D Rao, D. Singh, and R. DeFries. Sensitivity of grain yields to historical climate variability in India. *Environmental Research Letters*, 14(6):064013, 2019.
- J.M. Deines, A.D. Kendall, J.J. Butler, and D.W. Hyndman. Quantifying irrigation adaptation strategies in response to stakeholder-driven groundwater management in the US High Plains Aquifer. *Environmental Research Letters*, 14(4):044014, 2019.
- J.M. Deines, M.E. Schipanski, B. Golden, S.C. Zipper, S. Nozari, C. Rottler, B. Guerrero, and V. Sharda. Transitions from irrigated to dryland agriculture in the Ogallala Aquifer: Land use suitability and regional economic impacts. *Agricultural Water Management*, 233:106061, 2020.
- J.M. Deines, R. Patel, S. Liang, W. Dado, and D.B. Lobell. A million kernels of truth: Insights into scalable satellite maize yield mapping and yield gap analysis from an extensive ground dataset in the US Corn Belt. *Remote Sensing of Environment*, 253:112174, 2021.
- P. Döll, H. Müller Schmied, C. Schuh, F.T. Portmann, and A. Eicker. Global-scale assessment of groundwater depletion and related groundwater abstractions: Combining hydrological modeling with information from well observations and grace satellites. *Water Resources Research*, 50(7):5698–5720, 2014.
- M. Dowle and A. Srinivasan. *data.table: Extension of ‘data.frame’*, 2022. URL <https://CRAN.R-project.org/package=data.table>. R package version 1.14.4.
- J.I.R. Edreira, S. Mourtzinis, G. Azzari, J.F. Andrade, S.P. Conley, J.E. Specht, and P. Grassini. Combining field-level data and remote sensing to understand impact of management practices on producer yields. *Field Crops Research*, 257:107932, 2020.
- P. Edzer. Simple Features for R: Standardized Support for Spatial Vector Data. *The R Journal*, 10(1): 439–446, 2018.
- A.S. Elshall, A.D. Arik, A.I. El-Kadi, S. Pierce, M. Ye, K.M. Burnett, C.A. Wada, L.L. Bremer, and G. Chun. Groundwater sustainability: A review of the interactions between science and policy. *Environmental Research Letters*, 15(9):093004, 2020.
- J.S. Famiglietti. The global groundwater crisis. *Nature Climate Change*, 4(11):945–948, 2014.
- J.S. Famiglietti, M. Lo, S.L. Ho, J. Bethune, K.J. Anderson, T.H. Syed, S.C. Swenson, C.R. de Linage, and M. Rodell. Satellites measure recent rates of groundwater depletion in california’s central valley. *Geophysical Research Letters*, 38(3), 2011.
- W. Feng, CK Shum, M. Zhong, and Y. Pan. Groundwater storage changes in china from satellite gravity: An overview. *Remote Sensing*, 10(5):674, 2018.
- E.P. Fenichel, J.K. Abbott, J. Bayham, W. Boone, E. Haacker, and L. Pfeiffer. Measuring the value of groundwater and other forms of natural capital. *Proceedings of the National Academy of Sciences*, 113(9): 2382–2387, 2016.
- M. Flörke, C. Schneider, and R.I. McDonald. Water competition between cities and agriculture driven by climate change and urban growth. *Nature Sustainability*, 1(1):51–58, 2018.
- T. Foster, N. Brozović, and A.P. Butler. Modeling irrigation behavior in groundwater systems. *Water resources research*, 50(8):6370–6389, 2014.
- T. Foster, N. Brozović, and A.P. Butler. Analysis of the impacts of well yield and groundwater depth on irrigated agriculture. *Journal of Hydrology*, 523:86–96, 2015a.
- T. Foster, N. Brozović, and A.P. Butler. Why well yield matters for managing agricultural drought risk. *Weather and Climate Extremes*, 10:11–19, 2015b.
- T. Foster, N. Brozović, and A.P. Butler. Effects of initial aquifer conditions on economic benefits from groundwater conservation. *Water Resources Research*, 53(1):744–762, 2017.

- T. Foster, I.Z. Gonçalves, I. Campos, C.M.U. Neale, and N. Brozović. Assessing landscape scale heterogeneity in irrigation water use with remote sensing and in situ monitoring. *Environmental Research Letters*, 14(2):024004, 2019.
- J. Gibson, T.E. Franz, T. Wang, J. Gates, P. Grassini, H. Yang, and D. Eisenhauer. A case study of field-scale maize irrigation patterns in western Nebraska: implications for water managers and recommendations for hyper-resolution land surface modeling. *Hydrology and Earth System Sciences*, 21(2):1051–1062, 2017.
- K.E.B. Gibson, J.P. Gibson, and P. Grassini. Benchmarking irrigation water use in producer fields in the US central Great Plains. *Environmental Research Letters*, 14(5):054009, 2019.
- T.J. Glose, S. Zipper, D.W. Hyndman, A.D. Kendall, J.M. Deines, and J.J. Butler. Quantifying the impact of lagged hydrological responses on the effectiveness of groundwater conservation. *Water Resources Research*, 58(7):e2022WR032295, 2022.
- R.Q. Grafton, J. Williams, C.J. Perry, F. Molle, C. Ringler, P. Steduto, B. Udall, S.A. Wheeler, Y. Wang, D. Garrick, et al. The paradox of irrigation efficiency. *Science*, 361(6404):748–750, 2018.
- E. Haacker, A.D. Kendall, and D.W. Hyndman. Water level declines in the high plains aquifer: Predevelopment to resource senescence. *Groundwater*, 54(2):231–242, 2016.
- E. Haacker, A.D. Kendall, and D.W. Hyndman. High plains/ogallala water table elevations annual estimates. *HydroShare*, 2023.
- G.R. Hecox, P.A. Macfarlane, and B.B. Wilson. Calculation of yield for high plains wells: Relationship between saturated thickness and well yield. Technical report, Kansas Geological Survey, 2002.
- R.A. Hrozencik, D.T. Manning, J.F. Suter, C. Goemans, and R.T. Bailey. The heterogeneous impacts of groundwater management policies in the republican river basin of colorado. *Water Resources Research*, 53(12):10757–10778, 2017.
- D.R. Hyslop and G.W. Imbens. Bias from classical and other forms of measurement error. *Journal of Business & Economic Statistics*, 19(4):475–481, 2001.
- M. Jain, R. Fishman, P. Mondal, G.L. Galford, N. Bhattarai, S. Naeem, U. Lall, Balwinder-Singh, and R.S. DeFries. Groundwater depletion will reduce cropping intensity in India. *Science Advances*, 7(9):eabd2849, 2021.
- M.T Kahil, A. Dinar, and J. Albiac. Modeling water scarcity and droughts for policy adaptation to climate change in arid and semiarid regions. *Journal of Hydrology*, 522:95–109, 2015.
- Leonard F Konikow. Long-term groundwater depletion in the united states. *Groundwater*, 53(1):2–9, 2015.
- Leonard F Konikow and Eloise Kendy. Groundwater depletion: A global problem. *Hydrogeology Journal*, 13(1):317–320, 2005.
- J.T. Korus and H.J. Hensen. Depletion percentage and nonlinear transmissivity as design criteria for groundwater-level observation networks. *Environmental Earth Sciences*, 79(16):382, 2020.
- P. Kreins, M. Henseler, J. Anter, F. Herrmann, and F. Wendland. Quantification of climate change impact on regional agricultural irrigation and groundwater demand. *Water Resources Management*, 29(10):3585–3600, 2015.
- Y. Kuwayama, A. Thompson, R. Bernknopf, B. Zaitchik, and P. Vail. Estimating the impact of drought on agriculture using the US Drought Monitor. *American Journal of Agricultural Economics*, 101(1):193–210, 2019.
- X. Li and T.J. Troy. Changes in rainfed and irrigated crop yield response to climate in the western US. *Environmental Research Letters*, 13(6):064031, 2018.
- Y. Li, K. Guan, G.D. Schnitkey, E. DeLucia, and B. Peng. Excessive rainfall leads to maize yield loss of a comparable magnitude to extreme drought in the united states. *Global Change Biology*, 25(7):2325–2337, 2019.

- Y. Li, K. Guan, B. Peng, T.E. Franz, B. Wardlow, and M. Pan. Quantifying irrigation cooling benefits to maize yield in the us midwest. *Global Change Biology*, 26(5):3065–3078, 2020.
- B. Lindblad. *tidyUSDA: A Minimal Tool Set for Gathering USDA Quick Stat Data for Analysis and Visualization*, 2022. URL <https://bradlindblad.github.io/tidyUSDA/>. R package version 0.4.0.
- D.B. Lobell and C. Bonfils. The effect of irrigation on regional temperatures: A spatial and temporal analysis of trends in california, 1934–2002. *Journal of Climate*, 21(10):2063–2071, 2008.
- D.B. Lobell, M.J. Roberts, W. Schlenker, N. Braun, B.B. Little, R.M. Rejesus, and G.L. Hammer. Greater sensitivity to drought accompanies maize yield increase in the US Midwest. *Science*, 344(6183):516–519, 2014.
- J. Lu, G.J. Carbone, X. Huang, K. Lackstrom, and P. Gao. Mapping the sensitivity of agriculture to drought and estimating the effect of irrigation in the United States, 1950–2016. *Agricultural and Forest Meteorology*, 292:108124, 2020.
- W. Lu, W. Adamowicz, S.R. Jeffrey, G.G. Goss, and M. Faramarzi. Crop yield response to climate variables on dryland versus irrigated lands. *Canadian Journal of Agricultural Economics/Revue canadienne d’agroeconomie*, 66(2):283–303, 2018.
- X. Luan, R. Bommarco, A. Scaini, and G. Vico. Combined heat and drought suppress rainfed maize and soybean yields and modify irrigation benefits in the USA. *Environmental research letters*, 16(6):064023, 2021.
- M. Lubell, W. Blomquist, and L. Beutler. Sustainable groundwater management in California: A grand experiment in environmental governance. *Society & Natural Resources*, 33(12):1447–1467, 2020.
- D. MacEwan, M. Cayar, A. Taghavi, D. Mitchell, S. Hatchett, and R. Howitt. Hydroeconomic modeling of sustainable groundwater management. *Water Resources Research*, 53(3):2384–2403, 2017.
- D.T. Manning, C. Goemans, and A. Maas. Producer responses to surface water availability and implications for climate change adaptation. *Land Economics*, 93(4):631–653, 2017.
- L.T. Marston, S. Zipper, S.M. Smith, J.J. Allen, J.J. Butler, S. Gautam, and J.Y. David. The importance of fit in groundwater self-governance. *Environmental Research Letters*, 17(11):111001, 2022.
- T. Mieno and N. Brozović. Price elasticity of groundwater demand: attenuation and amplification bias due to incomplete information. *American Journal of Agricultural Economics*, 99(2):401–426, 2017.
- A. Mrad, G.G. Katul, D.F. Levia, A.J. Guswa, E.W. Boyer, M. Bruen, D.E. Carlyle-Moses, R. Coyte, I.F. Creed, N. Van De Giesen, et al. Peak grain forecasts for the us high plains amid withering waters. *Proceedings of the National Academy of Sciences*, 117(42):26145–26150, 2020.
- E.D. Mullens and R.A. McPherson. Quantitative scenarios for future hydrologic extremes in the US Southern Great Plains. *International Journal of Climatology*, 39(5):2659–2676, 2019.
- A. Ortiz-Bobea, H. Wang, C.M. Carrillo, and T.R. Ault. Unpacking the climatic drivers of US agricultural yields. *Environmental Research Letters*, 14(6):064003, 2019.
- L.S. Pereira, R.G. Allen, M. Smith, and D. Raes. Crop evapotranspiration estimation with fao56: Past and future. *Agricultural Water Management*, 147:4–20, 2015.
- C.D. Pérez-Blanco, A. Loch, F. Ward, C. Perry, and D. Adamson. Agricultural water saving through technologies: a zombie idea. *Environmental Research Letters*, 16(11):114032, 2021.
- R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2021. URL <https://www.R-project.org/>.
- M. Rouhi-Rad, A. Araya, and Z.T. Zambreski. Downside risk of aquifer depletion. *Irrigation Science*, 38: 577–591, 2020a.

- M. Rouhi-Rad, N. Brozović, T. Foster, and T. Mieno. Effects of instantaneous groundwater availability on irrigated agriculture and implications for aquifer management. *Resource and Energy Economics*, 59:101129, 2020b.
- M. Rouhi-Rad, E.M.K. Haacker, V. Sharda, S. Nozari, Z. Xiang, A. Araya, V. Uddameri, J.F. Suter, and P. Gowda. MOD\$AT: A hydro-economic modeling framework for aquifer management in irrigated agricultural regions. *Agricultural Water Management*, 238:106194, 2020c.
- P.J. Ruess, M. Konar, N. Wanders, and M. Bierkens. Irrigation by crop in the Continental United States from 2008 to 2020. *Water Resources Research*, page e2022WR032804, 2022.
- B.R. Scanlon, C.C. Faunt, L. Longuevergne, R.C. Reedy, W.M. Alley, V.L. McGuire, and P.B. McMahon. Groundwater depletion and sustainability of irrigation in the us high plains and central valley. *Proceedings of the national academy of sciences*, 109(24):9320–9325, 2012.
- B.R. Scanlon, S. Fakhreddine, A. Rateb, I. de Graaf, J. Famiglietti, T. Gleeson, R.Q. Grafton, E. Jobbagy, S. Kebede, S.R. Kolusu, et al. Global water resources and the role of groundwater in a resilient water future. *Nature Reviews Earth & Environment*, pages 1–15, 2023.
- W. Schlenker and D.B. Lobell. Robust negative impacts of climate change on African agriculture. *Environmental Research Letters*, 5(1):014010, 2010.
- W. Schlenker and M.J. Roberts. Nonlinear temperature effects indicate severe damages to us crop yields under climate change. *Proceedings of the National Academy of sciences*, 106(37):15594–15598, 2009.
- S.J. Smidt, E.M.K. Haacker, A.D. Kendall, J.M. Deines, L. Pei, K.A. Cotterman, H. Li, X. Liu, B. Basso, and D.W. Hyndman. Complex water management in modern agriculture: Trends in the water-energy-food nexus over the High Plains Aquifer. *Science of the Total Environment*, 566:988–1001, 2016.
- J.F. Suter, M. Rouhi-Rad, D.T. Manning, C. Goemans, and M.R. Sanderson. Depletion, climate, and the incremental value of groundwater. *Resource and Energy Economics*, 63:101143, 2021.
- A.M. Ukkola, M.G. De Kauwe, M.L. Roderick, G. Abramowitz, and A.J. Pitman. Robust future changes in meteorological drought in CMIP6 projections despite uncertainty in precipitation. *Geophysical Research Letters*, 47(11):e2020GL087820, 2020.
- M. Venkatappa, N. Sasaki, P. Han, and I. Abe. Impacts of droughts and floods on croplands and crop production in Southeast Asia—An application of Google Earth Engine. *Science of the Total Environment*, 795:148829, 2021.
- Y. Wada and M.F. Bierkens. Sustainability of global water use: past reconstruction and future projections. *Environmental Research Letters*, 9(10):104003, 2014.
- Y. Wada, L.P.H. Van Beek, C.M. Van Kempen, J.W.T.M. Reckman, S. Vasak, and M.F.P. Bierkens. Global depletion of groundwater resources. *Geophysical research letters*, 37(20), 2010.
- Y. Wada, D. Wisser, S. Eisner, M. Flörke, D. Gerten, I. Haddeland, N. Hanasaki, Y. Masaki, F.T. Portmann, T. Stacke, et al. Multimodel projections and uncertainties of irrigation water demand under climate change. *Geophysical Research Letters*, 40(17):4626–4632, 2013.
- H. Wickham. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, 2016. ISBN 978-3-319-24277-4. URL <https://ggplot2.tidyverse.org>.
- J. Yoon, C. Klassert, P. Selby, T. Lachaut, S. Knox, N. Avisse, J.J. Harou, A. Tilmant, B. Klauer, D. Mustafa, et al. A coupled human–natural system analysis of freshwater security under climate and population change. *Proceedings of the National Academy of Sciences*, 118(14):e2020431118, 2021.
- W. Zhou, K. Guan, N. Peng, J. Shi, C. Jiang, B. Wardlow, M. Pan, J.S. Kimball, T.E. Franz, P. Gentine, et al. Connections between the hydrological cycle and crop yield in the rainfed US Corn Belt. *Journal of Hydrology*, 590:125398, 2020.

- 641 Y. Zhou, F. Zwahlen, Y. Wang, and Y. Li. Impact of climate change on irrigation requirements in terms of
642 groundwater resources. *Hydrogeology journal*, 18(7):1571–1582, 2010.
- 643 P. Zhu and J. Burney. Untangling irrigation effects on maize water and heat stress alleviation using satellite
644 data. *Hydrology and Earth System Sciences*, 26(3):827–840, 2022.
- 645 P. Zhu, Q. Zhuang, S.V. Archontoulis, C. Bernacchi, and C. Müller. Dissecting the nonlinear response of maize
646 yield to high temperature stress with model-data integration. *Global Change Biology*, 25(7):2470–2484,
647 2019.
- 648 P. Zhu, J. Burney, J. Chang, Z. Jin, N.D. Mueller, Q. Xin, J. Xu, L. Yu, D. Makowski, and P. Ciais. Warming
649 reduces global agricultural production by decreasing cropping frequency and yields. *Nature Climate Change*,
650 12(11):1016–1023, 2022.
- 651 S.C. Zipper, J. Qiu, and C.J. Kucharik. Drought effects on US maize and soybean production: spatiotemporal
652 patterns and historical changes. *Environmental Research Letters*, 11(9):094021, 2016.