

Calculus I

Linearization and differentials

Todor Milev

2019

Outline

1 Linear Approximations

Outline

1 Linear Approximations

2 Differentials

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Linear Approximations and Differentials

- Main idea: A curve is very close to its tangent line at the point of tangency.
- We can use the tangent line at $(a, f(a))$ as an approximation to the curve $y = f(x)$.
- This approximation works well as long as x is near a .



Definition (Linearization of f at a)

The linear function whose graph is the tangent line at $(a, f(a))$ is called the linearization of f at a . Its equation is

$$L(x) = f(a) + f'(a)(x - a).$$

Definition (Linear Approximation of $f(x)$ near a)

The approximation

$$f(x) \approx f(a) + f'(a)(x - a)$$


is called the linear approximation of f at a .

Let $y = f(x)$, $\Delta y := f(x) - f(a)$, and $\Delta x := x - a$.

Definition (Linear approx. $y = f(x)$ near a , alternative notation)


$$\Delta y \approx \frac{dy}{dx} \Delta x \quad .$$

Linear approximations




Function	f	L
Run	Δx	Δx
Rise	Δf	ΔL
Formula	$\Delta f = f(x + \Delta x) - f(x)$	$\Delta L = (\Delta x)f'(x)$

Linear approximations



Function	f	L
Run	Δx	Δx
Rise	Δf	ΔL
Formula	$\Delta f = f(x + \Delta x) - f(x)$	$\Delta L = (\Delta x)f'(x)$


Linear approximations



Function	f	L
Run	Δx	Δx
Rise	Δf	ΔL
Formula	$\Delta f = f(x + \Delta x) - f(x)$	$\Delta L = (\Delta x)f'(x)$

Example


Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?



Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?


- $f'(x) = ?$
- $f(1) = ?$
- $f'(1) = ?$
- Linearization:



Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = ?$
- $f(1) = ?$
- $f'(1) = ?$
- Linearization:



Example


Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.

- $f(1) = ?$

- $f'(1) = ?$


- Linearization:



Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?


- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = ?$
- $f'(1) = ?$
- Linearization:



Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?


- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = ?$
- Linearization:



Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?


- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = ?$
- Linearization:



Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$L(x) = ? + ? (x - ?)$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$L(x) = ? + ? (x - 1)$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$L(x) = \textcolor{red}{?} + \textcolor{red}{?} (x - 1)$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$L(x) = 2 + ? (x - 1)$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$L(x) = 2 + ? (x - 1)$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$L(x) = 2 + \frac{1}{4}(x - 1)$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$




Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$




- $\sqrt{3.98} = f(0.98) \approx ?$
- $\sqrt{4.05} = f(1.05) \approx ?$

Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$




- $\sqrt{3.98} = f(0.98) \approx ?$
- $\sqrt{4.05} = f(1.05) \approx ?$

Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$




- $\sqrt{3.98} = f(0.98) \approx \frac{7}{4} + \frac{0.98}{4} = 1.995$.
- $\sqrt{4.05} = f(1.05) \approx ?$

Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$




- $\sqrt{3.98} = f(0.98) \approx \frac{7}{4} + \frac{0.98}{4} = 1.995$.
- $\sqrt{4.05} = f(1.05) \approx ?$

Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$




- $\sqrt{3.98} = f(0.98) \approx \frac{7}{4} + \frac{0.98}{4} = 1.995$.
- $\sqrt{4.05} = f(1.05) \approx \frac{7}{4} + \frac{1.05}{4} = 2.0125$.

Example

Find the linearization of the function $f(x) = \sqrt{x+3}$ at $a = 1$ and use it to approximate the numbers $\sqrt{3.98}$ and $\sqrt{4.05}$. Are these approximations overestimates or underestimates?

- $f'(x) = \frac{1}{2\sqrt{x+3}}$.
- $f(1) = \sqrt{1+3} = 2$.
- $f'(1) = \frac{1}{2\sqrt{1+3}} = \frac{1}{4}$.
- Linearization:

$$\begin{aligned} L(x) &= 2 + \frac{1}{4}(x - 1) \\ &= \frac{7}{4} + \frac{x}{4} \end{aligned}$$



The graph of the linearization is above the curve, so these are overestimates.

- $\sqrt{3.98} = f(0.98) \approx \frac{7}{4} + \frac{0.98}{4} = 1.995$.
- $\sqrt{4.05} = f(1.05) \approx \frac{7}{4} + \frac{1.05}{4} = 2.0125$.

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) =$
- $f(2.05) =$
- $\Delta y =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) =$
- $f(2.05) =$
- $\Delta y =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) =$
- $\Delta y =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$

- $f(2.05) =$

- $\Delta y =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) =$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) =$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) = 3x^2 + 2x - 2.$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x = (3x^2 + 2x - 2)\Delta x.$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) = 3x^2 + 2x - 2.$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x = (3x^2 + 2x - 2)\Delta x.$
- When $x = 2$ and $\Delta x = 0.05$, we have:
- $\Delta L =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) = 3x^2 + 2x - 2.$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x = (3x^2 + 2x - 2)\Delta x.$
- When $x = 2$ and $\Delta x = 0.05$, we have:
- $\Delta L = (3(2)^2 + 2(2) - 2)(0.05) =$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) = 3x^2 + 2x - 2.$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x = (3x^2 + 2x - 2)\Delta x.$
- When $x = 2$ and $\Delta x = 0.05$, we have:
- $\Delta L = (3(2)^2 + 2(2) - 2)(0.05) = 0.7.$

Example

Compute Δy and $\Delta L = f'(x)\Delta x$ if $y = f(x) = x^3 + x^2 - 2x + 1$ and x changes from 2 to 2.05.

- $f(2) = 2^3 + 2^2 - 2(2) + 1 = 9.$
- $f(2.05) = (2.05)^3 + (2.05)^2 - 2(2.05) + 1 = 9.717625.$
- $\Delta y = f(2.05) - f(2) = 9.717625 - 9 = 0.717625.$
- $f'(x) = 3x^2 + 2x - 2.$
- $\Delta y \simeq \Delta L = f'(x)\Delta x = f'(x)\Delta x = (3x^2 + 2x - 2)\Delta x.$
- When $x = 2$ and $\Delta x = 0.05$, we have:
- $\Delta L = (3(2)^2 + 2(2) - 2)(0.05) = 0.7.$
- Therefore $\Delta L \approx \Delta y = 0.7$, an approximation of $\Delta y = 0.717625.$

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $\Delta y \approx \frac{dy}{dx} \Delta x$

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $\Delta y \approx \frac{dy}{dx} \Delta x$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy \approx \frac{dy}{dx} \Delta x$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a **formal identity**.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.
- Define the *differential* d and the *differential forms* $dx, d(f(x))$ by requesting that d and dx satisfy the transformation law

$$d(f(x)) = f'(x)dx$$

for any differentiable function $f(x)$. In abbreviated notation:

$$df = f'dx$$

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.
- Define the *differential* d and the *differential forms* $dx, d(f(x))$ by requesting that d and dx satisfy the transformation law

$$d(f(x)) = f'(x)dx$$

for any differentiable function $f(x)$. In abbreviated notation:

$$df = f' dx$$

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.
- Define the *differential* d and the *differential forms* $dx, d(f(x))$ by requesting that **d and dx satisfy the transformation law**

$$d(f(x)) = f'(x)dx$$

for any differentiable function $f(x)$. In abbreviated notation:

$$df = f'dx$$

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.
- Define the *differential* d and the *differential forms* $dx, d(f(x))$ by requesting that d and dx satisfy the transformation law

$$d(f(x)) = f'(x)dx$$

for any differentiable function $f(x)$. In abbreviated notation:

$$df = f'dx$$

Expressions containing expression of the form $d(\text{something})$ are called *differential forms*.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.
- Define the *differential d* and the *differential forms* $dx, d(f(x))$ by requesting that d and dx satisfy the transformation law

$$d(f(x)) = f'(x)dx$$

for any differentiable function $f(x)$. In abbreviated notation:

$$df = f'dx$$

Expressions containing expression of the form $d(\text{something})$ are called differential forms.

Differentials

- Recall $\Delta y, \Delta x$ stand for change of x, y . Recall: $\Delta y \approx \frac{dy}{dx} \Delta x$
- $dy = \frac{dy}{dx} dx = dy$
- If we substitute Δy by the formal expression dy and Δx by the formal expression dx , the expression dx appears to “cancel” to give a formal identity.
- Define the *differential* d and the *differential forms* $dx, d(f(x))$ by requesting that d and dx satisfy the transformation law

$$d(f(x)) = f'(x)dx$$

for any differentiable function $f(x)$. In abbreviated notation:

$$df = f'dx$$

Expressions containing expression of the form $d(\text{something})$ are called differential forms.

- $df(x) = f'(x)dx.$

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are **formal expressions related by a transformation law**.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.
- Courses such as “Integration and Manifolds” or “Differential geometry” usually give precise definitions and fill in the details.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are **formal expressions related by a transformation law**.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.
- Courses such as “Integration and Manifolds” or “Differential geometry” usually give precise definitions and fill in the details.
- Nonetheless, **what we studied** is completely sufficient for practical purposes and carrying out computations.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.
- Courses such as “Integration and Manifolds” or “Differential geometry” usually give precise definitions and fill in the details.
- Nonetheless, what we studied is **completely sufficient** for practical purposes and **carrying out computations**.

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.
- Courses such as “Integration and Manifolds” or “Differential geometry” usually give precise definitions and fill in the details.
- Nonetheless, what we studied is completely sufficient for practical purposes and carrying out computations.
- **Do not confuse differentials with derivatives.**

$$df(x) = f'(x)$$

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.
- Courses such as “Integration and Manifolds” or “Differential geometry” usually give precise definitions and fill in the details.
- Nonetheless, what we studied is completely sufficient for practical purposes and carrying out computations.
- **Do not confuse differentials with derivatives.**



~~$df(x) = f'(x)dx$~~

- $df(x) = f'(x)dx$.
- On the previous slide we stated the differential d and the differential forms $dx, df(x)$ are formal expressions related by a transformation law.
- The precise definitions of differential forms and differentials are outside of the scope of Calculus I and II.
- Differential forms “encode” linear approximations which in turn “encode” “infinitesimal” lengths of segments.
- Courses such as “Integration and Manifolds” or “Differential geometry” usually give precise definitions and fill in the details.
- Nonetheless, what we studied is completely sufficient for practical purposes and carrying out computations.
- **Do not confuse differentials with derivatives.** The correct equality is this.

~~$$df(x) = f'(x)$$~~

$$df(x) = f'(x)dx$$

Example

Compute the differential (via dx).

$$d(x^2)$$

Example

Compute the differential (via dx).

$$d(x^2) = (x^2)' dx$$

Example

Compute the differential (via dx).

$$d(x^2) = (x^2)' dx = 2x dx \quad .$$

Example

Compute the differential (via dx).

$$d(\sqrt{x})$$

Example

Compute the differential (via dx).

$$d(\sqrt{x}) = (\sqrt{x})' dx$$

Example

Compute the differential (via dx).

$$d(\sqrt{x}) = (\sqrt{x})' dx = \frac{1}{2\sqrt{x}} dx .$$

- All rules for computing with derivatives have analogues for computing with differential forms.

- All rules for computing with derivatives have analogues for computing with differential forms.
- The rules for computing differential forms are a direct consequence of the corresponding derivative rules and the transformation law $d(f(x)) = f'(x)dx$.

Rule name: **product rule.**

Differential rule

Derivative rule

$$(fg)' = f'g + fg'$$

Rule name: **product rule.**

Differential rule

$$d(fg) = gdf + fdg$$

Derivative rule

$$(fg)' = f'g + fg'$$

Rule name: **constant derivative rule.**

Differential rule

$$d(fg) = gdf + fdg$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

c-const.

Rule name: **constant derivative rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

c-const.

c-const.

Rule name: **sum rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

c-const.

c-const.

Rule name: **sum rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

c-const.

c-const.

Rule name: **chain rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

c-const.

c-const.

$$(f(g(x)))' = f'(g(x))g'(x)$$

Rule name: **chain rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

c-const.

c-const.

Rule name: **power rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

c-const.

c-const.

Rule name: **power rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

c-const.

c-const.

Rule name: **exponent derivative rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

c-const.

c-const.

Rule name: **exponent derivative rule.**

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

$$(\sin x)' = \cos x$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

$$d(\sin x) = \cos x dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

$$(\sin x)' = \cos x$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

$$d(\sin x) = \cos x dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

$$d(\sin x) = \cos x dx$$

$$d(\cos x) = -\sin x dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

$$d(\sin x) = \cos x dx$$

$$d(\cos x) = -\sin x dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

$$(\ln x)' = \frac{1}{x}$$

c-const.

c-const.

Rule name:

Differential rule

$$d(fg) = gdf + fdg$$

$$dc = 0 = 0dx$$

$$d(cf) = cdf$$

$$d(f + g) = df + dg$$

$$\begin{aligned} df(g(x)) &= f'(g(x))dg(x) \\ &= f'(g(x))g'(x)dx \end{aligned}$$

$$df(g) = f'(g)dg$$

$$d(x^n) = nx^{n-1}dx$$

$$d(e^x) = e^x dx$$

$$d(\sin x) = \cos x dx$$

$$d(\cos x) = -\sin x dx$$

$$(d \ln x) = \frac{1}{x} dx$$

Derivative rule

$$(fg)' = f'g + fg'$$

$$(c)' = 0$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

c-const.

c-const.

$$(x^n)' = nx^{n-1}$$

$$(e^x)' = e^x$$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

$$(\ln x)' = \frac{1}{x}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

$$d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$.

$$d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) = d(\ln u)$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$.

$$d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) = d(\ln u) = ? du$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$.

$$d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) = d(\ln u) = \frac{1}{u}du$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$.

$$d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) = d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) =$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$.

$$\begin{aligned}d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) = \\&= \frac{1}{u}d\left(\sqrt{1 + x^2}\right) =\end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) = \\&= \frac{1}{u}d\left(\sqrt{1 + x^2}\right) = \frac{1}{u}d\left(v^{\frac{1}{2}}\right) =\end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) = \\&= \frac{1}{u}d\left(\sqrt{1 + x^2}\right) = \frac{1}{u}d\left(v^{\frac{1}{2}}\right) = \frac{1}{u}\frac{1}{2}v^{-\frac{1}{2}}dv\end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned} d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u} du = \frac{1}{u} d\left(1 + \sqrt{1 + x^2}\right) = \\ &= \frac{1}{u} d\left(\sqrt{1 + x^2}\right) = \frac{1}{u} d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2} v^{-\frac{1}{2}} dv \\ &= \frac{1}{2uv^{\frac{1}{2}}} d\left(1 + x^2\right) = \end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) = \\&= \frac{1}{u}d\left(\sqrt{1 + x^2}\right) = \frac{1}{u}d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2}v^{-\frac{1}{2}}dv \\&= \frac{1}{2uv^{\frac{1}{2}}}d\left(1 + x^2\right) =\end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) = \\&= \frac{1}{u}d\left(\sqrt{1 + x^2}\right) = \frac{1}{u}d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2} v^{-\frac{1}{2}} dv \\&= \frac{1}{2uv^{\frac{1}{2}}} d\left(1 + x^2\right) = \frac{2x}{2uv^{\frac{1}{2}}} dx =\end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}
 d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u}du = \frac{1}{u}d\left(1 + \sqrt{1 + x^2}\right) = \\
 &= \frac{1}{u}d\left(\sqrt{1 + x^2}\right) = \frac{1}{u}d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2} v^{-\frac{1}{2}} dv \\
 &= \frac{1}{2uv^{\frac{1}{2}}}d\left(1 + x^2\right) = \frac{\cancel{2}x}{\cancel{2}uv^{\frac{1}{2}}}dx = \frac{x}{uv^{\frac{1}{2}}}dx
 \end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}
 d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u} du = \frac{1}{u} d\left(1 + \sqrt{1 + x^2}\right) = \\
 &= \frac{1}{u} d\left(\sqrt{1 + x^2}\right) = \frac{1}{u} d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2} v^{-\frac{1}{2}} dv \\
 &= \frac{1}{2uv^{\frac{1}{2}}} d\left(1 + x^2\right) = \frac{2x}{2uv^{\frac{1}{2}}} dx = \frac{x}{uv^{\frac{1}{2}}} dx \\
 &= \frac{x}{\left(1 + \sqrt{1 + x^2}\right) \sqrt{1 + x^2}} dx
 \end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}
 d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u} du = \frac{1}{u} d\left(1 + \sqrt{1 + x^2}\right) = \\
 &= \frac{1}{u} d\left(\sqrt{1 + x^2}\right) = \frac{1}{u} d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2} v^{-\frac{1}{2}} dv \\
 &= \frac{1}{2uv^{\frac{1}{2}}} d\left(1 + x^2\right) = \frac{2x}{2uv^{\frac{1}{2}}} dx = \frac{x}{u\sqrt{1 + x^2}} dx \\
 &= \frac{x}{\left(1 + \sqrt{1 + x^2}\right) \sqrt{1 + x^2}} dx
 \end{aligned}$$

Differentials are especially efficient at “encoding” the chain rule.

Example

Compute the differential $d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right)$.

Set $u = 1 + \sqrt{1 + x^2}$. Set $v = 1 + x^2$.

$$\begin{aligned}
 d\left(\ln\left(1 + \sqrt{1 + x^2}\right)\right) &= d(\ln u) = \frac{1}{u} du = \frac{1}{u} d\left(1 + \sqrt{1 + x^2}\right) = \\
 &= \frac{1}{u} d\left(\sqrt{1 + x^2}\right) = \frac{1}{u} d\left(v^{\frac{1}{2}}\right) = \frac{1}{u} \frac{1}{2} v^{-\frac{1}{2}} dv \\
 &= \frac{1}{2uv^{\frac{1}{2}}} d\left(1 + x^2\right) = \frac{2x}{2uv^{\frac{1}{2}}} dx = \frac{x}{uv^{\frac{1}{2}}} dx \\
 &= \frac{x}{\left(1 + \sqrt{1 + x^2}\right) \sqrt{1 + x^2}} dx
 \end{aligned}$$