#### **Precalculus**

# Equations involving logarithms and exponents

**Todor Miley** 

2019

# **Outline**

- Equations involving logarithms
- Equations involving exponents
- 3 Inverse function problems and exponents
- Basic exponential inequalities

### License to use and redistribute

These lecture slides and their LATEX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share to copy, distribute and transmit the work,
- to Remix to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides: https://github.com/tmilev/freecalc
- Should the link be outdated/moved, search for "freecalc project".
- Creative Commons license CC BY 3.0:
   https://creativecommons.org/licenses/by/3.0/us/and the links therein.

$$\log_3(2x^2+1) = 2$$
 | Exponentiate base 3  $3^{\log_3(2x^2+1)} = 3^2$   $2x^2+1 = 9$   $2x^2 = 8$   $x^2 = \frac{8}{2} = 4$   $x = \pm\sqrt{4} = \pm 2$   $x = 2$  or  $x = -2$  | final answer

The logarithmic property  $\log_a(xy) = \log_a x + \log_a y$  holds only for positive x, y. Failure to check the positivity of x, y can result in extraneous (fake) solutions to logarithmic equations.

### Example

$$\log_2(x+2) + \log_2(x-1) = 2$$
 Domain:  $x > 1$   
 $\log_2((x+2)(x-1)) = 2$  Exponentiate base 2  
 $(x+2)(x-1) = 2^2$   
 $x^2 + x - 2 = 4$   
 $x^2 + x - 6 = 0$   
 $(x-2)(x+3) = 0$   
 $x = 2$  or  $x = 3$   
 $x = -3$  not a solution (outside of domain)

# Example (Solve exponential equation without logarithms)

Solve for t.

$$e^{x-3}=2e^{2x-1}$$
 Divide by  $e^{2x-1}$ 
 $\frac{e^{x-3}}{e^{2x-1}}=2$ 
 $e^{x-3-(2x-1)}=2$ 
 $e^{-x-2}=2$  Apply In
 $-x-2=\ln 2$ 
 $-x=\ln 2+2$ 
 $x=-(\ln 2+2)$ 
 $x=-\ln 2-2$  Final answer
 $x\approx -2.693$  Calculator

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1 - 5\log_2 3}{2\log_2 3 + 1}$$

$$x \approx -1.1038$$

 $x \approx -1.1038$ 

#### Common base

$$a = b^{\log_b a}$$

Apply log<sub>2</sub>

Calculator

$$e^{5-3x}=10$$
 apply In  $=$   $=$   $x=$  Calculator:  $x\approx0.8991$ .

# Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for *t*:

=

=

=

=

t =

# Example (Solving a quadratic exponential equation)

Solve for x.

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$  | Substitute  $u = 3^{x}$ 
 $u^{2} - 2u - 63 = 0$ 
 $(u - 9)(u + 7) = 0$ 
 $u = 9 \text{ or } u = -7$ 
 $3^{x} = 9 \text{ or } 3^{x} = -7$ 
 $x = 2 \text{ no real solution}$ 

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set  $e^x = u$ . Then  $e^{2x} = u^2$ .

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

$$u=4$$
 or  $u=-1$   
 $e^x=4$  or  $e^x=-1$   
 $x=\ln 4$  or no real solution  
 $x\approx 1.3863$ 

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set  $u = 2^x$ . Then  $4^{x+1} = 4u^2$ ,  $2^{x+2} = 4u$ .
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln\left(\frac{3}{2}\right)}{\ln 2} \approx 0.58496 \text{ or no real solution}$$

# Example (Exponential equation that reduces to quadratic)

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$


$$2x = \log_{3} 9$$

$$2x = 2$$

$$x = 1$$

$$>$$
 2 terms  $\Rightarrow$  transfer one side  $3^{2x} = u$   $3^{-2x} = (3^{2x})^{-1} = u^{-1}$  Multiply  $\cdot u$ 

Find 
$$f^{-1}(x)$$
 for  $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ .  
 $f = \tanh = \text{hyperbolic}$  tangent function.



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln \left(\frac{1 + y}{1 - y}\right)$$

Set 
$$u = e^x$$
  
 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$ 

Take In

# Solve the inequality.

$$2^{-3x-5} < 7$$

$$\log_2 2^{-3x-5} < \log_2 7$$

 $x \in \left(-\frac{5 + \log_2 7}{3}, \infty\right)$ 

Logarithms preserve inequalities: apply log<sub>2</sub>

Division by negative number flips inequalities

